

$\tan^2 1^\circ + \tan^2 2^\circ + \dots + \tan^2 89^\circ$

Created by Mr. Francis Hung

Last updated: 21 April 2011

To find $\tan^2 1^\circ + \tan^2 2^\circ + \dots + \tan^2 89^\circ$.

Consider the equation $\left(\frac{1+ix}{1-ix}\right)^{180} = 1$, where $i = \sqrt{-1}$

$$\frac{1+ix}{1-ix} = \cos \frac{2k\pi}{180} + i \sin \frac{2k\pi}{180}, k = 0, 1, 2, \dots, 179$$

$$\frac{1+ix}{1-ix} = \cos \frac{k\pi}{90} + i \sin \frac{k\pi}{90}$$

$$\text{Let } \omega = \cos \frac{k\pi}{90} + i \sin \frac{k\pi}{90}$$

$$\frac{1+ix}{1-ix} = \omega$$

$$1+ix = \omega(1-ix)$$

$$(1+\omega)ix = \omega - 1$$

$$x = \frac{\omega - 1}{i(\omega + 1)} = \frac{\omega^{\frac{1}{2}} - \omega^{-\frac{1}{2}}}{i(\omega^{\frac{1}{2}} + \omega^{-\frac{1}{2}})} = \frac{2i \sin \frac{k\pi}{180}}{2i \cos \frac{k\pi}{180}} = \tan \frac{k\pi}{180}, k = 0, 1, 2, \dots, 179$$

But the denominator $\neq 0$, $x \neq -i$

Also when $k = 90$, $x = \tan \frac{\pi}{2}$ which is undefined, therefore rejected.

From the equation: $\left(\frac{1+ix}{1-ix}\right)^{180} = 1 \Rightarrow (1+ix)^{180} - (1-ix)^{180} = 0$

Let $f(x) = (1+ix)^{180} - (1-ix)^{180}$, $f(x)$ can be expanded by binomial theorem

$f(x) = a_{180}x^{180} + a_{179}x^{179} + \dots + a_1x + a_0$, where $a_r = {}_{180}C_r (i)^r [1 - (-1)^r]$, $0 \leq r \leq 180$

According to the formula, $a_r = 0$ when r is even, $a_r = {}_{180}C_r \times 2(i)^r$ when r is odd

$$\therefore f(x) = a_{179}x^{179} + a_{177}x^{177} + \dots + a_3x^3 + a_1x$$

The roots of this equation are: $x = \tan \frac{k\pi}{180}, k = 0, 1, 2, \dots, 89, 91, \dots, 179$

$$\text{Let } g(x) = \frac{f(x)}{x} = a_{179}x^{178} + a_{177}x^{176} + \dots + a_3x^2 + a_1,$$

The roots of $g(x)$ are $\tan \frac{k\pi}{180}, k = 1, 2, \dots, 89, 91, \dots, 179$

$$\text{Consider } g(\sqrt{x}) = a_{179}x^{89} + a_{177}x^{88} + \dots + a_3x + a_1 = 0$$

The roots are $\tan^2 1^\circ, \tan^2 2^\circ, \dots, \tan^2 89^\circ$

$$\tan^2 1^\circ + \tan^2 2^\circ + \dots + \tan^2 89^\circ = \text{sum of roots of the equation of } g(\sqrt{x}) = 0$$

$$\tan^2 1^\circ + \tan^2 2^\circ + \dots + \tan^2 89^\circ = -\frac{a_{177}}{a_{179}} = -\frac{{}_{180}C_{177} \times 2i^{177}}{{}_{180}C_{179} \times 2i^{179}} = \frac{15931}{3} = 5310.333$$