

## Individual Events

<b>SI</b>	<b>a</b>	900	<b>I1</b>	<b>a</b>	10	<b>I2</b>	<b>a</b>	$\frac{1}{2} (=0.5)$	<b>I3</b>	<b>a</b>	-7	<b>I4</b>	<b>a</b>	15	<b>I5</b>	<b>a</b>	80
	<b>b</b>	7		<b>b</b>	1		<b>b</b>	5		<b>b</b>	6		<b>b</b>	8		<b>b</b>	4
	<b>c</b>	2		<b>c</b>	4		<b>c</b>	10		<b>x</b>	$\frac{1}{2} (=0.5)$		<b>c</b>	4		<b>N</b>	10
	<b>d</b>	9		<b>d</b>	-5		<b>d</b>	15		<b>y</b>	-1		<b>d</b>	12		<b>x</b>	144

## Group Events

<b>SG</b>	<b>a</b>	2	<b>G6</b>	<b>p</b>	10	<b>G7</b>	<b>p</b>	75	<b>G8</b>	<b>M</b>	1	<b>G9</b>	<b>x</b>	$\frac{1}{100}$	<b>G10</b>	<b>A</b>	50
	<b>b</b>	*136 see the remark		<b>q</b>	15		<b>q</b>	$\frac{1}{2} (=0.5)$		<b>N</b>	6		<b>A</b>	52		<b>S</b>	2
	<b>c</b>	-6		<b>r</b>	24		<b>a</b>	2		<b>R</b>	8		<b>m</b>	501		<b>n</b>	7
	<b>d</b>	7		<b>s</b>	27		<b>m</b>	14		<b>Y</b>	2		<b>P</b>	36		<b>d</b>	5

## Sample Individual Event (1988 Sample Individual Event)


**SI.1** In the given diagram, the sum of the three marked angles is  $a^\circ$ . Find  $a$ .

**Reference: 1987 FSG.3, 1989 FSI.1**

Sum of interior angles of a triangle =  $180^\circ$

angle sum of three vertices =  $3 \times 360^\circ = 1080^\circ$

$a = 1080 - 180 = 900$



**SI.2** The sum of the interior angles of a regular  $b$ -sided polygon is  $a^\circ$ . Find  $b$ .

**Reference 1989 FSI.2**

$a = 900 = 180 \times (b - 2)$

$b = 7$

**SI.3** If  $8^b = c^{21}$ , find  $c$ .

$8^7 = c^{21}$

$2^{21} = c^{21}$

$c = 2$

**SI.4** If  $c = \log_d 81$ , find  $d$ .

$2 = c = \log_d 81$  and  $d > 0$

$d^2 = 81$

$d = 9$

## Individual Event 1

**I1.1** If  $100a = 35^2 - 15^2$ , find  $a$ .

**Reference: 1987 FSG.1, 1988 FI2.2**

$100a = (35 + 15)(35 - 15) = 50 \times 20 = 1000$

$a = 10$

**I1.2** If  $(a - 1)^2 = 3^{4b}$ , find  $b$ .

$9^2 = 3^{4b}$

$4b = 4$

$\Rightarrow b = 1$

**I1.3** If  $b$  is a root of  $x^2 + cx - 5 = 0$ , find  $c$ .

Put  $x = 1$  into the equation:  $1 + c - 5 = 0$

$c = 4$

**I1.4** If  $x + c$  is a factor of  $2x^2 + 3x + 4d$ , find  $d$ .

$x + 4$  is a factor

Put  $x = -4$  into the polynomial:  $2(-4)^2 + 3(-4) + 4d = 0$

$d = -5$

**Individual Event 2**

**I2.1** If  $\alpha, \beta$  are roots of  $x^2 - 10x + 20 = 0$ , find  $a$ , where  $a = \frac{1}{\alpha} + \frac{1}{\beta}$ .

$$\alpha + \beta = 10, \alpha\beta = 20$$

$$a = \frac{1}{\alpha} + \frac{1}{\beta} = \frac{\alpha + \beta}{\alpha\beta} = \frac{10}{20} = \frac{1}{2}$$

**I2.2** If  $\sin \theta = a$  ( $0^\circ < \theta < 90^\circ$ ), and  $10 \cos 2\theta = b$ , find  $b$ .

$$\sin \theta = \frac{1}{2}$$

$$\Rightarrow \theta = 30^\circ$$

$$b = 10 \cos 60^\circ = 5$$

**I2.3** The point  $A(b, c)$  lies on the line  $2y = x + 15$ . Find  $c$ .

**Reference: 1984 FI2.3**

Put  $x = b = 5, y = c$  into the line:  $2c = 5 + 15$

$$c = 10$$

**I2.4** If  $x^2 - cx + 40 \equiv (x + k)^2 + d$ , find  $d$ .

**Reference: 1985 FG10.2, 1986 FG7.3, 1987 FSI.1, 1988 FG9.3**

$$x^2 - 10x + 40 \equiv (x - 5)^2 + 15$$

$$k = -5, d = 15$$

**Individual Event 3**

**I3.1** If  $a$  is the remainder when  $2x^3 - 3x^2 + x - 1$  is divided by  $x + 1$ , find  $a$ .

$$a = 2(-1)^3 - 3(-1)^2 - 1 - 1 = -7$$

**I3.2** If  $b \text{ cm}^2$  is the total surface area of a cube of side  $(8 + a) \text{ cm}$ , find  $b$ .

**Similar Questions: 1984 FG9.2, 1985 FSI.2**

$$8 + a = 1$$

$$b = 6$$

**I3.3** One ball is taken at random from a bag containing  $b + 4$  red balls and  $2b - 2$  white balls.

If  $x$  is the probability that the ball is white, find  $x$ .

There are  $b + 4 = 10$  red balls and  $2b - 2 = 10$  white balls

$$x = \frac{1}{2}$$

**I3.4** If  $\sin \theta = x$  ( $90^\circ < \theta < 180^\circ$ ) and  $\tan(\theta - 15^\circ) = y$ , find  $y$ .

$$\sin \theta = \frac{1}{2}$$

$$\Rightarrow \theta = 150^\circ$$

$$y = \tan(\theta - 15^\circ) = \tan 135^\circ = -1$$


### Individual Event 4

**I4.1** In figure 1,  $DE \parallel BC$ . If  $AD = 4$ ,  $DB = 6$ ,  $DE = 6$  and  $BC = a$ , find  $a$ .

$\triangle ADE \sim \triangle ABC$  (equiangular)

$$\frac{a}{6} = \frac{10}{4} \quad (\text{ratio of sides, } \sim \Delta \text{'s})$$


$$a = 15$$



**I4.2**  $\theta$  is an acute angle such that  $\cos \theta = \frac{a}{17}$ . If  $\tan \theta = \frac{b}{15}$ , find  $b$ .

$$b^2 = 17^2 - 15^2$$

$$b = 8$$



**I4.3** If  $c^3 = b^2$ , find  $c$ .

$$c^3 = 8^2 = 64 = 4^3$$

$$\Rightarrow c = 4$$

**I4.4** The area of an equilateral triangle is  $c\sqrt{3} \text{ cm}^2$ . If its perimeter is  $d \text{ cm}$ , find  $d$ .

**Reference:** 1985 FSL.4, 1986 FSG.3, 1987 FG6.2, 1988 FG9.1

$$\text{Each side} = \frac{d}{3} \text{ cm}$$

$$\frac{1}{2} \cdot \left(\frac{d}{3}\right)^2 \sin 60^\circ = c\sqrt{3} = 4\sqrt{3}$$


$$d = 12$$

### Individual Event 5

**I5.1** In Figure 2, find  $a$ .

$$100 + (180 - a) + 50 + 110 = 360 \quad (\text{sum of ext. } \angle \text{ of } \Delta)$$

$$a = 80$$



**I5.2** If  $b = \log_2 \left( \frac{a}{5} \right)$ , find  $b$ .

$$2^b = 16$$

$$b = 4$$

**I5.3** A piece of string, 20 m long, is divided into 3 parts in the ratio of  $b - 2 : b : b + 2$ . If  $N \text{ m}$  is the length of the longest portion, find  $N$ .

$$b - 2 : b : b + 2 = 2 : 4 : 6 = 1 : 2 : 3$$

$$N = 20 \times \frac{3}{1+2+3} = 10$$

**I5.4** Each interior angle of an  $N$ -sided regular polygon is  $x^\circ$ . Find  $x$ .

$$x = \frac{180 \times (10 - 2)}{10} = 144$$

**Sample Group Event**

**SG.1** The sum of 2 numbers is 20, their product is 10. If the sum of their reciprocals is  $a$ , find  $a$ .

**Reference:** 1983 FG6.3, 1985 FSI.1, 1986 FSG.1

Let the 2 numbers be  $x$  and  $y$ .

$$x + y = 20 \text{ and } xy = 10$$

$$a = \frac{1}{x} + \frac{1}{y} = \frac{x+y}{xy} = 2$$

**SG.2**  $1^2 - 1 = 0 \times 2$ ,  $2^2 - 1 = 1 \times 3$ ,  $3^2 - 1 = 2 \times 4$ , ...,  $b^2 - 1 = 135 \times 137$ . If  $b > 0$ , find  $b$ .

**Reference:** 1983 FI10.1, 1991 FI2.1

$$135 \times 137 = (136 - 1) \times (136 + 1) = 136^2 - 1$$

$$b = 136$$

**Remark** The original question is:

$$1^2 - 1 = 0 \times 2, 2^2 - 1 = 1 \times 3, 3^2 - 1 = 2 \times 4, \dots, b^2 - 1 = 135 \times 137, \text{ find } b.$$

$b = 136$  or  $-136$ , there are 2 different answers!

**SG.3** If the lines  $x + 2y + 1 = 0$  and  $cx + 3y + 1 = 0$  are perpendicular, find  $c$ .

**Reference:** 1983 FG9.3, 1985 FI4.1, 1986 FSG.2, 1987 FG10.2, 1988 FG8.2

$$-\frac{1}{2} \times \left(-\frac{c}{3}\right) = -1 \Rightarrow c = -6$$

**SG.4** The points  $(2, -1)$ ,  $(0, 1)$ ,  $(c, d)$  are collinear. Find  $d$ .

**Reference:** 1984 FG7.3, 1986 FG6.2, 1987 FG7.4, 1989 HG8

$$\frac{d-1}{-6} = \frac{1-(-1)}{0-2}$$

$$d = 7$$

**Group Event 6**

**G6.1** If  $p = \frac{21^3 - 11^3}{21^2 + 21 \times 11 + 11^2}$ , find  $p$ . (**Similar questions:** 1985 FG7.1)

$$p = \frac{21^3 - 11^3}{21^2 + 21 \times 11 + 11^2} = \frac{(21-11)(21^2 + 21 \times 11 + 11^2)}{21^2 + 21 \times 11 + 11^2} = 10$$

**G6.2** If  $p$  men can do a job in 6 days and 4 men can do the same job in  $q$  days, find  $q$ .

10 men can do a job in 6 days.

1 man can do a job in 60 days

4 men can do a job in 15 days  $\Rightarrow q = 15$

**G6.3** If the  $q^{\text{th}}$  day of March in a year is Wednesday and the  $r^{\text{th}}$  day of March in the same year is Friday, where  $18 < r < 26$ , find  $r$ . (**Reference:** 1985 FG9.3, 1987 FG6.4, 1988 FG10.2)

15<sup>th</sup> March is Wednesday

17<sup>th</sup> March is Friday

24<sup>th</sup> March is Friday  $\Rightarrow r = 24$

**G6.4** If  $a*b = ab + 1$ , and  $s = (3*4)*2$ , find  $s$ . (**Reference:** 1985 FSG.1)

$$3*4 = 3 \times 4 + 1 = 13$$

$$s = (3*4)*2 = 13*2 = 13 \times 2 + 1 = 27$$

**Group Event 7 (1988 Sample Group Event)****G7.1** The acute angle between the 2 hands of a clock at 3:30 a.m. is  $p^\circ$ . Find  $p$ .**Reference:** 1985 FI3.1 1987 FG7.1, 1989 FI1.1, 1990 FG6.3, 2007 HI1At 3:00 a.m., the angle between the arms of the clock =  $90^\circ$ From 3:00 a.m. to 3:30 a.m., the hour-hand had moved  $360^\circ \times \frac{1}{12} \times \frac{1}{2} = 15^\circ$ .The minute hand had moved  $180^\circ$ .

$$p = 180 - 90 - 15 = 75$$

**G7.2** In  $\triangle ABC$ ,  $\angle B = \angle C = p^\circ$ . If  $q = \sin A$ , find  $q$ .

$$\angle B = \angle C = 75^\circ, \angle A = 180^\circ - 75^\circ - 75^\circ = 30^\circ$$

$$q = \sin 30^\circ = \frac{1}{2}$$

**G7.3** The 3 points (1, 3), (a, 5), (4, 9) are collinear. Find  $a$ .**Reference:** 1984 FSG.4, 1986FG6.2, 1987 FG7.4, 1989 HG8

$$\frac{9-5}{4-a} = \frac{9-3}{4-1} = 2$$

$$\Rightarrow a = 2$$

**G7.4** The average of 7, 9,  $x$ ,  $y$ , 17 is 10. If  $m$  is the average of  $x+3$ ,  $x+5$ ,  $y+2$ , 8,  $y+18$ , find  $m$ .

$$\frac{7+9+x+y+17}{5} = 10$$

$$\Rightarrow x+y = 17$$

$$m = \frac{x+3+x+5+y+2+8+y+18}{5}$$

$$= \frac{2(x+y)+36}{5} = 14$$

**Group Event 8**

In the addition shown, each letter represents a different digit ranging from

zero to nine. It is already known that

 $S = 9$ ,  $O = \text{zero}$ ,  $E = 5$ .

$$\begin{array}{rcccc} & S & E & N & D \\ + & & M & O & R & E \\ \hline M & O & N & E & Y \end{array}$$

Find the numbers represented by

(i)  $M$ , (ii)  $N$ , (iii)  $R$ , (iv)  $Y$ 

Consider the thousands digit and the ten thousands digits.

$$0 \leq S, M \leq 9, 9 + M = 10M + 0 \text{ or } 9 + M + 1 = 10M + 0$$

 $\Rightarrow M = 1$  and there is no carry digit.Consider the hundreds digit.  $5 + 0 + 1 = N$  $\Rightarrow N = 6$  and there is a carry digit.For the tens digit.  $6 + R = 10 + 5$  $\Rightarrow R = 9$  (same as  $S$ , rejected) or  $6 + R + 1 = 10 + 5$  $\Rightarrow R = 8$ 

There is a carry digit in the unit digit

$$D + 5 = 10 + Y, (D, Y) = (7, 2) \Rightarrow Y = 2$$

 $\therefore M = 1, N = 6, R = 8, Y = 2$

## Group Event 9

**G9.1** If  $x = \left(1 - \frac{1}{2}\right)\left(1 - \frac{1}{3}\right)\left(1 - \frac{1}{4}\right) \cdots \left(1 - \frac{1}{100}\right)$ , find  $x$  in the simplest fractional form.

**Reference: 1985 FSG.3, 1986 FG10.4**

$$x = \frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} \times \cdots \times \frac{99}{100} = \frac{1}{100}$$

**G9.2** The length, width and height of a rectangular block are 2, 3 and 4 respectively. Its total surface area is  $A$ , find  $A$ .

**Similar Questions: 1984 FI3.2, 1985 FSI.2**

$$A = 2 \times (2 \times 3 + 3 \times 4 + 2 \times 4) = 52$$

**G9.3** The average of the integers 1, 2, 3, ..., 1001 is  $m$ . Find  $m$ .

$$\begin{aligned} m &= \frac{1}{1001} (1 + 2 + 3 + \cdots + 1001) \\ &= \frac{1}{1001} \cdot \frac{(1 + 1001) \cdot 1001}{2} = 501 \end{aligned}$$

**G9.4** The area of a circle inscribed in an equilateral triangle is  $12\pi$ .

If  $P$  is the perimeter of this triangle, find  $P$ .

**Reference: 1990 FI2.3**

Let the radius be  $r$  and the centre be  $O$ .

$$\pi r^2 = 12\pi$$


$$\Rightarrow r = 2\sqrt{3}$$

The length of one side of the equilateral triangle is  $\frac{P}{3}$ .

$$\frac{P}{3} = 2r \cot 30^\circ$$

$$= 2\sqrt{3}r = 12$$

$$P = 36$$



**Group Event 10**


**G10.1** If  $A$  is the area of a square inscribed in a circle of diameter 10, find  $A$ .

**Reference: 1985 FSG.4, 1989 FI3.3**

Let the square be  $BCDE$ .

$$BC = 10 \cos 45^\circ = 5\sqrt{2}$$

$$A = (5\sqrt{2})^2 = 50$$



**G10.2** If  $a + \frac{1}{a} = 2$ , and  $S = a^3 + \frac{1}{a^3}$ , find  $S$ .

**Reference: 1998 HG1**

$$a^2 + \frac{1}{a^2} = \left(a + \frac{1}{a}\right)^2 - 2 = 4 - 2 = 2$$

$$S = a^3 + \frac{1}{a^3}$$

$$= \left(a + \frac{1}{a}\right) \left(a^2 - 1 + \frac{1}{a^2}\right)$$

$$= 2(2 - 1) = 2$$

**G10.3** An  $n$ -sided convex polygon has 14 diagonals. Find  $n$ .

**Reference: 1985 FG8.3, 1988 FG6.2, 1989 FG6.1, 1991 FI2.3, 2001 FI4.2, 2005 FI1.4**

$$\text{Number of diagonals} = C_2^n - n = \frac{n(n-1)}{2} - n = 14$$

$$n^2 - 3n - 28 = 0$$

$$(n-7)(n+4) = 0$$

$$\Rightarrow n = 7$$

**G10.4** If  $d$  is the distance between the 2 points  $(2, 3)$  and  $(-1, 7)$ , find  $d$ .

**Reference: 1986 FG9.4**

$$d = \sqrt{[2 - (-1)]^2 + (3 - 7)^2} = 5$$