

Individual Events

SI	<i>h</i>	4	I1	<i>a</i>	5	I2	<i>p</i>	3	I3	<i>a</i>	1000	I4	<i>a</i>	5	I5	<i>a</i>	17
	<i>k</i>	32		<i>b</i>	4		<i>q</i>	36		<i>b</i>	8		<i>b</i>	12		<i>b</i>	5
	<i>p</i>	3		<i>c</i>	10		<i>k</i>	12		<i>c</i>	16		<i>c</i>	4		<i>c</i>	23
	<i>q</i>	16		<i>d</i>	34		<i>m</i>	150		<i>d</i>	1		<i>d</i>	12		<i>d</i>	9

Group Events

SG	<i>a</i>	2	G6	<i>a</i>	150	G7	<i>C</i>	47	G8	<i>A</i>	2	G9	<i>S</i>	1000	G10	<i>A</i>	1584
	<i>b</i>	-3		<i>b</i>	10		<i>K</i>	2		<i>B</i>	3		<i>K</i>	98		<i>k</i>	14
	<i>p</i>	60		<i>k</i>	37.5		<i>A</i>	1		<i>C</i>	7		<i>t</i>	20		<i>x</i>	160
	<i>q</i>	136		<i>d</i>	6		<i>B</i>	5		<i>k</i>	9		<i>d</i>	5		<i>n</i>	15

Sample Individual Event (1986 Final Individual Event 2)

SI.1 Given that $3x^2 - 4x + \frac{h}{3} = 0$ has equal roots, find h .

$$\Delta = (-4)^2 - 4(3) \cdot \frac{h}{3} = 0$$

$$h = 4$$

SI.2 If the height of a cylinder is doubled and the new radius is h times the original, then the new volume is k times the original. Find k .

Let the old height be x , old radius be r , then the old volume is $\pi r^2 x$.

The new height is $2x$, the new radius is $4r$,

then the new volume is $\pi(4r)^2(2x) = 32\pi r^2 x$

$$k = 32$$

SI.3 If $\log_{10} 210 + \log_{10} k - \log_{10} 56 + \log_{10} 40 - \log_{10} 120 + \log_{10} 25 = p$, find p .

$$p = \log_{10} \left(\frac{210 \times 32 \times 40 \times 25}{56 \times 120} \right)$$

$$= \log_{10} 1000 = 3$$

SI.4 If $\sin A = \frac{p}{5}$ and $\frac{\cos A}{\tan A} = \frac{q}{15}$, find q .

$$\sin A = \frac{3}{5}$$

$$\frac{\cos A}{\tan A} = \frac{q}{15}$$

$$\frac{\cos^2 A}{\sin A} = \frac{q}{15}$$

$$\frac{1 - \sin^2 A}{\sin A} = \frac{1 - \left(\frac{3}{5}\right)^2}{\frac{3}{5}} = \frac{16}{15} = \frac{q}{15}$$

$$q = 16$$

Individual Event 1**I1.1** Find a if $2t + 1$ is a factor of $4t^2 + 12t + a$.

$$\text{Let } f(t) = 4t^2 + 12t + a$$

$$f\left(-\frac{1}{2}\right) = 4\left(-\frac{1}{2}\right)^2 + 12\left(-\frac{1}{2}\right) + a = 0$$

$$a = 5$$

I1.2 \sqrt{K} denotes the nonnegative square root of K , where $K \geq 0$. If b is the root of the equation

$$\sqrt{a-x} = x-3, \text{ find } b.$$

$$(\sqrt{5-x})^2 = (x-3)^2$$

$$\Rightarrow 5-x = x^2 - 6x + 9$$

$$\Rightarrow x^2 - 5x + 4 = 0$$

$$\Rightarrow x = 1 \text{ or } 4$$

$$\text{When } x = 1, \text{ LHS} = 2 \neq -1 = \text{RHS}$$

$$\text{When } x = 4, \text{ LHS} = 1 = \text{RHS.}$$

$$\therefore x = b = 4$$

I1.3 If c is the greatest value of $\frac{20}{b+2\cos\theta}$, find c .

$$\frac{20}{b+2\cos\theta} = \frac{20}{4+2\cos\theta} = \frac{10}{2+\cos\theta}$$

$$c = \text{the greatest value} = \frac{10}{2-1} = 10$$

I1.4 A man drives a car at $3c$ km/h for 3 hours and then $4c$ km/h for 2 hours. If his average speed for the whole journey is d km/h, find d .

$$\text{Total distance travelled} = (30 \times 3 + 40 \times 2) \text{ km} = 170 \text{ km}$$

$$d = \frac{170}{3+2} = 34$$

Individual Event 2

I2.1 If $0^\circ \leq \theta < 360^\circ$, the equation in θ : $3 \cos \theta + \frac{1}{\cos \theta} = 4$ has p roots. Find p .

$$\begin{aligned} 3 \cos^2 \theta + 1 &= 4 \cos \theta \\ \Rightarrow 3 \cos^2 \theta - 4 \cos \theta + 1 &= 0 \\ \Rightarrow \cos \theta &= \frac{1}{3} \text{ or } 1 \end{aligned}$$

$$p = 3$$

I2.2 If $x - \frac{1}{x} = p$ and $x^3 - \frac{1}{x^3} = q$, find q .

Reference: 2009 FI2.3

$$x - \frac{1}{x} = 3; \left(x - \frac{1}{x}\right)^2 = 9$$

$$\Rightarrow x^2 + \frac{1}{x^2} = 11$$

$$q = x^3 - \frac{1}{x^3}$$

$$= \left(x - \frac{1}{x}\right) \left(x^2 + 1 + \frac{1}{x^2}\right)$$

$$= 3(11 + 1) = 36$$

I2.3 A circle is inscribed in an equilateral triangle of perimeter q cm. If the area of the circle is $k\pi \text{ cm}^2$, find k .

Reference: 1984 FG9.4

Let the equilateral triangle be ABC , the centre of the inscribed circle is O , which touches the triangle at D and E , with radius r cm

Perimeter = 36 cm

\Rightarrow Each side = 12 cm

$\angle ACB = 60^\circ$ (\angle s of an equilateral Δ)

$\angle ODC = 90^\circ$ (tangent \perp radius)

$\angle OCD = 30^\circ$ (tangent from ext. pt.)

$CD = 6$ cm (tangent from ext. pt.)

$$r = 6 \tan 30^\circ = 2\sqrt{3}$$

$$\text{Area of circle} = \pi(2\sqrt{3})^2 \text{ cm}^2 = 12\pi \text{ cm}^2$$


$$k = 12$$

I2.4 Each interior angle of a regular polygon of k sides is m° . Find m .

Angle sum of 12-sides polygon = $180^\circ(12 - 2) = 1800^\circ$

Each interior angle = $m^\circ = 1800^\circ \div 12 = 150^\circ$

$$m = 150$$



Individual Event 3**I3.1** If $998a + 1 = 999^2$, find a .

$$\begin{aligned}998a &= 999^2 - 1 \\&= (999 - 1)(999 + 1) \\&= 998 \times 1000 \\a &= 1000\end{aligned}$$

I3.2 If $\log_{10}a = \log_2b$, find b .

$$\begin{aligned}\log_{10} 1000 &= \log_2 b \\ \log_2 b &= 3 \\ \Rightarrow b &= 2^3 = 8\end{aligned}$$

I3.3 The area of the triangle formed by the x -axis, the y -axis and the line $2x + y = b$ is c sq. units. Find c .**Reference: 1994 FI5.3**

$$2x + y = 8; x\text{-intercept} = 4, y\text{-intercept} = 8$$

$$c = \text{area} = \frac{1}{2} \cdot 4 \times 8 = 16$$

I3.4 If $64t^2 + ct + d$ is a perfect square, find d .

$$64t^2 + 16t + d \text{ has a double root}$$

$$\Delta = 16^2 - 4 \times 64d = 0$$

$$d = 1$$

Individual Event 4**I4.1** Solve for a in the equation $2^{a+1} + 2^a + 2^{a-1} = 112$.

$$2^a \cdot (2 + 1 + \frac{1}{2}) = 112$$

$$2^a = 32$$

$$a = 5$$

Method 2

$$112 = 64 + 32 + 16 = 2^6 + 2^5 + 2^4$$

$$a = 5$$

I4.2 If a is one root of the equation $x^2 - bx + 35 = 0$, find b .One root of $x^2 - bx + 35 = 0$ is 5

$$\Rightarrow 5^2 - 5b + 35 = 0$$

$$\Rightarrow b = 12$$

I4.3 If $\sin \theta = \frac{-b}{15}$, where $180^\circ < \theta < 270^\circ$, and $\tan \theta = \frac{c}{3}$, find c .

$$\sin \theta = -\frac{12}{15} = -\frac{4}{5}$$

$$\Rightarrow \tan \theta = \frac{4}{3}$$

$$\Rightarrow c = 4$$

I4.4 The probability of getting a sum of c in throwing two dice is $\frac{1}{d}$. Find d .

$$P(\text{sum} = 4) = P((1,3), (2, 2), (3, 1))$$


$$= \frac{3}{36} = \frac{1}{12} = \frac{1}{d}$$

$$\Rightarrow d = 12$$

Individual Event 5**I5.1** In the figure, find a .

$$a^2 - 8^2 = 12^2 + 9^2 \text{ (Pythagoras' Theorem)}$$

$$a = 17$$

**I5.2** If the lines $ax + by = 1$ and $10x - 34y = 3$ are perpendicular to each other, find b .

$$17x + by = 1 \text{ is perpendicular to } 10x - 34y = 3$$

$$\Rightarrow \text{product of slopes} = -1$$

$$-\frac{17}{b} \times \frac{10}{34} = -1$$

$$\Rightarrow b = 5$$

I5.3 If the b^{th} day of May in a year is Friday and the c^{th} day of May in the same year is Tuesday, where $16 < c < 24$, find c .5th May is a Friday \Rightarrow 9th May is Tuesday \Rightarrow 16th May is Tuesday \Rightarrow 23rd May is Tuesday

$$c = 23$$

I5.4 c is the d^{th} prime number. Find d .**Reference: 1985 FSG.2, 1989 FSG.3**

The first few prime numbers are: 2, 3, 5, 7, 11, 13, 17, 19, 23

23 is the 9th prime number

$$d = 9$$

Sample Group Event (1986 Sample Group Event)**SG.1** The sum of two numbers is 50, and their product is 25.If the sum of their reciprocals is a , find a .Let the 2 numbers be x, y .

$$x + y = 50, xy = 25$$

$$\Rightarrow a = \frac{1}{x} + \frac{1}{y}$$

$$= \frac{x+y}{xy}$$

$$= \frac{50}{25} = 2$$

SG.2 If the lines $ax + 2y + 1 = 0$ and $3x + by + 5 = 0$ are perpendicular, find b . $2x + 2y + 1 = 0$ is \perp to $3x + by + 5 = 0$ \Rightarrow product of slopes $= -1$

$$-\frac{2}{2} \times \frac{-3}{b} = -1$$

$$\Rightarrow b = -3$$

SG.3 The area of an equilateral triangle is $100\sqrt{3} \text{ cm}^2$. If its perimeter is $p \text{ cm}$, find p .Let the length of one side be $x \text{ cm}$.

$$\frac{1}{2}x^2 \sin 60^\circ = 100\sqrt{3}$$

$$\Rightarrow x = 20$$

$$\Rightarrow p = 60$$

SG.4 If $x^3 - 2x^2 + px + q$ is divisible by $x + 2$, find q .

$$\text{Let } f(x) = x^3 - 2x^2 + 60x + q$$

$$f(-2) = -8 - 8 - 120 + q = 0$$

$$q = 136$$

Group Event 6

G6.1 If $a = \frac{(68^3 - 65^3) \cdot (32^3 + 18^3)}{(32^2 - 32 \times 18 + 18^2) \cdot (68^2 + 68 \times 65 + 65^2)}$, find a .

$$a = \frac{(32+18)(32^2 - 32 \times 18 + 18^2) \cdot (68-65)(68^2 + 68 \times 65 + 65^2)}{(32^2 - 32 \times 18 + 18^2) \cdot (68^2 + 68 \times 65 + 65^2)}$$

$$= 50 \times 3 = 150$$

G6.2 If the 3 points (a, b) , $(10, -4)$ and $(20, -3)$ are collinear, find b .

The slopes are equal: $\frac{b+4}{150-10} = \frac{-3+4}{20-10}$

$$\Rightarrow b = 10$$

G6.3 If the acute angle formed by the hands of a clock at 4:15 is k° , find k .

Reference 1984 FG7.1, 1985 FI3.1, 1987 FG7.1, 1989 FI1.1, 2007 HI1


$$k = 30 + 30 \times \frac{1}{4} = 37.5$$

G6.4 In the figure, $PQ = 10$, $RS = 15$, $QS = 20$. If $XY = d$, find d .

Reference: 1985 FI2.4, 1989 HG8

$$\frac{1}{d} = \frac{1}{10} + \frac{1}{15} = \frac{25}{150} = \frac{1}{6}$$

$$d = 6$$



Group Event 7**G7.1** 2 apples and 3 oranges cost 6 dollars.

4 apples and 7 oranges cost 13 dollars.

16 apples and 23 oranges cost C dollars. Find C .Let the cost of one apple be $\$x$ and one orange be $\$y$.

$$2x + 3y = 6 \dots\dots (1)$$

$$4x + 7y = 13 \dots\dots (2)$$

$$(2) - 2(1): y = 1, x = 1.5$$

$$C = 16x + 23y = 24 + 23 = 47$$

G7.2 If $K = \frac{6 \cos \theta + 5 \sin \theta}{2 \cos \theta + 3 \sin \theta}$ and $\tan \theta = 2$, find K .**Reference: 1986 FG10.3, 1987 FG8.1, 1989 FSG.4, 1989 FG10.3**

$$\begin{aligned} K &= \frac{6 \frac{\cos \theta}{\cos \theta} + 5 \frac{\sin \theta}{\cos \theta}}{2 \frac{\cos \theta}{\cos \theta} + 3 \frac{\sin \theta}{\cos \theta}} \\ &= \frac{6 + 5 \tan \theta}{2 + 3 \tan \theta} \\ &= \frac{6 + 5 \times 2}{2 + 3 \times 2} = 2 \end{aligned}$$

G7.3 and **G7.4** A, B are positive integers less than 10 such that $21A104 \times 11 = 2B8016 \times 9$.**Similar Questions 1985 FG8.1-2, 1988 FG8.3-4****G7.3** Find A .11 and 9 are relatively prime, $21A104$ is divisible by 9.

$$2 + 1 + A + 1 + 0 + 4 = 9m$$

$$\Rightarrow 8 + A = 9m$$

$$\Rightarrow A = 1$$

G7.4 Find B . $2B8016$ is divisible by 11.

$$2 + 8 + 1 - (B + 0 + 6) = 11n$$

$$\Rightarrow 11 - (B + 6) = 11n$$

$$\Rightarrow B = 5$$

Group Event 8

In the multiplication shown, the letters A , B , C and K ($A < B$) represent different integers from 1 to 9.

$$\begin{array}{r} \\ B \\ \hline K K \end{array}$$

(Hint: $KKK = K \times 111$.)

G8.1 Find A .

$$1^2 = 1, 2^2 = 4, 3^2 = 9, 4^2 = 16, 5^2 = 25, 6^2 = 36, 7^2 = 49, 8^2 = 64, 9^2 = 81$$

Possible $K = 1, 4, 5, 6, 9$

$$100K + 10K + K = 111K = 3 \times 37K, 37 \text{ is a prime number}$$

Either $10A + C$ or $10B + C$ is divisible by 37

$$10B + C = 37 \text{ or } 74$$

$$\text{When } B = 3, C = 7, K = 9$$

$$999 \div 37 = 27$$

$$\therefore A = 2$$

G8.2 Find B .

$$B = 3$$

G8.3 Find C .

$$C = 7$$

G8.4 Find K .

$$K = 9$$

Group Event 9**G9.1** If $S = ab - 1 + a - b$ and $a = 101$, $b = 9$, find S .**Reference: 1985 FG8.4, 1986 FG9.3, 1988 FG6.3**

$$S = (a - 1)(b + 1) = 100 \times 10 = 1000$$

G9.2 If $x = 1.9\dot{8}\dot{9}$ and $x - 1 = \frac{K}{99}$, find K .

$$x = 1.9 + \frac{89}{990}$$

$$\begin{aligned} x - 1 &= \frac{K}{99} = \frac{9}{10} + \frac{89}{990} \\ &= \frac{9 \times 99 + 89}{990} = \frac{980}{990} = \frac{98}{99} \end{aligned}$$

$$K = 98$$

G9.3 The average of p , q and r is 18. The average of $p + 1$, $q - 2$, $r + 3$ and t is 19. Find t .

$$\frac{p + q + r}{3} = 18$$

$$\Rightarrow p + q + r = 54$$

$$\frac{p + 1 + q - 2 + r + 3 + t}{4} = 19$$

$$\Rightarrow p + q + r + 2 + t = 76$$

$$\Rightarrow 54 + 2 + t = 76$$

$$t = 20$$

G9.4 In the figure, \widehat{QR} , \widehat{RP} , \widehat{PQ} are 3 arcs, centres at X , Y and Z respectively, touching one another at P , Q and R . If $ZQ = d$, $XR = 3$, $YP = 12$, $\angle X = 90^\circ$, find d .**Reference: 1986 FG7.1**

$$XZ = 3 + d, XY = 3 + 12 = 15, YZ = 12 + d$$


$$XZ^2 + XY^2 = YZ^2 \text{ (Pythagoras' theorem)}$$

$$(3 + d)^2 + 15^2 = (12 + d)^2$$

$$9 + 6d + d^2 + 225 = 144 + 24d + d^2$$

$$18d = 90$$

$$\Rightarrow d = 5$$



Group Event 10**G10.1** If $A = 1 + 2 - 3 + 4 + 5 - 6 + 7 + 8 - 9 + \dots + 97 + 98 - 99$, find A .**Reference: 1985 FG7.4, 1988 FG6.4, 1991 FSI.1, 1992 FI1.4**

$$A = (1 + 2 - 3) + (4 + 5 - 6) + (7 + 8 - 9) + \dots + (97 + 98 - 99)$$

$$A = 0 + 3 + 6 + \dots + 96 = \frac{3+96}{2} \times 32 = 99 \times 16 = 1584$$

G10.2 If $\log_{10}(k-1) - \log_{10}(k^2 - 5k + 4) + 1 = 0$, find k .

$$10(k-1) = k^2 - 5k + 4$$

$$k^2 - 15k + 14 = 0$$

$$k = 1 \text{ or } 14$$

When $k = 1$, LHS is undefined \therefore rejected

When $k = 14$, LHS = $\log_{10} 13 - \log_{10}(14-1)(14-4) + 1 = \text{RHS}$

$$\therefore k = 14$$

G10.3 and **G10.4** One interior angle of a convex n -sided polygon is x° . The sum of the remaining interior angles is 2180° .**Reference: 1989 HG2, 1992 HG3, 2002 FI3.4, 2013 HI6****G10.3** Find x .

$$2180 + x = 180(n-2) \text{ (}\angle\text{s sum of polygon)}$$

$$2160 + 20 + x = 180 \times 12 + 20 + x = 180(n-2)$$

$$\therefore x < 180$$

$$\therefore 20 + x = 180$$

$$x = 160$$

G10.4 Find n .

$$n - 2 = 12 + 1$$

$$n = 15$$