

**Individual Events**

| I1 | a | 1             | I2 | R | * $\frac{9}{25}$<br>see the remark | I3 | S | 10             | I4 | k | 1   |
|----|---|---------------|----|---|------------------------------------|----|---|----------------|----|---|---|
|    | b | $\frac{1}{2}$ |    | S | 1                                  |    | R | 30             |    | r | 2   |
|    | C | 10            |    | T | 1                                  |    | T | 6              |    | s | $\frac{1}{\sqrt{2}} (= \frac{\sqrt{2}}{2})$ |
|    | D | -2            |    | W | $\sqrt{5}$                         |    | P | $\sqrt{7} + 2$ |    | w | 9   |

**Group Events**

| G1 | k | 1               | G2 | w | 45            | G3 | r | 2006                    | G4 | R | $12\sqrt{3}$         |
|----|---|-----------------|----|---|---------------|----|---|-------------------------|----|---|----------------------|
|    | B | $\frac{16}{15}$ |    | z | -13           |    | x | $\frac{7}{4} (= 1.75)$  |    | S | *8<br>see the remark |
|    | C | $\frac{1}{4}$   |    | s | $\frac{1}{4}$ |    | z | 30                      |    | T | $\frac{1}{2}$        |
|    | a | 1               |    | t | 14            |    | R | $\frac{15}{4} (= 3.75)$ |    | W | 2013021              |

**Individual Event 1**

- I1.1** If  $a$  is a real number satisfying  $\log_2(x+3) - \log_2(x+1) = 1$ , find the value of  $a$ .

$$\log_2 \frac{x+3}{x+1} = \log_2 2$$

$$x+3 = 2x+2$$

$$x=1 \Rightarrow a=1$$

- I1.2** In Figure 1,  $O$  is the centre of the circle with radius 1 cm. If the length of the arc  $AB$  is equal to  $a$  cm and the area of the shaded sector  $OAB$  is equal to  $b$   $\text{cm}^2$ , find the value of  $b$ . (Take  $\pi = 3$ )

$$b = \frac{1}{2}rs = \frac{1}{2}1 \cdot 1 = \frac{1}{2}$$




Figure 1

- I1.3** An interior angle of a regular  $C$ -sided polygon is  $288b^\circ$ , find the value of  $C$ .  
Each interior angle =  $288b^\circ = 144^\circ$

$$\text{Each exterior angle} = 36^\circ = \frac{360^\circ}{C}$$

$$C = 10$$

- I1.4** Given that  $C$  is a root of the equation  $kx^2 + 2x + 5 = 0$ , where  $k$  is a constant. If  $D$  is another root, find the value of  $D$ .

$$100k + 20 + 5 = 0 \Rightarrow k = -\frac{1}{4}$$

$$C + D = \text{sum of roots} = -\frac{2}{k}$$

$$10 + D = 8 \Rightarrow D = -2$$

**Individual Event 2**

**I2.1** Given that  $a : b : c = 6 : 3 : 1$ . If  $R = \frac{3b^2}{2a^2 + bc}$ , find the value of  $R$ .

$$\text{Let } a = 6k, b = 3k, c = k, \text{ then } R = \frac{3(3k)^2}{2(6k)^2 + (3k)k} = \frac{9}{25}$$

**Remark:** original question is Given that  $a, b$  and  $c$  are three numbers not equal to 0 and  $a : b : c = 6 : 3 : 1 \dots\dots$ , the condition  $a : b : c = 6 : 3 : 1$  already implies that  $a, b$  and  $c$  are not zero.

**I2.2** Given that  $\frac{|k+R|}{|R|} = 0$ . If  $S = \frac{|k+2R|}{|2k+R|}$ , find the value of  $S$ .

$$k + \frac{9}{25} = 0 \Rightarrow k = -\frac{9}{25}$$

$$S = \frac{|k+2R|}{|2k+R|} = \frac{\left|-\frac{9}{25} + \frac{18}{25}\right|}{\left|-\frac{18}{25} + \frac{9}{25}\right|} = 1$$

**I2.3** Given that  $T = \sin 50^\circ \times (S + \sqrt{3} \times \tan 10^\circ)$ , find the value of  $T$ .

$$\begin{aligned} T &= \sin 50^\circ \times \left(1 + \sqrt{3} \cdot \frac{\sin 10^\circ}{\cos 10^\circ}\right) \\ &= \frac{\sin 50^\circ}{\cos 10^\circ} \cdot (\cos 10^\circ + \sqrt{3} \sin 10^\circ) \\ &= \frac{2 \sin 50^\circ}{\cos 10^\circ} \left(\frac{1}{2} \cos 10^\circ + \frac{\sqrt{3}}{2} \sin 10^\circ\right) \\ &= \frac{2 \sin 50^\circ}{\cos 10^\circ} (\cos 60^\circ \cdot \cos 10^\circ + \sin 60^\circ \cdot \sin 10^\circ) \\ &= \frac{2 \sin 50^\circ \cos 50^\circ}{\cos 10^\circ} = \frac{\sin 100^\circ}{\cos 10^\circ} = 1 \end{aligned}$$

**I2.4** Given that  $x_0$  and  $y_0$  are real numbers satisfying the system of equations  $\begin{cases} y = \frac{T}{x} \\ y = |x| + T \end{cases}$ .

If  $W = x_0 + y_0$ , find the value of  $W$ .

$$\begin{cases} y = \frac{1}{x} \\ y = |x| + 1 \end{cases}$$

$$\frac{1}{x} = |x| + 1 \dots\dots (*)$$

$$\frac{1}{x} = x + 1 \text{ or } \frac{1}{x} = -x + 1$$

$$1 = x^2 + x \text{ or } 1 = -x^2 + x$$

$$x^2 + x - 1 = 0 \text{ or } x^2 - x + 1 = 0$$

$$x = \frac{-1 \pm \sqrt{5}}{2} \text{ or no solution}$$

$$\text{Check: sub. } x = \frac{-1 - \sqrt{5}}{2} \text{ into (*)}$$

$$\text{LHS} = \frac{2}{-1 - \sqrt{5}} < 0, \text{ RHS} > 0 \text{ (rejected)}$$

$$\text{When } x = \frac{-1 + \sqrt{5}}{2}; \text{ LHS} = \frac{2}{-1 + \sqrt{5}} = \frac{\sqrt{5} + 1}{2}; \text{ RHS} = \frac{-1 + \sqrt{5}}{2} + 1 = \frac{1 + \sqrt{5}}{2} \text{ (accepted)}$$

$$y = \frac{1 + \sqrt{5}}{2} \Rightarrow W = x_0 + y_0 = \frac{-1 + \sqrt{5}}{2} + \frac{1 + \sqrt{5}}{2} = \sqrt{5}$$

**Individual Event 3**

- I3.1** Given that  $\frac{2x-3}{x^2-x} = \frac{A}{x-1} + \frac{B}{x}$ , where  $A$  and  $B$  are constants. If  $S = A^2 + B^2$ , find the value of  $S$

$$\frac{2x-3}{x^2-x} = \frac{Ax+B(x-1)}{(x-1)x}$$

$$A+B=2, -B=-3$$

$$A=-1, B=3$$

$$S=(-1)^2+3^2=10$$

- I3.2** In Figure 1,  $ABCD$  is an inscribed rectangle,  $AB = (S - 2)$  cm and  $AD = (S - 4)$  cm. If the circumference of the circle is  $R$  cm, find the value of  $R$ . (Take  $\pi = 3$ )

$$AB = 8 \text{ cm}, CD = 6 \text{ cm}$$

$$AC = 10 \text{ cm} (\text{Pythagoras' theorem})$$

$$R = 10\pi = 30$$




Figure 1

- I3.3** Given that  $x$  and  $y$  are integers satisfying the equation  $\frac{R}{2}xy = 21x + 20y - 13$ .

If  $T = xy$ , find the value of  $T$ .

$$15xy = 21x + 20y - 13$$

$$(3x-4)(5y-7) = 15$$

$$\begin{cases} 3x-4=1 \\ 5y-7=15 \end{cases} \text{ or } \begin{cases} 3x-4=3 \\ 5y-7=5 \end{cases} \text{ or } \begin{cases} 3x-4=5 \\ 5y-7=3 \end{cases} \text{ or } \begin{cases} 3x-4=15 \\ 5y-7=1 \end{cases} \text{ or } \begin{cases} 3x-4=-1 \\ 5y-7=-15 \end{cases} \text{ or } \begin{cases} 3x-4=-3 \\ 5y-7=-5 \end{cases} \text{ or } \begin{cases} 3x-4=-5 \\ 5y-7=-3 \end{cases} \text{ or } \begin{cases} 3x-4=-15 \\ 5y-7=-1 \end{cases}$$

For integral solution:  $3x-4=5, 5y-7=3$

$$x=3, y=2 \Rightarrow T=3\times 2=6$$

- I3.4** Let  $a$  be the positive root of the equation  $x^2 - 2x - T = 0$ .

If  $P = 3 + \frac{T}{2 + \frac{T}{2 + \frac{T}{2 + \frac{T}{a}}}}$ , find the value of  $P$ .

$$2 + \frac{T}{2 + \frac{T}{2 + \frac{T}{a}}}$$

$$x^2 - 2x - 6 = 0$$

$$a = 1 + \sqrt{7}$$

$$a^2 - 2a - 6 = 0 \Rightarrow a^2 = 2a + 6 \Rightarrow a = 2 + \frac{6}{a}$$

$$2 + \frac{T}{a} = 2 + \frac{6}{a} = a \Rightarrow 2 + \frac{T}{2 + \frac{T}{a}} = 2 + \frac{6}{a} = a \Rightarrow 2 + \frac{T}{2 + \frac{T}{2 + \frac{T}{a}}} = 2 + \frac{6}{a} = a$$

$$P = 3 + \frac{6}{a} = 1 + 2 + \frac{6}{a} = 1 + a = 1 + 1 + \sqrt{7} = 2 + \sqrt{7}$$

**Individual Event 4**

**I4.1** Let  $\frac{k}{4} = \left(1 + \frac{1}{2} + \frac{1}{3}\right) \times \left(\frac{1}{2} + \frac{1}{3} + \frac{1}{4}\right) - \left(1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}\right) \times \left(\frac{1}{2} + \frac{1}{3}\right)$ , find the value of  $k$ .

**Reference: 1995 FG6.2**

Let  $x = 1 + \frac{1}{2} + \frac{1}{3}$ ,  $y = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4}$ , then  $\frac{k}{4} = x(y-1) - y(x-1) = -x + y = \frac{1}{4} \Rightarrow k = 1$

**I4.2** Let  $x$  and  $y$  be real numbers satisfying the equation  $y^2 + 4y + 4 + \sqrt{x+y+k} = 0$ .

If  $r = |xy|$ , find the value of  $r$ .

**Reference: 2005 FI4.1, 2009 FG1.4, 2011 FI4.3, 2013 FI1.4, 2015 HG4, 2015 FI1.1**

The equation is equivalent to  $(y+2)^2 + \sqrt{x+y+k} = 0$

which is a sum of two non-negative numbers.

$\Rightarrow y+2=0$  and  $x+y+1=0$

$y=-2$  and  $x-2+1=0$ ,  $x=1$

$r=|-2 \times 1|=2$

**I4.3** In Figure 1, there are eight positive numbers in series. Starting from the 3<sup>rd</sup> number, each number is the product of the previous two numbers. Given that the 5<sup>th</sup> number is  $\frac{1}{r}$  and the 8<sup>th</sup> number is  $\frac{1}{r^4}$ . If the first number is  $s$ , find the value of  $s$ .

$s$

$\frac{1}{r}$

$\frac{1}{r^4}$

Let the 2<sup>nd</sup> number be  $a$ , then other numbers are as follows:

$s$

$a$

$sa$

$sa^2$

$s^2a^3 =$   
 $\frac{1}{r}$

$s^3a^5$

$s^5a^8$

$s^8a^{13} =$   
 $\frac{1}{r^4}$

$$s^2a^3 = \frac{1}{r} \quad \dots \dots (1)$$

$$s^8a^{13} = \frac{1}{r^4} \quad \dots \dots (2)$$

$$(2) \div (1)^4: a = 1$$

$$\text{Sub. } a = 1 \text{ into (1): } s^2 = \frac{1}{r} = \frac{1}{2}; s > 0 \Rightarrow s = \frac{1}{\sqrt{2}}$$

**I4.4** Let  $[x]$  be the largest integer not greater than  $x$ . For example,  $[2.5] = 2$ .

Let  $w = 1 + [10 \times s^2] + [10 \times s^4] + [10 \times s^6] + \dots + [10 \times s^{2n}] + \dots$ , find the value of  $w$ .

$$w = 1 + [10 \times \frac{1}{2}] + [10 \times \frac{1}{4}] + [10 \times \frac{1}{8}] + \dots + [10 \times \frac{1}{2^n}] + \dots$$

$$= 1 + 5 + 2 + 1 + 0 + 0 + \dots = 9$$

**Group Event 1**

**G1.1** Given that  $k$  is a real number. If  $x^2 + 2kx - 3k^2$  can be divisible by  $x - 1$ , find the greatest value of  $k$ .

By factor theorem,  $1^2 + 2k - 3k^2 = 0$

$$3k^2 - 2k - 1 = 0$$

$$(3k + 1)(k - 1) = 0$$

$$k = -\frac{1}{3} \text{ or } 1$$

Greatest value of  $k = 1$

**G1.2** Given that  $x = x_0$  and  $y = y_0$  satisfy the system of equations  $\begin{cases} \frac{x}{3} + \frac{y}{5} = 1 \\ \frac{x}{5} + \frac{y}{3} = 1 \end{cases}$ . If  $B = \frac{1}{x_0} + \frac{1}{y_0}$ , find the value of  $B$ .

$$\frac{x}{3} + \frac{y}{5} = \frac{x}{5} + \frac{y}{3} = 1 \Rightarrow \frac{x}{3} - \frac{x}{5} = \frac{y}{3} - \frac{y}{5} \Rightarrow x = y$$

$$\text{Sub. } x = y \text{ into the first equation: } \frac{x}{3} + \frac{x}{5} = 1 \Rightarrow x = y = \frac{15}{8} \Rightarrow B = \frac{16}{15}$$

**G1.3** Given that  $x = 2 + \sqrt{3}$  is a root of the equation  $x^2 - (\tan \alpha + \cot \alpha)x + 1 = 0$ .

If  $C = \sin \alpha \times \cos \alpha$ , find the value of  $C$ .

Let the other root be  $y$ ,  $(2 + \sqrt{3})y = \text{product of roots} = 1 \Rightarrow y = 2 - \sqrt{3}$

$\tan \alpha + \cot \alpha = \text{sum of roots} = 2 + \sqrt{3} + 2 - \sqrt{3} = 4$

$$\frac{\sin \alpha}{\cos \alpha} + \frac{\cos \alpha}{\sin \alpha} = 4 \Rightarrow \frac{1}{\sin \alpha \cos \alpha} = 4 \Rightarrow C = \sin \alpha \times \cos \alpha = \frac{1}{4}$$

**G1.4** Let  $a$  be an integer. If the inequality  $|x + 1| < a - 1.5$  has no integral solution, find the greatest value of  $a$ .

$\because |x + 1| \geq 0$ , In order that the equation has no integral solution, it is sufficient that  $a - 1.5 < 0$

$$a < 1.5$$

Greatest integral value of  $a = 1$

**Group Event 2**


**G2.1** In Figure 1,  $PRS$  is a straight line,  $PQ = PR = QS$  and  $\angle QPR = 30^\circ$ . If  $\angle RQS = w^\circ$ , find the value of  $w$ .

$\angle QPR = \angle QSP = 30^\circ$  (base  $\angle$ s isos.  $\Delta$ )

$\angle PQS = 120^\circ$  ( $\angle$ s sum of  $\Delta$ )

$\angle PQR = \angle PRQ = (180^\circ - 30^\circ) \div 2 = 75^\circ$  ( $\angle$ s sum of isos.  $\Delta$ )

$\angle RQS = 120^\circ - 75^\circ = 45^\circ \Rightarrow w = 45$



**G2.2** Let  $f(x) = px^7 + qx^3 + rx - 5$ , where  $p, q$  and  $r$  are real numbers.

If  $f(-6) = 3$  and  $z = f(6)$ , find the value of  $z$ .

**Reference: 1995 FI1.3**

$$f(-6) = 3 \Rightarrow -p \times 6^7 - q \times 6^3 - 6r - 5 = 3$$

$$f(6) = p \times 6^7 + q \times 6^3 + 6r - 5 = -(-p \times 6^7 - q \times 6^3 - 6r - 5) - 10 = -3 - 10 = -13$$

**G2.3** If  $n \neq 0$  and  $s = \left(\frac{20}{2^{2n+4} + 2^{2n+2}}\right)^{\frac{1}{n}}$ , find the value of  $s$ .

$$s = \left(\frac{20}{16 \cdot 2^{2n} + 4 \cdot 2^{2n}}\right)^{\frac{1}{n}} = \left(\frac{20}{20 \cdot 2^{2n}}\right)^{\frac{1}{n}} = \frac{1}{4}$$

**G2.4** Given that  $x$  and  $y$  are positive integers and  $x + y + xy = 54$ . If  $t = x + y$ , find the value of  $t$ .

**Reference: 2024 FI1.1**

$$1 + x + y + xy = 55$$

$$(1 + x)(1 + y) = 55$$

$$1 + x = 5, 1 + y = 11 \text{ or } 1 + x = 11, 1 + y = 5$$

$$x = 4, y = 10 \text{ or } x = 10, y = 4$$

$$t = 14$$

**Group Event 3**

**G3.1** Given that  $r = 2006 \times \frac{\sqrt{8} - \sqrt{2}}{\sqrt{2}}$ , find the value of  $r$ .

It is easy to show that  $r = 2006$ .

**G3.2** Given that  $6^{x+y} = 36$  and  $6^{x+5y} = 216$ , find the value of  $x$ .

$$x + y = 2 \dots\dots\dots (1)$$

$$x + 5y = 3 \dots\dots\dots (2)$$

$$5(1) - (2): 4x = 7 \Rightarrow x = \frac{7}{4}$$

**G3.3** Given that  $\tan x + \tan y + 1 = \cot x + \cot y = 6$ . If  $z = \tan(x + y)$ , find the value of  $z$ .

$$\tan x + \tan y + 1 = \frac{\tan y + \tan x}{\tan x \tan y} = 6$$

$$\tan x + \tan y = 5; \tan x \tan y = \frac{5}{6}$$

$$\tan(x + y) = \frac{\tan x + \tan y}{1 - \tan x \tan y} = \frac{5}{1 - \frac{5}{6}} = 30$$

**G3.4** In Figure 1,  $ABCD$  is a rectangle,  $F$  is the midpoint of  $CD$  and  $BE : EC = 1 : 3$ . If the area of the rectangle  $ABCD$  is  $12 \text{ cm}^2$  and the area of  $BEFD$  is  $R \text{ cm}^2$ , find the value of  $R$ .

$$\text{Area of } \triangle ABCD = 6 \text{ cm}^2$$

$$\text{Area of } \triangle CEF = \frac{3}{4} \cdot \frac{1}{2} \cdot 6 \text{ cm}^2 = \frac{9}{4} \text{ cm}^2$$

$$\text{Area of } BEFD = \left(6 - \frac{9}{4}\right) \text{ cm}^2 = \frac{15}{4} \text{ cm}^2$$




Figure 1

