

	1	2	2	3	3	30	4	10	5	1
16-17 Individual	6	-6049	7	79	8	$\frac{22}{35}$	9	90°	10	5
	11	44	12	$\frac{1}{42}$	13	20 or $3\sqrt{11}$ <small>see the remark</small>	14	16	15	$4\sqrt{5}$
16-17 Group	1	(8, 10)	2	60	3	45°	4	4	5	10
	6	16	7	4034	8	$\frac{13}{\text{see the remark}}$	9	3	10	11

Individual Events

I1 已知 $A2017B$ 是一個六位數，且可被 72 整除，求 A 的值。

Given that $A2017B$ is a 6-digit number which is divisible by 72, find the value of A .

Reference: 2001 FG1.3, 2003 FI4.1

$72 = 8 \times 9$, the number is divisible by 8 and 9.

$\overline{17B}$ is divisible by 8, i.e. $B = 6$.

$A + 2 + 0 + 1 + 7 + 6 = 9m$, where m is an integer.

$16 + A = 9m$, $A = 2$

$A = 2$

I2 已知 $0 \leq p \leq 1$ ，求 $Q = 3p^2(1-p) + 6p(1-p)^2 + 3(1-p)^3$ 的最大值。

Given that $0 \leq p \leq 1$, find the greatest value of $Q = 3p^2(1-p) + 6p(1-p)^2 + 3(1-p)^3$.

$$Q = 3p^2(1-p) + 6p(1-p)^2 + 3(1-p)^3$$

$$= 3(1-p)(p+1-p)^2$$

$$Q = 3(1-p) \leq 3$$

The maximum value of Q is 3.

I3 已知 ΔABC 的三條邊的長是 a 、 b 和 c ，其中 $3 \leq a \leq 5 \leq b \leq 12 \leq c \leq 15$ ，求當 ΔABC 的面積最大時，它的周界是多少？

Given that the three sides of ΔABC are of lengths a , b and c , where $3 \leq a \leq 5 \leq b \leq 12 \leq c \leq 15$, find the perimeter of ΔABC when its area attains the maximum value.
 c is the longest side.

$$\text{Area} = \frac{1}{2} \cdot ab \sin C \leq \frac{1}{2} \cdot ab \cdot 1 \quad (\text{Equality holds when } a^2 + b^2 = c^2)$$

The largest area is attained when $a = 5$, $b = 12$, $c = 13$

$$\text{Perimeter} = 5 + 12 + 13 = 30$$

I4 設 B 及 C 為正整數，求 C 的最 小 值使得 $B^2 = C + 134$ 。

Let B and C be positive integers. Find the least value of C satisfying $B^2 = C + 134$.

C is the least when B is the least.

$$B^2 = C + 134 \leq 144 = 12^2$$

When $B = 12$, $C = \underline{\underline{10}}$

I5 若把一組自然數之和 $1 + 2 + 3 + \dots + 2015 + 2016 + 2017$ 除以 9，餘數是甚麼？

Determine the remainder when the sum of natural numbers $1 + 2 + 3 + \dots + 2015 + 2016 + 2017$ is divided by 9.

$$\begin{aligned} 1 + 2 + \dots + 2017 &= \frac{1 + 2017}{2} \cdot 2017 \\ &= 1009 \times 2017 \\ &= (112 \times 9 + 1)(224 \times 9 + 1) \\ &= 9m + 1 \end{aligned}$$

The remainder when divided by 9 is 1.

- I6** 已知 $a_0 = 2$, $a_1 = -1$ 及 $a_{n+1} = 2a_n - a_{n-1}$, 其中 $n \geq 1$, 求 a_{2017} 的值。

Given that $a_0 = 2$, $a_1 = -1$ and $a_{n+1} = 2a_n - a_{n-1}$, where $n \geq 1$, determine the value of a_{2017} .

The characteristics equation is $\lambda^2 = 2\lambda - 1 \Rightarrow \lambda = 1$

The general solution is $a_n = (An + B) \cdot 1^n = An + B$

$$a_0 = 0 + B = 2, a_1 = A + 2 = -1 \Rightarrow A = -3$$

$$a_n = 2 - 3n \Rightarrow a_{2017} = 2 - 3 \times 2017 = -6049$$

- I7** 設 N 為完全立方數，已知 $N = 161x + 23y$, 其中 x 和 y 均為正整數。求 $x + y$ 的最小值。

Let N be a perfect cube number. Given that $N = 161x + 23y$, where x and y are positive integers.

Find the minimum value of $x + y$.

$$161x + 23y = 23(7x + y) = m^3$$

$$7x + y = 23^2 = 529 = 7 \times 75 + 4$$

$$x = 75, y = 4$$

$$\text{Minimum value of } x + y = 79$$

- I8** 已知 $\textcircled{2} = 1 \times 2 \times 3 \times 4$, $\textcircled{3} = 2 \times 3 \times 4 \times 5$, $\textcircled{4} = 3 \times 4 \times 5 \times 6$, ... 及 $\frac{1}{\textcircled{15}} - \frac{1}{\textcircled{17}} = \frac{1}{\textcircled{17}} \times A$, 求 A 的值。

Given that $\textcircled{2} = 1 \times 2 \times 3 \times 4$, $\textcircled{3} = 2 \times 3 \times 4 \times 5$, $\textcircled{4} = 3 \times 4 \times 5 \times 6$, ... and $\frac{1}{\textcircled{15}} - \frac{1}{\textcircled{17}} = \frac{1}{\textcircled{17}} \times A$,

find the value of A .

$$\frac{1}{14 \times 15 \times 16 \times 17} - \frac{1}{16 \times 17 \times 18 \times 19} = \frac{1}{16 \times 17 \times 18 \times 19} \times A$$

$$\frac{1}{14 \times 15} - \frac{1}{18 \times 19} = \frac{1}{18 \times 19} \times A$$

$$18 \times 19 - 14 \times 15 = 14 \times 15A$$

$$132 = 210A$$

$$A = \frac{22}{35}$$

- I9** 已知 $\sin x \cdot \cos x = 0$ 及 $\sin^3 x - \cos^3 x = 1$, 其中 $90^\circ \leq x < 180^\circ$, 求 x 的值。

Given that $\sin x \cdot \cos x = 0$ and $\sin^3 x - \cos^3 x = 1$, where $90^\circ \leq x < 180^\circ$, find the value of x .

$\sin x = 0$ or $\cos x = 0$

$x = 0^\circ$ (rejected), 180° (rejected) or 90°

When $x = 90^\circ$, $\sin^3 x - \cos^3 x = 1$


- I10** 如圖一, CM 是 $\angle ACB$ 的角平分線, 且 $AB = 2AC$ 。已知 ΔAMC

的外接圓與 BC 相交於 N 。若 $BN = 10$, 求 AM 的長度。

In Figure 1, CM is the angle bisector of $\angle ACB$ and $AB = 2AC$.

Given that the circumscribed circle of ΔAMC intersects BC at N . If $BN = 10$, find the length of AM .

Let $AC = x$, $AM = y$, then $AB = 2x$, $BM = 2x - y$



圖一 Figure 1

Let $\angle ACM = \angle BCM = \theta$

$\angle MAC = \theta$ (\angle s in the same segment)

$\angle ANM = \theta$ (\angle s in the same segment)

$MN = y$ (sides opposite equal \angle s)

$\Delta BMN \sim \Delta BCA$ (equiangular)

$$\frac{y}{10} = \frac{x}{2x} \Rightarrow AM = y = 5$$

I11 已知 x 為一實數，求 $\sqrt{x(x+3)(x+6)(x+9)+2017}$ 的最小值。

Given that x is a real number, find the least value of $\sqrt{x(x+3)(x+6)(x+9)+2017}$.

Reference 1993 HG6, 1995 FI4.4, 1996 FG10.1, 2000 FG3.1, 2004 FG3.1, 2012 FI2.3, 2013 HI5

$$\begin{aligned}\sqrt{x(x+3)(x+6)(x+9)+2017} &= \sqrt{x(x+9)(x+3)(x+6)+2017} \\ &= \sqrt{(x^2+9x)(x^2+9x+18)+2017} \\ &= \sqrt{(x^2+9x)^2+18(x^2+9x)+9^2+1936} \\ &= \sqrt{(x^2+9x+9)^2+44^2} \geq 44\end{aligned}$$

The minimum value is 44.

I12 已知 $\frac{x}{x^2-5x+1} = \frac{1}{2}$ ，求 $\frac{x^2}{x^4-5x^2+1}$ 的值。

Given $\frac{x}{x^2-5x+1} = \frac{1}{2}$, find the value of $\frac{x^2}{x^4-5x^2+1}$.

$$\frac{x^2-5x+1}{x} = 2 \Rightarrow \frac{x^2+1}{x} - 5 = 2 \Rightarrow x + \frac{1}{x} = 7$$

$$\left(x + \frac{1}{x}\right)^2 = 49 \Rightarrow x^2 + 2 + \frac{1}{x^2} = 49 \Rightarrow x^2 + \frac{1}{x^2} = 47$$

$$x^2 - 5 + \frac{1}{x^2} = 42 \Rightarrow \frac{x^4 - 5x^2 + 1}{x^2} = 42$$

$$\Rightarrow \frac{x^2}{x^4 - 5x^2 + 1} = \frac{1}{42}$$

I13 如圖二， O 是圓 ADB 的圓心。 BC 及 CD 分別是圓形在點 B 及 D 的切線。 $OC \parallel AD$ ， $OA = 15$ 。

若 $AD + OC = 43$ ，求 CD 的長。

As shown in Figure 2, O is the centre of the circle ADB . BC and CD are tangents to the circle at points B and D respectively. $OC \parallel AD$, $OA = 15$.

If $AD + OC = 43$, find the length of CD .

Join OD . $OD \perp DC$ (tangent \perp radius)

Draw $OJ \perp AJ$. $\triangle OAJ \cong \triangle ODJ$ (R.H.S.)

Let $AJ = JD = x$ (corr. sides \cong Δ s), $OC = 43 - 2x$

Let $\angle ODA = \theta$, $\angle COD = \theta$ (alt. \angle s $AD \parallel OC$)

$$\cos \theta = \frac{x}{15} = \frac{15}{43-2x}$$

$$43x - 2x^2 = 225$$

$$2x^2 - 43x + 225 = 0$$

$$(x-9)(2x-25) = 0$$

$$x = 9 \text{ or } 12.5$$

When $x = 9$, $OC = 43 - 2x = 25$


$CD^2 = OC^2 - OD^2 = 25^2 - 15^2$ (Pythagoras' theorem)

$$CD = 20$$

When $x = 12.5$, $OC = 43 - 2x = 18$

$CD^2 = OC^2 - OD^2 = 18^2 - 15^2 = 99$ (Pythagoras' theorem)

$CD = 3\sqrt{11}$ (**Remark:** Candidates give answer with either 20 or $3\sqrt{11}$ will score the mark)



圖二 Figure 2

I14 若 $a + \log_2 b = a^2 + \log_2 b^3 - 10 = 3$ ，其中 $b > 1$ ，求 b 的值。

If $a + \log_2 b = a^2 + \log_2 b^3 - 10 = 3$, where $b > 1$, find the value of b .

$$\log_2 b = 3 - a \dots\dots (1)$$

$$a^2 + \log_2 b^3 - 10 = 3$$

$$a^2 + 3 \log_2 b - 10 = 3$$

Sub. (1) into the equation: $a^2 + 3(3 - a) - 10 = 3$

$$a^2 - 3a - 4 = 0$$

$$(a - 4)(a + 1) = 0$$

$$a = 4 \text{ or } -1$$

Sub. $a = 4$ into (1): $\log_2 b = 3 - 4 = -1 \Rightarrow b = 2^{-1} < 1$ (rejected)

Sub. $a = -1$ into (1): $\log_2 b = 3 + 1 = 4 \Rightarrow b = 2^4 = 16$

I15 在圖三中，已知 $ABCDEF$ 為正六邊形，且它的面積是 $90\sqrt{3}$ ，

求 GJ 的值。

In Figure 3, given that $ABCDEF$ is a regular hexagon and its area is $90\sqrt{3}$, find the length of GJ .

Let O be the centre. Let $AB = a$, OA, OB, OC, OD, OE, OF divides the hexagon $ABCDEF$ into 6 congruent equilateral triangles with sides a .

$$\frac{6}{2} \cdot a^2 \sin 60^\circ = 90\sqrt{3}$$

$$\Rightarrow a^2 = 60$$

$$\Rightarrow a = \sqrt{60}$$


In $\triangle OFG$, $\angle GOF = 30^\circ = \angle GFO$, $OF = AB = a$

$$\frac{a}{2OG} = \cos 30^\circ$$

$$\frac{\sqrt{60}}{2} = \frac{\sqrt{3}}{2} OG$$

$$OG = \sqrt{20} = 2\sqrt{5}$$

$$GJ = 2OG = 4\sqrt{5}$$



圖三 Figure 3

Group Events

G1 設 ΔABC 為一等腰直角三角形，頂點 A 及 B 的座標分別為 $(-2, 0)$ 及 $(18, 0)$ ，且 C 的座標是正數。當 ΔABC 的面積為最小時，求 C 的座標。

Suppose that ΔABC is an isosceles right-angled triangle with the coordinates of the vertices A and B as $(-2, 0)$ and $(18, 0)$, respectively, and the coordinates of C having positive values. Determine the coordinates of C when the area of ΔABC attains its minimum.

When the area of ΔABC attains its minimum, AB is the hypotenuse, $AC = BC$, $AC \perp BC$.

Let M be the mid-point of $AB = (8, 0)$. Let the coordinates of C be $(8, y)$.

$$\frac{y}{8+2} \cdot \frac{y}{8-18} = -1$$

$$y^2 = 100$$

$y = 10$, the coordinates of C is $(8, 10)$.


G2 如圖一所示，點 A 、 B 、 C 、 D 、 E 及 F 均在一直線上。點 G 、 H 、 D 及 I 在另一直線上。揀選三點，可形成多少個三角形？

As shown in Figure 1, points A, B, C, D, E and F lie on the same straight line, and G, H, D and I lie on another straight line. How many triangles can be made by connecting any three points?

Number of triangles without $D = C_2^5 \cdot C_1^3 + C_1^5 \cdot C_2^3 = 45$

Number of triangles with $D = C_1^5 \cdot C_1^3 = 15$

Total number of triangles = $45 + 15 = 60$




圖一 Figure 1

G3 如圖二所示， P 、 Q 分別是正方形 $ABCD$ 的邊 BC 及 CD 上的點。

已知 ΔPCQ 的周界的長等於正方形 $ABCD$ 的周界的長的 $\frac{1}{2}$ ，

求 $\angle PAQ$ 的值。

As shown in Figure 2, P, Q are points on the sides BC and CD of a square $ABCD$. Given that the perimeter of ΔPCQ is $\frac{1}{2}$ of that of the square $ABCD$, find the value of $\angle PAQ$.



圖二 Figure 2

Reference: Dropbox/Data/My%20Web/Home_Page/Geometry/transform/Q5.pdf, 2006 HG7

Let $AB = BC = CD = DA = a$, perimeter of $\Delta PCQ = 2a$

Let $CP = x$, $CQ = y$, $BP = a - x$, $CQ = a - y$, $PQ = 2a - x - y$

Rotate ΔABP about A in clockwise direction by 90° to ΔADE

Then $\Delta ABP \cong \Delta ADE$; $DE = a - x$, $AP = AE$ (corr. sides \cong Δ 's)

$AQ = AQ$ (common side)


$$PQ = 2a - x - y = (a - y) + (a - x) = QE$$

$\therefore \Delta APQ \cong \Delta AEQ$ (S.S.S.)

$\angle PAE = 90^\circ$ (by rotation)

$\angle PAQ = \angle EAQ$ (corr. \angle s. \cong Δ 's)

$\angle PAQ = 45^\circ$



- G4** 在圖三中， O 是圓心。弦 AB 及半徑 OD 的延綫相交於 C 。已知 $OA = 25$ 、 $AB = 30$ 及 $BC = 6$ 。求 CD 的長。

In Figure 3, O is the centre of the circle. Chord AB and radius OD are produced to meet at C . Given that $OA = 25$, $AB = 30$ and $BC = 6$, find the length of CD .

Produce CO to meet the circle again at E . $DE = \text{diameter} = 50$


By intersecting chords theorem, $CB \times CA = CD \times CE$

$$6 \times 36 = CD \times (CD + 50)$$

$$CD^2 + 50CD - 216 = 0$$

$$(CD - 4)(CD + 54) = 0$$

$$CD = 4$$



圖三 Figure 3

- G5** 設 Q 為所有能滿足不等式 $\frac{9p^2}{(\sqrt{3p+1}-1)^2} < 3p+10$ 的整數 p 之和，求 Q 的值。

Let Q be the sum of all integers p satisfying the inequality $\frac{9p^2}{(\sqrt{3p+1}-1)^2} < 3p+10$,

find the value of Q .

$$3p+1 \geq 0 \text{ and } 3p+10 > 0 \text{ and } \sqrt{3p+1}-1 \neq 0 \text{ and } 9p^2 < (3p+10)(3p+1-2\sqrt{3p+1}+1)$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } 9p^2 < (3p+10)(3p+2)-2(3p+10)\sqrt{3p+1}$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } 9p^2 < 9p^2 + 36p + 20 - 2(3p+10)\sqrt{3p+1}$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } (3p+10)\sqrt{3p+1} < 18p + 10$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } (3p+10)^2(3p+1) < (18p+10)^2$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } 27p^3 + 189p^2 + 360p + 100 < 324p^2 + 360p + 100$$

$$p \geq -\frac{1}{3} \text{ and } p \neq 0 \text{ and } 27p^3 - 135p^2 < 0$$

$$p \geq -\frac{1}{3} \text{ and } p < 5$$

$$p = 1, 2, 3 \text{ or } 4$$

$$\text{Sum of all integers } p = 1 + 2 + 3 + 4 = 10$$

- G6** 在圖四中，正方形 $ABCD$ 的邊長為 20。已知 $DK : KA = AH : HB = 1 : 3$ 及 $BK \parallel GD$, $HC \parallel AN$ ，求陰影部分 $PQRS$ 的面積。

In Figure 4, square $ABCD$ has sides of length 20.

Given that $DK : KA = AH : HB = 1 : 3$ and $BK \parallel GD$, $HC \parallel AN$, find the area of shaded region $PQRS$. (Reference 2009 HG6)

$AK = 15 = HB$, $DK = 5 = AH$, $\angle KAB = 90^\circ = \angle HBC$, $AB = BC$
 $\Delta ABK \cong \Delta BCH$ (S.A.S.)


Let $\angle ABK = \theta = \angle BCH$ (corr. \angle 's, $\cong \Delta$'s)

$\angle BHC = 90^\circ - \theta$ (\angle sum of Δ)

In ΔBQH , $\angle BQH = 180^\circ - \theta - (90^\circ - \theta) = 90^\circ$ (\angle sum of Δ)

$\because BK \parallel GD$, $HC \parallel AN$ and $\angle BQH = 90^\circ$

$\therefore PQRS$ is a rectangle



圖四 Figure 4

$$BK = \sqrt{15^2 + 20^2} = 25, \cos \theta = \frac{20}{25} = \frac{4}{5}$$

$$PQ = AH \cos \theta = 5 \times \frac{4}{5} = 4$$

$$PS = DK \cos \theta = 5 \times \frac{4}{5} = 4$$

$$\text{Area of } PQRS = 4 \times 4 = 16$$

G7 已知對於實數 $x_1, x_2, x_3, \dots, x_{2017}$,

$$\sqrt{x_1 - 1} + \sqrt{x_2 - 1} + \sqrt{x_3 - 1} + \dots + \sqrt{x_{2017} - 1} = \frac{1}{2}(x_1 + x_2 + x_3 + \dots + x_{2017}),$$

求 $x_1 + x_2 + x_3 + \dots + x_{2017}$ 的值。

It is given that for real numbers $x_1, x_2, x_3, \dots, x_{2017}$,

$$\sqrt{x_1 - 1} + \sqrt{x_2 - 1} + \sqrt{x_3 - 1} + \dots + \sqrt{x_{2017} - 1} = \frac{1}{2}(x_1 + x_2 + x_3 + \dots + x_{2017}),$$

Find the value of $x_1 + x_2 + x_3 + \dots + x_{2017}$.

$x_1 \geq 1, x_2 \geq 1, \dots, x_{2017} \geq 1$ (otherwise, $\sqrt{x_1 - 1} + \sqrt{x_2 - 1} + \sqrt{x_3 - 1} + \dots + \sqrt{x_{2017} - 1}$ is undefined)

For $1 \leq i \leq 2017$, $\sqrt{x_i - 1} \leq \frac{1}{2}(x_i - 1 + 1) = \frac{1}{2}x_i$ (A.M. \geq G.M., equality holds when $x_i = 2$)

$$\sqrt{x_1 - 1} + \sqrt{x_2 - 1} + \sqrt{x_3 - 1} + \dots + \sqrt{x_{2017} - 1} = \frac{1}{2}(x_1 + x_2 + x_3 + \dots + x_{2017})$$

$$\Rightarrow x_1 = x_2 = \dots = x_{2017} = 2$$

$$x_1 + x_2 + x_3 + \dots + x_{2017} = 2 \times 2017 = 4034$$

G8 設正整數 T 能滿足條件： T 的數字的積是 $T^2 - 11T - 23$ 。求該等正整數之和， S 的值。

Let positive integers, T , satisfy the condition: the product of the digits of T is $T^2 - 11T - 23$.

Find the sum S , of all such positive integers.

Let $y = T^2 - 11T - 23 = (T - 5.5)^2 - 53.25$, y is decreasing for $T < 5.5$, increasing for $T > 5.5$

If $1 \leq T \leq 5$, then $T = T^2 - 11T - 23 < 1^2 - 11 - 23 < 0$, which is impossible

$$y > 0 \Leftrightarrow (T - 5.5)^2 - 53.25 > 0 \Leftrightarrow T - 5.5 > \sqrt{53.25} > \sqrt{42.25} = 6.5 \Leftrightarrow T > 12$$

When $T = 13$, $y = 13^2 - 11 \times 13 - 23 = 3 = 1 \times 3 = \text{product of digits}$

$\therefore T = 13$ is one possible solution

Δ of $y (= T^2 - 11T - 23)$ is $(-11)^2 - 4(-23) = 213$, which is not a perfect square

$\therefore y$ cannot be a composite number

However, $y = T^2 - 11T - 23 = \text{product of its digits of } T$

$\Rightarrow y = 1 \times \text{prime number}$

$\therefore 1 < y = \text{prime number} < 9$

$\therefore y$ is strictly increasing for $T > 5.5$

When $T = 14$, $T^2 - 11T - 23 = 14^2 - 11 \times 14 - 23 = 19$, which is a two-digit number

\therefore There is no solution for $T \geq 14$

\therefore There is only one possible solution $T = 13$ which satisfies $1 < y < 9$

$S = \text{sum of all such positive integers} = 13$

Remark Original version ... product of the digits of $T = T^2 - 11T - 23 \dots$

Somebody will confuse that $T = T^2 - 11T - 23$.

G9 在圖五中， ABC 是一個等邊三角形且與一圓相交於六點： P 、 Q 、 R 、 S 、 T 及 U 。若 $AS = 3$ ， $SR = 13$ ， $RC = 2$ 及 $UT = 8$ ，求 $BP - QC$ 的值。

In Figure 5, ABC is an equilateral triangle intersecting the circle at six points P, Q, R, S, T and U . If $AS = 3$, $SR = 13$, $RC = 2$ and $UT = 8$, find the value of $BP - QC$.

Reference: 2015 HG9

Let $AT = a$, $BU = b$, $BP = x$, $QC = y$, $PQ = 18 - x - y$

By intersecting chord theorem,

$$a(a + 8) = 3 \times (3 + 13)$$

$$a^2 + 8a - 48 = 0$$

$$(a - 4)(a + 12) = 0$$

$a = 4$ or -12 (rejected)

$$AC = 3 + 13 + 2 = 18 = AB = BC$$

$$b = 18 - 4 - 8 = 6$$

$$x(x + 18 - x - y) = 6 \times (6 + 8)$$

$$x(18 - y) = 84 \dots\dots (1)$$

$$y(y + 18 - x - y) = 2 \times (2 + 13)$$

$$y(18 - x) = 30 \dots\dots (2)$$

$$(1) - (2): 18(x - y) = 54$$

$$BP - QC = x - y = 3$$

G10 已知方程 $a^2x^2 - (4a - 3a^2)x + 2a^2 - a - 21 = 0$ (其中 $a > 0$) 最少有一個整數根，求所有 a 的可能整數值之和。

It is given that the equation $a^2x^2 - (4a - 3a^2)x + 2a^2 - a - 21 = 0$ (where $a > 0$) has at least one integral root. Find the sum of all possible integral values of a .

$$\Delta = (4a - 3a^2)^2 - 4a^2(2a^2 - a - 21)$$

$$\Delta = 16a^2 - 24a^3 + 9a^4 - 8a^4 + 4a^3 + 84a^2$$

$$\Delta = a^4 - 20a^3 + 100a^2 = a^2(a - 10)^2$$

$$x = \frac{(4a - 3a^2) \pm \sqrt{a^2(a - 10)^2}}{2a^2}$$

$$= \frac{(4a - 3a^2) \pm a(a - 10)}{2a^2}$$


$$= \frac{(4 - 3a) \pm (a - 10)}{2a}$$

$$x = \frac{-6 - 2a}{2a} \text{ or } \frac{14 - 4a}{2a}$$

$$x = -\frac{3}{a} - 1 \text{ or } \frac{7}{a} - 2$$

$$a = 1, 3, 7$$

$$\text{Sum of all possible integral values of } a = 1 + 3 + 7 = 11$$




圖五 Figure 5

Geometrical Construction

1. 如下圖，已知一圓 O 的其中一條直徑為 AC 。
求作圓上兩點 B 、 D 使得 $ABCD$ 成為一個正方形。

As shown in the figure below, given that O is a circle with a diameter AC . Construct two points B, D on the circle such that $ABCD$ form a square.




作圖方法如下(圖一)：

- (1) 作 AC 的垂直平分線，交圓 O 於 B 及 D 。
- (2) 連接 AB 、 BC 、 CD 及 DA 。

$ABCD$ 便是所需的正方形，作圖完畢。

證明從略。





圖一

2. 已知 $\triangle ABC$ ，如下圖所示。

求作一點 M ，使得 MA 、 MB 及 MC 三條綫段將 $\triangle ABC$ 的面積三等分。

Given $\triangle ABC$ as shown in the figure below. Construct a point M such that the line segments MA , MB , MC will divide the area of $\triangle ABC$ into 3 equal parts.




作圖方法如下(圖二)：

- (1) 作 BC 的垂直平分綫， D 為中點，作 AC 的垂直平分綫， E 為中點，作 AB 的垂直平分綫， F 為中點。
 - (2) 連接中綫 AB 、 BE 及 CF ，交於形心 M 。
- MA 、 MB 及 MC 三條綫段將 $\triangle ABC$ 的面積三等分，作圖完畢。
證明從略。


3. 已知 A 、 B 兩點和直綫 ℓ ，如下圖所示。求作一圓過 A 、 B 兩點且與 ℓ 相切。

Given two points A , B and a straight line ℓ as shown in the figure below. Construct a circle which passes through A and B , and is tangent to the straight line ℓ .

Reference: C:\Users\twhung.CLSMSS.002\Dropbox\Data\My Web\Home Page\Geometry\7 Construction by ruler and compasses\circle\circle_through_A_B_touch_L.pdf



圖三



圖四

作圖方法如下(圖三、圖四及圖五)：

- (1) 連接 AB ，其延長綫交 L 於 D 。
- (2) 作 AB 的垂直平分線，交 L 於 G ， H 為 AB 的中點。
- (3) 以 G 為圓心， GA 為半徑作一圓。(圖四)
- (4) 作 GD 的垂直平分線， M 為 GD 的中點。
- (5) 以 M 為圓心， MG 為半徑作一圓，交步驟(3)的圓於 E 。
- (6) 連接 EG 、 DE 。(圖)
- (7) 以 D 為圓心， DE 為半徑作一圓，交 L 於 F (在 D 與 G 之間)及 C (在 GD 之延長部分)。
- (8) 過 F 作一綫段垂直於 L ，且交 GH 的延綫於 O ，過 C 作一綫段垂直於 L ，且交 GH 的延長綫於 Q 。
- (9) 以 O 為圓心， OA 為半徑作一圓；以 Q 為圓心， QA 為半徑作一圓。(圖五)

作圖完畢，證明如下：

$$\angle AHG = \angle BHG = 90^\circ \text{ (由作圖所得)}$$

$$GH = GH \quad (\text{公共邊})$$

$$AH = HB \quad (\text{由作圖所得})$$

$$\therefore \Delta AGH \cong \Delta BGH \quad (\text{S.A.S.})$$

$$GA = GB \quad (\text{全等三角形的對應邊})$$

\therefore 步驟(3)的圓經過 A 、 B 。

利用相同方法，可證明步驟(9)的二圓皆經過 A 、 B 。

$$\angle GED = 90^\circ \quad (\text{半圓上的圓周角})$$

$\therefore DE$ 切步驟(3)的圓於 E 。

(切線垂直於半徑的逆定理)

$$DA \times DB = DE^2 \quad (\text{相交弦定理})$$

$$\therefore DE = DF = DC \quad (\text{半徑})$$

$$\therefore DA \times DB = DF^2 \text{ 及 } DA \times DB = DC^2$$

$$\therefore DF$$
 切圓 ABF 於 F 及 DC 切圓 ABC 於 C

(相交弦定理的逆定理)




圖 2