

Integral Cyclic Quadrilateral

Created by Mr. Francis Hung

Last updated: December 26, 2024

It is given that a triangle with one angle is θ , where $\cos \theta = \frac{3}{5}$. If all sides are integers, find all solutions.

$$c^2 = a^2 + b^2 - 2ab \cos \theta \Rightarrow c^2 = a^2 + b^2 - 2ab \times \frac{3}{5}$$

$$5c^2 = 5a^2 - 10ab + 5b^2 + 4ab$$

$$5[c^2 - (a - b)^2] = 4ab$$

$$\frac{5(c+a-b)}{b} = \frac{4a}{c-a+b} = k$$

$$c+a-b = \frac{bk}{5} \dots\dots(1) \text{ and } c-a+b = \frac{4a}{k} \dots\dots(2)$$

$$[(1)-(2)] \div 2: a-b = \frac{bk}{10} - \frac{2a}{k} = \frac{bk^2 - 20a}{10k}$$

$$10ka - 10kb = bk^2 - 20a$$

$$(10k+20)a = (k^2 + 10k)b$$

$$\frac{a}{b} = \frac{k^2 + 10k}{10k + 20}$$

Let $a = (k^2 + 10k)p$, $b = (10k + 20)p$

$$c = b - a + \frac{bk}{5} = [10k + 20 - k^2 - 10k + (2k + 4)k]p = (k^2 + 4k + 20)p. \text{ Let } p = 1.$$

k	$a = k^2 + 10k$	$b = 10k + 20$	$c = k^2 + 4k + 20$	Remark
1	11	30	25	
2	24	40	32	3-4-5
3	39	50	41	
4	56	60	52	13-14-15

It is given that a triangle with one angle is θ , where $\cos \theta = \frac{4}{5}$. If all sides are integers, find all solutions.

$$c^2 = a^2 + b^2 - 2ab \cos \theta \Rightarrow c^2 = a^2 + b^2 - 2ab \times \frac{4}{5}$$

$$5c^2 = 5a^2 - 10ab + 5b^2 + 2ab$$

$$5[c^2 - (a - b)^2] = 2ab$$

$$\frac{5(c+a-b)}{b} = \frac{2a}{c-a+b} = k$$

$$c+a-b = \frac{bk}{5} \dots\dots(1) \text{ and } c-a+b = \frac{2a}{k} \dots\dots(2)$$

$$[(1)-(2)] \div 2: a-b = \frac{bk}{10} - \frac{a}{k} = \frac{bk^2 - 10a}{10k}$$

$$10ka - 10kb = k^2b - 10a$$

$$(10k+10)a = (k^2 + 10k)b$$

$$\frac{a}{b} = \frac{k^2 + 10k}{10k + 10}$$

Let $a = (k^2 + 10k)p$, $b = (10k + 10)p$

$$c = b - a + \frac{bk}{5} = [10k + 10 - k^2 - 10k + (2k + 2)k]p = (k^2 + 2k + 10)p. \text{ Let } p = 1.$$

k	$a = k^2 + 10k$	$b = 10k + 10$	$c = k^2 + 2k + 10$	Remark
1	11	20	13	
2	24	30	18	3-4-5
3	39	40	25	
4	56	50	34	17-25-28
5	75	60	45	3-4-5
6	96	70	58	29-35-48

It is given that a triangle with one angle is θ , where $\cos \theta = \frac{7}{25}$. If all sides are integers, find all solutions.

$$c^2 = a^2 + b^2 - 2ab \cos \theta \Rightarrow c^2 = a^2 + b^2 - 2ab \times \frac{7}{25}$$

$$25c^2 = 25a^2 - 50ab + 25b^2 + 36ab$$

$$25[c^2 - (a-b)^2] = 36ab$$

$$\frac{25(c+a-b)}{b} = \frac{36a}{c-a+b} = k$$

$$c+a-b = \frac{bk}{25} \dots\dots(1) \text{ and } c-a+b = \frac{36a}{k} \dots\dots(2)$$

$$[(1)-(2)] \div 2: a-b = \frac{bk}{50} - \frac{18a}{k} = \frac{bk^2 - 900a}{50k}$$

$$50ka - 50kb = k^2b - 900a$$

$$(50k+900)a = (k^2+50k)b$$

$$\frac{a}{b} = \frac{k^2+50k}{50k+900}$$

Let $a = (k^2 + 50k)p$, $b = (50k + 900)p$

$$c = b - a + \frac{bk}{25} = [50k + 900 - k^2 - 50k + (2k + 36)k]p = (k^2 + 36k + 900)p. \text{ Let } p = 1.$$

k	$a = k^2 + 50k$	$b = 50k + 900$	$c = k^2 + 36k + 900$	Remark
1	51	950	937	
2	104	1000	976	13-122-125
3	159	1050	1017	
4	216	1100	1060	54-275-265
5	275	1150	1105	55-221-230
6	336	1200	1152	7-24-25
7	399	1250	1201	
8	464	1300	1252	116-313-325
9	531	1350	1305	59-145-150
10	600	1400	1360	15-34-35
11	671	1450	1417	
12	744	1500	1476	62-123-125
13	819	1550	1537	
14	896	1600	1600	14-25-25
15	975	1650	1665	65-110-111
16	1056	1700	1732	264-425-433
17	1139	1750	1801	
18	1224	1800	1872	17-25-26
19	1311	1850	1945	
20	1400	1900	2020	70-95-101

It is given that the three sides of a triangle are positive integers.

Let θ be an acute angle such that $\sin \theta = \frac{p}{r}$, $\cos \theta = \frac{q}{r}$ and $\tan \theta = \frac{p}{q}$ are rational numbers in the simplest forms.

Then (p, q, r) is a primitive Pythagorean triple. $(p, q, r) = (2uv, u^2 - v^2, u^2 + v^2)$ or $(u^2 - v^2, 2uv, u^2 + v^2)$.

Let ABC be a triangle ($BC = a$, $AC = b$, $AB = c$). $\angle ACB = \theta$.

Area of $\Delta ABC = \frac{1}{2}ab \sin C = \frac{1}{2}bc \sin A = \frac{1}{2}ca \sin B$, which must be rational.

Denote the area of ΔABC by S , then $\sin A = \frac{2S}{bc}$, $\sin B = \frac{2S}{ca}$ and $\sin C = \frac{2S}{ab}$ which are also rational.

By cosine formula, $\cos A = \frac{b^2 + c^2 - a^2}{2bc}$, $\cos B = \frac{c^2 + a^2 - b^2}{2ac}$, $\cos C = \frac{a^2 + b^2 - c^2}{2ab} \Rightarrow \cos A, \cos B, \cos C \in \mathbb{Q}$.

$\tan A = \frac{\sin A}{\cos A}$, $\tan B = \frac{\sin B}{\cos B}$, $\tan C = \frac{\sin C}{\cos C}$ which are still rational. Find all integral solutions for a , b and c .

$$c^2 = a^2 + b^2 - 2ab \cos C \Rightarrow c^2 = a^2 + b^2 - 2ab \times \frac{q}{r}$$

Case 1 $q = 2uv$

$$(u^2 + v^2)c^2 = (u^2 + v^2)a^2 + (u^2 + v^2)b^2 - 4abuv$$

$$(u^2 + v^2)[c^2 - (a^2 - 2ab + b^2)] = -4abuv + 2ab(u^2 + v^2)$$

$$(u^2 + v^2)(c + a - b)(c - a + b) = 2ab(u - v)^2$$

$$\frac{(u^2 + v^2)(c + a - b)}{b} = \frac{2(u - v)^2 a}{c - a + b} = k$$

$$c + a - b = \frac{bk}{u^2 + v^2} \dots\dots(1) \text{ and } c - a + b = \frac{2(u - v)^2 a}{k} \dots\dots(2)$$

$$(1) - (2): 2(a - b) = \frac{bk}{u^2 + v^2} - \frac{2(u - v)^2 a}{k}$$

$$2(a - b)(u^2 + v^2)k = bk^2 - 2a(u - v)^2(u^2 + v^2)$$

$$2(u^2 + v^2)ka + 2(u - v)^2(u^2 + v^2)a = 2(u^2 + v^2)kb + k^2b$$

$$\frac{a}{b} = \frac{(2u^2 + 2v^2 + k)k}{2(u^2 + v^2)[(u - v)^2 + k]} \Rightarrow a = (2u^2 + 2v^2 + k)k, b = 2(u^2 + v^2)[(u - v)^2 + k].$$

$$\begin{aligned} \text{Sub. into (2): } c &= (2u^2 + 2v^2 + k)k - 2(u^2 + v^2)[(u - v)^2 + k] + 2(u - v)^2(2u^2 + 2v^2 + k) \\ &= k^2 + (2u^2 + 2v^2)k - 2(u^2 + v^2)(u - v)^2 - 2(u^2 + v^2)k + 4(u - v)^2(u^2 + v^2) + 2(u - v)^2k \\ &= k^2 + 2(u - v)^2k + 2(u - v)^2(u^2 + v^2) = k^2 + 2(u - v)^2(u^2 + v^2 + k) \end{aligned}$$

k	u	v	$a = (2u^2 + 2v^2 + k)k$	$b = 2(u^2 + v^2)[(u - v)^2 + k]$	$c = k^2 + 2(u - v)^2(u^2 + v^2 + k)$	$\cos C = \frac{2uv}{u^2 + v^2}$
1	2	1	11	20	13	$\frac{4}{5}$
1	3	1	21	100	89	$\frac{6}{10} = \frac{3}{5}$
1	3	2	27	52	29	$\frac{12}{13}$
1	4	1	35	340	325	$\frac{8}{17}$
1	4	3	51	100	53	$\frac{24}{25}$
3	2	1	39	40	25	$\frac{4}{5}$

Note that $u > v$ are relatively prime positive integers. $\cos C = \frac{2uv}{u^2 + v^2} > 0$ and so $\angle C$ must be acute.

$$\text{Area of } \Delta ABC = \frac{1}{2} (2u^2 + 2v^2 + k)k \cdot 2(u^2 + v^2)[(u-v)^2 + k] \cdot \frac{u^2 - v^2}{u^2 + v^2} = k(u^2 - v^2)(2u^2 + 2v^2 + k)[(u-v)^2 + k]$$

$$\text{Case 2 } p = 2uv, \cos \theta = \frac{q}{r} = \frac{u^2 - v^2}{u^2 + v^2}$$

$$c^2 = a^2 + b^2 - 2ab \cos C \Rightarrow c^2 = a^2 + b^2 - 2ab \cdot \frac{u^2 - v^2}{u^2 + v^2}$$

$$(u^2 + v^2)c^2 = (u^2 + v^2)(a^2 - 2ab + b^2) + 2ab(u^2 + v^2) - 2ab(u^2 - v^2)$$

$$(u^2 + v^2)(c + a - b)(c - a + b) = 4v^2 ab$$

$$\frac{(u^2 + v^2)(c + a - b)}{b} = \frac{4av^2}{c - a + b} = k$$

$$c + a - b = \frac{bk}{u^2 + v^2} \dots\dots (1) \text{ and } c - a + b = \frac{4av^2}{k} \dots\dots (2)$$

$$(1) - (2): 2(a - b) = \frac{bk}{u^2 + v^2} - \frac{4av^2}{k}$$

$$2k(u^2 + v^2)a - 2k(u^2 + v^2)b = k^2b - 4v^2(u^2 + v^2)a$$

$$2(u^2 + v^2)(k + 2v^2)a = [k^2 + 2(u^2 + v^2)k]b$$

$$\frac{a}{b} = \frac{k^2 + 2(u^2 + v^2)k}{2(u^2 + v^2)(k + 2v^2)} \Rightarrow a = k^2 + 2(u^2 + v^2)k, b = 2(u^2 + v^2)(k + 2v^2)$$

$$\text{Sub. into (2): } c = k^2 + 2(u^2 + v^2)k - 2(u^2 + v^2)(k + 2v^2) + 4v^2[k + 2(u^2 + v^2)] \\ = k^2 + 4v^2(u^2 + v^2) + 4v^2k = k^2 + 4v^2(u^2 + v^2 + k)$$

k	u	v	$a = (2u^2 + 2v^2 + k)k$	$b = 2(u^2 + v^2)(2v^2 + k)$	$c = k^2 + 4v^2(u^2 + v^2 + k)$	$\cos C = \frac{u^2 - v^2}{u^2 + v^2}$
1	1	2	11	90	97	$-\frac{3}{5}$
1	2	1	11	30	25	$\frac{3}{5}$
1	2	3	27	494	505	$-\frac{5}{13}$
1	3	2	27	234	225	$\frac{5}{13}$
1	4	1	35	102	73	$\frac{15}{17}$
1	4	3	51	950	937	$\frac{7}{25}$
3	2	1	39	50	41	$\frac{3}{5}$

Note that u, v are distinct relatively prime positive integers.

If $u > v$, then $\cos C = \frac{u^2 - v^2}{u^2 + v^2} > 0$ and so C is acute. If $u < v$, then $\cos C = \frac{u^2 - v^2}{u^2 + v^2} < 0$ and so C is obtuse.

$$\text{Area of } \Delta ABC = \frac{1}{2} ab \sin C = \frac{1}{2} (2u^2 + 2v^2 + k)k \cdot 2(u^2 + v^2)(2v^2 + k) \cdot \frac{2uv}{u^2 + v^2}$$

$$= 2kuv(2u^2 + 2v^2 + k)(2v^2 + k), \text{ which is an integer}$$


Given that the three sides of a triangle ACB are positive integers. $\angle ACB = \theta$ such that $\sin \theta = \frac{p}{r}$, $\cos \theta = \frac{q}{r}$ and $\tan \theta = \frac{p}{q}$ are all rational numbers. If (p, q, r) is a Pythagorean triple, from the above analysis,

$$(a, b, c) = ((2u^2 + 2v^2 + k)k, 2(u^2 + v^2)[(u-v)^2 + k], k^2 + 2(u-v)^2(u^2 + v^2 + k)) \text{ or } ((2u^2 + 2v^2 + k)k, 2(u^2 + v^2)(2v^2 + k), k^2 + 4v^2(u^2 + v^2 + k))$$

where $k \in \mathbb{Z}^+$, u, v are distinct relatively prime positive integers.

In this section, we are going to find a cyclic quadrilateral for which all sides and all diagonals are integers.

Idea: In February 2008, I asked Dr. Man Keung Siu from the University of Hong Kong about how to find a solution to the above question. He quoted a paper "Normal Trigrade and cyclic quadrilateral with integral sides and diagonals" from April, 1951 American Mathematical Monthly. I didn't understand the content of the paper. However, the author gave an example of integral cyclic quadrilateral $ABCD$, $AB = 52$, $BC = 60$, $CD = 39$, $DA = 25$, $AC = 56$, $BD = 63$.



It takes me years of time to investigate how to find another integral cyclic quadrilateral. Still, I failed to find any other solutions. After retirement, I use cosine formula to find $\cos \angle ACB$.

$$\cos \angle ACB = \frac{56^2 + 60^2 - 52^2}{2 \times 56 \times 60} = \frac{3}{5} \text{ and so } \sin \angle ACB = \frac{4}{5}, \tan \angle ACB = \frac{4}{3} \text{ which are all rational.}$$

I started to investigate a triangle with integral sides and rational trigonometric ratios of each angle.

Suppose the diagonals intersect at Q . Let $\angle BQC = \theta$.

$$\text{It can be proved that } \tan \theta = \frac{4\sqrt{(s-a)(s-b)(s-c)(s-d)}}{a^2 + c^2 - b^2 - d^2},$$

where $a = 52$, $b = 60$, $c = 39$, $d = 25$, $s = \frac{1}{2}(a+b+c+d)$.

$$a^2 + c^2 - b^2 - d^2 = 52^2 + 39^2 - 60^2 - 25^2 = 0 \Rightarrow \text{denominator} = 0 \Rightarrow \theta = 90^\circ$$

$$CQ = BC \cos \angle BCQ = 60 \times \frac{3}{5} = 36, BQ = BC \sin \angle BCQ = 48. \Delta BCQ \text{ is a 3-4-5 } \Delta.$$

$$\therefore DQ = BD - BQ = 63 - 48 = 15, AQ = AC - CQ = 56 - 36 = 20.$$

ΔABQ is a 5-12-13 Δ . ΔADQ is a 3-4-5 Δ . ΔCDQ is a 5-12-13 Δ .

This is a **special case** of integral cyclic quadrilateral.



Will there be any other example(s) of integral cyclic quadrilateral $ABCD$ with perpendicular diagonals?

$\angle ACB = \angle ADB, \angle CAD = \angle CBD, \angle ACD = \angle ABD, \angle BDC = \angle BAC$ (\angle s in the same segment)



$\therefore \Delta ABQ \sim \Delta DCQ, \Delta BCQ \sim \Delta ADQ$ (equiangular)

We try to find **two pairs of right-angled triangles with common sides**.

If ΔABQ is a 5-12-13 Δ , ΔBCQ is a 8-15-17 Δ . $AQ = 12m, BQ = 5m, AB = 13m, BQ = 8n, CQ = 15n, BC = 17n$.



$BC = 5m = 8n$, let $m = 8, n = 5$, then $AB = 13 \times 8 = 104, AQ = 12 \times 8 = 96, BC = 17 \times 5 = 85, CQ = 15 \times 5 = 75$



Construct the circumscribed circle ABC . Extend BQ to cut the circumscribed circle at D .

$\Delta DCQ \sim \Delta ABQ$ which are 5-12-13 Δ s. $DQ = \frac{75}{5} \times 12 = 180, CD = \frac{75}{5} \times 13 = 195$.

$\Delta ADQ \sim \Delta BCQ$ which are 8-15-17 Δ s. $DQ = \frac{96}{8} \times 15 = 180, AD = \frac{96}{8} \times 17 = 204$.


$\therefore ABCD$ is another integral cyclic quadrilateral with

$AB = 104, BC = 85, CD = 195, DA = 204, AC = 96 + 75 = 171, BD = 40 + 180 = 220$.


Question:

Can we find an integral cyclic quadrilateral $ABCD$ so that the diagonals are not necessarily perpendicular?

We see from page 3 the last line that 25-39-40 is an integral triangle ΔABQ .



Using B as centre, BA as radius, draw a circular arc AC . Extend AQ , cutting the circular arc at C . Join BC .



Using B as centre, BQ as radius, draw a circular arc, cutting AQ at P . $BP = BQ = 25$ (radii)

Let M be the foot of perpendicular from B to PQ . $\Delta BPM \cong \Delta BQM$ (R.H.S.). Join BM .

$$AM = AB \cos \angle BAQ = 40 \times \frac{4}{5} = 32, QM = AQ - AM = 39 - 32 = 7 = PM \text{ (corr. sides, } \cong \Delta \text{s)}$$

$$AP = AQ - QM - PM = 39 - 7 - 7 = 25$$


$$\angle BAC = \angle BCA, \angle BPQ = \angle BQP \text{ (base } \angle \text{s, isos. } \Delta \text{)}$$

$$\angle APB = 180^\circ - \angle BPM = 180^\circ - \angle BQM = \angle BQC \text{ (adj. } \angle \text{s on st. line)}$$

$$\Delta ABP \cong \Delta CBQ \text{ (A.A.S.)}$$

$$CQ = AP = 25 \text{ (corr. sides, } \cong \Delta \text{s)}$$

Construct a circumscribed circle through A, B and C . Extend BQ to cut the circle again at D . Join AD and CD .



It is easy to show that $\Delta ABQ \cong \Delta DCQ$ (A.A.S.)

$$DQ = AQ = 39, DC = AB = 40 \text{ (corr. sides, } \cong \Delta \text{s)}$$

$$\Delta ADQ \sim \Delta BCQ \text{ (equiangular)}$$


$$\frac{AD}{BC} = \frac{AQ}{BQ} \text{ (corr. sides, } \sim \Delta \text{s)} \Rightarrow AD = 40 \times \frac{39}{25} = \frac{312}{5}$$

Multiply every side by 5 to give integral sides. $AB = BC = CD = 200, AD = 312, BD = AC = 320$.


Again, this is a **special case for three equal adjacent sides of integral cyclic quadrilateral**.

Second example:

We see from page 3 the first line in the table that 11-13-20 is an integral triangle ΔABQ .



Using B as centre, BA as radius, draw a circular arc AC . Extend AQ , cutting the circular arc at C . Join BC .



Using B as centre, BQ as radius, draw a circular arc, cutting AQ at P . $BP = BQ = 13$ (radii)

Let M be the foot of perpendicular from B to PQ . $\Delta BPM \cong \Delta BQM$ (R.H.S.). Join BM .

$$AM = AB \cos \angle BAQ = 20 \times \frac{4}{5} = 16, QM = AM - AQ = 16 - 11 = 5 = PM \text{ (corr. sides, } \cong \Delta \text{s)}$$


$$AP = AM + PM = 16 + 5 = 21$$

$$\angle BAC = \angle BCA, \angle BPQ = \angle BQP \text{ (base } \angle \text{s, isos. } \Delta)$$

$$\Delta ABP \cong \Delta CBQ \text{ (A.A.S.)}$$

$$CQ = AP = 21 \text{ (corr. sides, } \cong \Delta \text{s)}$$

Construct a circumscribed circle through A, B and C . Extend BQ to cut the circle again at D . Join AD and CD .



It is easy to show that $\Delta ABQ \sim \Delta DCQ$ (equiangular)

$$\frac{DQ}{AQ} = \frac{CQ}{BQ} = \frac{CD}{AB} \text{ (corr. sides, } \sim \Delta \text{s)}$$

$$DQ = 11 \times \frac{21}{13} = \frac{231}{13}, CQ = 20 \times \frac{21}{13} = \frac{420}{13}, AC = 11 + 21 = 32, BD = 13 + \frac{231}{13} = \frac{400}{13}$$

$\Delta ADQ \sim \Delta BCQ$ (equiangular)


$$\frac{AD}{BC} = \frac{AQ}{BQ} \text{ (corr. sides, } \sim \Delta \text{s)}$$

$$AD = 20 \times \frac{11}{13} = \frac{220}{13}$$


Multiply every side by 13 to give integral sides. $AB = BC = 260, CD = 420, AD = 220, AC = 416, BD = 400$. Again, this is a **special case for two equal adjacent sides of integral cyclic quadrilateral**.

Question: Can find an integral cyclic quadrilateral $ABCD$ so that all adjacent sides are unequal?

We see from page 3 the first line in the table that 11-13-20 is an integral triangle ΔABQ .



We find another triangle ΔBCQ so that $BC = q$, $QC = b$ and A, Q, C are collinear.



Apply cosine rule on ΔBCQ : $q^2 = b^2 + 11^2 + 22b \times \frac{4}{5}$

$$5q^2 = 5(b^2 + 22b + 11^2) - 110b + 88b$$

$$22b = 5[(b+11)^2 - q^2] = 5(b+q+11)(b-q+11)$$

$$\frac{b+q+11}{b} = \frac{22}{5(b-q+11)} = t$$

$$b+q+11 = bt \dots\dots (1), \quad b-q+11 = \frac{22}{5t} \dots\dots (2)$$

$$(1) + (2): 2b + 22 = bt + \frac{22}{5t}$$

$$10tb + 110t = 5t^2b + 22$$

$$110t - 22 = (5t^2 - 10t)b$$

$$b = \frac{22(5t-1)}{5t(t-2)}$$

$$(1) - (2): 2q = bt - \frac{22}{5t} = \frac{22t(5t-1)}{5t(t-2)} - \frac{22}{5t}$$


$$q = 11 \cdot \frac{t(5t-1) - (t-2)}{5t(t-2)} = 11 \cdot \frac{5t^2 - 2t + 2}{5t(t-2)}$$

$$\text{Put } t = 3, b = \frac{22 \times 14}{15} = \frac{308}{15}, q = 11 \times \frac{41}{15} = \frac{451}{15}$$

Multiply every side by 15 to give integral sides.

$$AB = 13 \times 15 = 195, BQ = 11 \times 15 = 165, AQ = 20 \times 15 = 300.$$

Construct a circumscribed circle through A , B and C . Extend BQ to cut the circle again at D . Join AD and CD .



It is easy to show that $\triangle ABQ \sim \triangle DCQ$ (equiangular)


$$\frac{DQ}{AQ} = \frac{CQ}{BQ} = \frac{CD}{AB} \quad (\text{corr. sides, } \sim\Delta\text{s})$$

$$DQ = 300 \times \frac{308}{165} = 560, CD = 195 \times \frac{308}{165} = 364, AC = 300 + 308 = 608, BD = 165 + 560 = 725$$

$\triangle ADQ \sim \triangle BCQ$ (equiangular)

$$\frac{AD}{BC} = \frac{AQ}{BQ} \quad (\text{corr. sides, } \sim\Delta\text{s})$$

$$AD = 451 \times \frac{300}{165} = 820$$



Question: Can we find a general formula for integral cyclic quadrilateral for which the diagonals are not necessarily perpendicular and the adjacent sides are not necessarily equal?

Let the cyclic quadrilateral be $ABCD$. The diagonals AC and BD intersect at Q .

Let $AQ = a, BQ = b, AB = c, DQ = d, CQ = e, CD = f, AD = g, BC = h$ as shown in the figure.

$\Delta ABQ \sim \Delta DCQ$ (equiangular)

$$\frac{a}{d} = \frac{b}{e} = \frac{c}{f} = m, \text{ where } m \text{ is a constant (corr. sides, } \sim \Delta \text{s)}$$

$$a = dm, b = em, c = fm$$

$\Delta ADQ \sim \Delta BCQ$ (equiangular)

$$\frac{a}{b} = \frac{d}{e} = \frac{g}{h} = n, \text{ where } n \text{ is a constant. (corr. sides, } \sim \Delta \text{s)}$$

$$a = bn = dm, d = en, g = hn$$

There are five variables e, f, h, m, n in the figure.

Let ΔBCQ be an obtuse-angled triangle with $\angle BQC > 90^\circ$.



Then ΔCDQ is an acute-angled triangle.

By the formula on Page 4, the only possible solution for ΔCDQ is:

$$CD = k^2 + 4v^2(u^2 + v^2 + k), CQ = (2u^2 + 2v^2 + k)k, DQ = 2(u^2 + v^2)(2v^2 + k)$$

$$\cos \angle CQD = \frac{u^2 - v^2}{u^2 + v^2} > 0,$$

where $u > v$ are distinct relatively prime positive integers.



Again, ΔBCQ is another triangle adjacent to ΔCDQ with $\cos \angle BQC = \frac{v^2 - u^2}{u^2 + v^2} < 0$

The roles of u and v are interchanged. $BC = h = k^2 + 4u^2(u^2 + v^2 + k), BQ = em = 2(u^2 + v^2)(2u^2 + k)$


$$n = \frac{en}{e} = \frac{DQ}{CQ} = \frac{2(u^2 + v^2)(2v^2 + k)}{(2u^2 + 2v^2 + k)k}$$

$$\begin{aligned} AQ &= emn = BQ \times n \\ &= 2(u^2 + v^2)(2u^2 + k) \cdot \frac{2(u^2 + v^2)(2v^2 + k)}{(2u^2 + 2v^2 + k)k} \\ &= 4 \cdot \frac{(u^2 + v^2)^2(2u^2 + k)(2v^2 + k)}{(2u^2 + 2v^2 + k)k} \end{aligned}$$

$$AD = hn = [k^2 + 4u^2(u^2 + v^2 + k)] \cdot \frac{2(u^2 + v^2)(2v^2 + k)}{(2u^2 + 2v^2 + k)k}$$

$$m = \frac{em}{e} = \frac{BQ}{CQ} = \frac{2(u^2 + v^2)(2u^2 + k)}{(2u^2 + 2v^2 + k)k}$$

$$AB = fm = [k^2 + 4v^2(u^2 + v^2 + k)] \cdot \frac{2(u^2 + v^2)(2v^2 + k)}{(2u^2 + 2v^2 + k)k}$$



Multiply every side by $(2u^2 + 2v^2 + k)k$ to give integral sides:

$$CQ = (2u^2 + 2v^2 + k)^2 k^2$$

$$CD = (2u^2 + 2v^2 + k)k[k^2 + 4v^2(u^2 + v^2 + k)]$$

$$DQ = 2k(u^2 + v^2)(2v^2 + k)(2u^2 + 2v^2 + k)$$

$$BC = (2u^2 + 2v^2 + k)k[k^2 + 4u^2(u^2 + v^2 + k)]$$

$$BQ = 2(u^2 + v^2)(2u^2 + k)(2u^2 + 2v^2 + k)k$$

$$AQ = 4(u^2 + v^2)^2(2u^2 + k)(2v^2 + k)$$

$$AD = 2(u^2 + v^2)(2v^2 + k)[k^2 + 4u^2(u^2 + v^2 + k)]$$

$$AB = 2(u^2 + v^2)(2u^2 + k)[k^2 + 4v^2(u^2 + v^2 + k)]$$

For example, take $k = 1, u = 2, v = 1$:

$$CQ = 121$$

$$CD = 11 \times (1 + 4 \times 6) = 275$$

$$DQ = 2 \times 5 \times 3 \times 11 = 330$$

$$BC = 11 \times (1 + 16 \times 6) = 1067$$

$$BQ = 2 \times 5 \times 9 \times 11 = 990$$


$$AQ = 4 \times 5^2 \times 9 \times 3 = 2700$$

$$AD = 2 \times 5 \times 3 \times (1 + 16 \times 6) = 2910$$

$$AB = 2 \times 5 \times 9 \times (1 + 4 \times 6) = 2250$$


$$AC = AQ + CQ = 2700 + 121 = 2821$$

$$BD = BQ + DQ = 990 + 330 = 1320$$




Think further: From the document: https://twhung78.github.io/Number_Theory/120triangle.pdf, we know that 3-5-7 is a 120° triangle, whereas 5-7-8 is a 60° triangle.

Combine the common side '5' to give a bigger triangle as shown:



Construct a circumscribed circle through B, C and D . Extend CQ to cut the circle again at A . Join AB and AD .



It is easy to show that $\Delta ABQ \sim \Delta DCQ$ (equiangular)

$$\frac{DQ}{AQ} = \frac{CQ}{BQ} = \frac{CD}{AB} \quad (\text{corr. sides, } \sim \Delta s)$$

$$AQ = 3 \times \frac{8}{5} = \frac{24}{5}, AB = 7 \times \frac{8}{5} = \frac{56}{5}, AC = \frac{24}{5} + 5 = \frac{49}{5}, BD = 8 + 3 = 11$$

$\Delta ADQ \sim \Delta BCQ$ (equiangular)

$$\frac{AD}{BC} = \frac{AQ}{BQ} \quad (\text{corr. sides, } \sim \Delta s)$$

$$AD = 7 \times \frac{3}{5} = \frac{21}{5}$$

Multiply every side by 5 to give integral sides. $BC = CD = 35, AD = 21, AB = 56, AC = 49, BD = 55$.


\therefore We can construct another integral cyclic quadrilateral with a simpler formula, but the area of each smaller triangle inside (and hence the cyclic quadrilateral) are not integers.

Again, this is a **special case for two equal adjacent sides of integral cyclic quadrilateral and the angle between the two diagonals is 60°** .

Question: Given any triangle ΔABQ with integral sides, can we construct an integral cyclic quadrilateral, while the angle between the diagonals are not necessarily 60° using a similar method?

Let ΔABC be a 2-3-4 triangle with $AB = 4$, $BQ = 2$, $CQ = 3$.

We can construct (method on page 8) another triangle ΔBCQ (with common sides BQ) so that $BC = 4$, $CQ = 4$.



$$\cos \angle AQB = \frac{2^2 + 3^2 - 4^2}{2 \times 2 \times 3} = -\frac{1}{4}$$

$$\cos \angle BQC = \frac{2^2 + 4^2 - 4^2}{2 \times 2 \times 4} = \frac{1}{4} = -\cos \angle AQB$$

$$\angle AQB + \angle BQC = 180^\circ$$



$$\therefore A, Q, C \text{ are collinear.}$$

Construct a circumscribed circle through A, B and C . Extend BQ to cut the circle again at D . Join AD and CD .

Using a similar method, we can prove that $DQ = 6$, $AD = 6$, $CD = 8$.

In this case, $AB = BC = 4$

(two equal adjacent sides, whereas the angle between the diagonals $\neq 60^\circ$.)



Given any triangle ΔABQ with integral sides ($AB = c$, $BQ = a$, $AQ = b$). We can use the method on page 9 to find another triangle ΔBCQ with a common side BQ and A, Q, C are collinear. $BC = c$, $CQ = x$.

$$\cos \angle AQB = \frac{a^2 + b^2 - c^2}{2ab}, \cos \angle BQC = -\frac{a^2 + b^2 - c^2}{2ab}$$


$$\text{If } \angle AQB \text{ is obtuse, then } x = b + 2a \cos \angle BQC = b - \frac{a^2 + b^2 - c^2}{2ab} = \frac{c^2 - a^2}{b}.$$

$$\text{If } \angle AQB \text{ is acute, then } x = b - 2a \cos \angle AQB = b - \frac{a^2 + b^2 - c^2}{2ab} = \frac{c^2 - a^2}{b}.$$

$$\text{If } \angle AQB = 90^\circ, \text{ then } x = b = \frac{c^2 - a^2}{b}.$$

Finally, a cyclic quadrilateral with integral sides is formed.

The only necessary condition is $c > a$ and a, b, c obey triangle inequality.



In a more general case, let $CQ = x$, $BC = y$.

$$y^2 = x^2 + a^2 - 2ax \cos \angle BQC = x^2 + a^2 + 2ax \cos \angle AQB = x^2 + a^2 + 2ax \cdot \frac{a^2 + b^2 - c^2}{2ab}$$

$$y^2 = x^2 + 2ax + a^2 + \frac{x}{b}(a^2 + b^2 - c^2) - 2ax$$

$$(x+y+a)(x-y+a) = \frac{x}{b}[c^2 - (a^2 - 2ab + b^2)] = \frac{x}{b}[c^2 - (a-b)^2]$$

$$\frac{x+y+a}{x} = \frac{c^2 - (a-b)^2}{b(x-y+a)} = k$$



$$x+y+a = kx \dots\dots (1), x-y+a = \frac{c^2 - (a-b)^2}{bk} \dots\dots (2)$$

$$(1) + (2): 2x+2a = kx + \frac{c^2 - (a-b)^2}{bk} \Rightarrow x = \frac{2abk + (a-b)^2 - c^2}{bk(k-2)}$$

$$(1) - (2): 2y = kx + \frac{(a-b)^2 - c^2}{bk} = \frac{2abk + (a-b)^2 - c^2}{b(k-2)} + \frac{(a-b)^2 - c^2}{bk}$$

$$y = \frac{ak}{k-2} - \frac{k-1}{bk(k-2)}[c^2 - (a-b)^2]$$

Multiply every side by $abk(k-2)$, construct a circumscribed circle through ABC , extend BQ to the circle at D .



$$\frac{DQ}{a[2abk + (a-b)^2 - c^2]} = \frac{ab^2k(k-2)}{a^2bk(k-2)} = \frac{AD}{a\{abk^2 - (k-1)[c^2 - (a-b)^2]\}} \quad (\text{corr. sides, } \Delta ADQ \sim \Delta BCQ)$$

$$DQ = b[2abk + (a-b)^2 - c^2], AD = b\{abk^2 - (k-1)[c^2 - (a-b)^2]\}$$

$$\frac{DQ}{ab^2k(k-2)} = \frac{CD}{abck(k-2)} \quad (\text{corr. sides, } \Delta CDQ \sim \Delta BAQ)$$

$$CD = c[2abk + (a-b)^2 - c^2]$$

Necessary condition: a, b, c obey triangle inequality and $k > 2$.

The following is the feedback from Doctor M. K. Siu, retired professor from the University of Hong Kong:
We know quite well how to compute the diagonals of a cyclic quadrilateral from the four sides. Further analysis follows. However, it is always good to try to solve a problem by oneself. The pleasure does not lie so much in the solution than in the process of solving it.

Author's email: twhung78-hkame@yahoo.com

Reference:

1. Pythagorean Triple: https://twhung78.github.io/Number_Theory/Pythagorean_triple.pdf
2. Angle between two diagonals in a cyclic quadrilateral:
https://twhung78.github.io/Geometry/6%20Circles/2%20Cyclic%20quadrilateral/Angle_diagonals_cyclic_quadrilateral.pdf
3. “Normal Trigrade and cyclic quadrilateral with integral sides and diagonals” from April, 1951 American Mathematical Monthly.
4. 120° triangle: https://twhung78.github.io/Number_Theory/120triangle.pdf