

# Divisible by 11

Created by Mr. Francis Hung

Last updated: July 1, 2023

Given a 5-digits integer  $x = \overline{abcde}$ .

If  $a + c + e - (b + d) = 11k$ , where  $k$  is an integer, prove that  $x$  is divisible by 11.

$$\begin{aligned}\text{Proof: } x &= 10000a + 1000b + 100c + 10d + e \\ &= 9999a + a + 1001b - b + 99c + c + 11d - d + e \\ &= 11(909a + 91b + 9c + d) + a - b + c - d + e \\ &= 11(900a + 100b + 9c + d) + 11k, \text{ which is divisible by 11.}\end{aligned}$$

## HKHLE General Mathematics 1976 Q8(b)

Prove that, for any positive integer  $n$ , the integer  $10^n + (-1)^{n-1}$  is divisible by 11.

Hence deduce a necessary and sufficient condition for an integer to be divisible by 11 by considering only the sum and difference of the digits of the integer.

Induction on  $n$ .

$n = 1$ ,  $10^1 + (-1)^0 = 11$  which is obviously divisible by 11.

Suppose  $10^k + (-1)^{k-1} = 11m$ , where  $m$  is an integer, for some positive integer  $k$ .

$$10^{k+1} + (-1)^k = 10(10^k) + (-1)^k = 10[11m - (-1)^{k-1}] + (-1)^k = 110m + (-1)^k(1 + 10) = 11[10m + (-1)^k]$$

which is divisible by 11.

So, by M.I.,  $10^n + (-1)^{n-1}$  is divisible by 11 for any positive integer  $n$ .

The necessary and sufficient condition is: Let  $S_1$  be the sum of all odd digits of an integer  $N$ ,  $S_2$  be the sum of all even digits of  $N$ .  $S_1 - S_2$  is divisible by 11 if and only if  $N$  is divisible by 11.

Proof: Let  $N = a_n \times 10^n + a_{n-1} \times 10^{n-1} + \dots + a_1 \times 10 + a_0$ , where  $0 \leq a_r \leq 9$  and  $a_r$  are integers,  $0 \leq r \leq n$ .

$$S_1 - S_2 = (-1)^n a_n + (-1)^{n-1} a_{n-1} + \dots - a_1 + a_0$$

If  $N$  is divisible by 11,  $a_n \times 10^n + a_{n-1} \times 10^{n-1} + \dots + a_1 \times 10 + a_0 = 11m$ , where  $m$  is an integer.

$$\begin{aligned}N &= 11m = a_n \times [10^n + (-1)^{n-1}] + a_{n-1} \times [10^{n-1} + (-1)^{n-2}] + \dots + a_1 \times [10 + 1] + a_0(1 - 1) \\ &\quad - [(-1)^{n-1} a_n + (-1)^{n-2} a_{n-1} + \dots + a_1 - a_0] \\ &= a_n \times 11k_n + a_{n-1} \times 11k_{n-1} + \dots + a_1 \times 11k_1 + a_0 \cdot 11k_0 + S_1 - S_2, \text{ where } k_r \text{ are integers, } 0 \leq r \leq n \\ \Rightarrow S_1 - S_2 &= 11m - [a_n \times 11k_n + a_{n-1} \times 11k_{n-1} + \dots + a_1 \times 11k_1 + a_0 \cdot 11k_0] \\ &= 11[m - a_n k_n + a_{n-1} k_{n-1} + \dots + a_1 k_1 + a_0 k_0], \text{ which is divisible by 11.}\end{aligned}$$

If  $S_1 - S_2$  is divisible by 11, then  $(-1)^n a_n + (-1)^{n-1} a_{n-1} + \dots - a_1 + a_0 = 11m$ , where  $m$  is an integer.

$$\begin{aligned}N &= a_n \times 10^n + a_{n-1} \times 10^{n-1} + \dots + a_1 \times 10 + a_0 \\ &= a_n \times [10^n + (-1)^{n-1}] + a_{n-1} \times [10^{n-1} + (-1)^{n-2}] + \dots + a_1 \times [10 + 1] + a_0(1 - 1) \\ &\quad + [(-1)^n a_n + (-1)^{n-1} a_{n-1} + \dots - a_1 + a_0] \\ &= a_n \times 11k_n + a_{n-1} \times 11k_{n-1} + \dots + a_1 \times 11k_1 + a_0 \cdot 11k_0 + S_1 - S_2 \\ &= 11[a_n k_n + a_{n-1} k_{n-1} + \dots + a_1 k_1 + a_0 k_0] + 11m, \text{ which is divisible by 11.}\end{aligned}$$

The problem is solved.