

# Compound angled formulae

Created by Mr. Francis Hung on 20090205

Last updated: March 19, 2023

## The formula $\sin(A + B) = \sin A \cos B + \cos A \sin B$

$0^\circ < A < 90^\circ, 0^\circ < B < 90^\circ$


Consider a triangle  $KLN$  with a right angle at  $M$ , as shown in the figure.

Let  $\angle LKM = A, \angle MKN = B$ , then  $\angle LKN = A + B$ .

$$\text{Area of } \triangle LKM = \frac{1}{2} pq \sin A$$

$$\text{Area of } \triangle MKN = \frac{1}{2} qr \sin B$$

$$\text{Area of } \triangle LKN = \frac{1}{2} pr \sin(A + B)$$



$$\text{Area of } \triangle LKM + \text{Area of } \triangle MKN = \text{Area of } \triangle LKN$$

$$\therefore \frac{1}{2} pq \sin A + \frac{1}{2} qr \sin B = \frac{1}{2} pr \sin(A + B)$$

$$\sin(A + B) = \frac{q}{r} \sin A + \frac{q}{p} \sin B$$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

**Method 2**  $0^\circ < A < 90^\circ, 0^\circ < B < 90^\circ, 0^\circ < A + B < 90^\circ$

In  $\triangle PQN$ ,  $\angle PNQ = 90^\circ$ ,  $\angle NPQ = A$

$PQRS$  is a rectangle.  $PR = r, PQ = q, PN = p$ .

$\angle RPQ = B, \angle RPN = A + B$ .

$RL \perp PN, KQ \perp RL$ .

$\angle KQP = A$  (alt.  $\angle$ s  $KQ \parallel PN$ )

$\angle RQK = 90^\circ - A$

$\angle KRQ = A$  ( $\angle$  sum of  $\Delta$ )


$KQNL$  is a rectangle

$RL = RK + KL = RQ \cos A + QN$

$$r \sin(A + B) = r \sin B \cos A + q \sin A$$

$$r \sin(A + B) = r \sin B \cos A + r \cos B \sin A$$

$$\therefore \sin(A + B) = \sin A \cos B + \cos A \sin B$$




**Method 3** In  $\triangle ABC$ ,  $A + B + C = 180^\circ$

$$\text{By sine rule, } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = 2R \Rightarrow \sin A = \frac{a}{2R}, \sin B = \frac{b}{2R}, \sin C = \frac{c}{2R}$$

$$\text{By cosine rule, } \cos A = \frac{b^2 + c^2 - a^2}{2bc}; \cos B = \frac{a^2 + c^2 - b^2}{2ac}$$

$$\begin{aligned} \sin A \cos B + \cos A \sin B &= \frac{a}{2R} \cdot \frac{a^2 + c^2 - b^2}{2ac} + \frac{b^2 + c^2 - a^2}{2bc} \cdot \frac{b}{2R} \\ &= \frac{2c^2}{4cR} = \frac{c}{2R} = \sin C = \sin(180^\circ - (A + B)) (\angle \text{s sum of } \Delta) \\ &= \sin(A + B) \end{aligned}$$

### The formula $\sin(A - B) = \sin A \cos B - \cos A \sin B$



Suppose  $PQRS$  is a semi-circle, with diameter  $PQ = d$ ,  $\angle QPS = A$ ,  $\angle QPR = B$ .

$\angle PRQ = 90^\circ$ ,  $\angle PSQ = 90^\circ$  ( $\angle$  in semi-circle)

$$QR = d \sin B$$

$$QS = d \sin A$$

$$PR = d \cos B$$

$$PS = d \cos A$$

$$\angle RPS = A - B$$

By Sine rule on  $\Delta PRS$ ,  $\frac{RS}{\sin(A - B)} = \frac{PS}{\sin \angle PRS}$

$$\frac{RS}{\sin(A - B)} = \frac{PS}{\sin \angle PQS} \quad (\because \angle PRS = \angle PQS, \angle s in the same segment)$$

$$\frac{RS}{\sin(A - B)} = \frac{d \cos A}{\sin(90^\circ - A)} \quad (\because \angle PQS = 90^\circ - \alpha, \angle s sum of \Delta PQS)$$

$$RS = d \sin(A - B) \quad (\because \sin(90^\circ - A) = \cos A)$$

By Ptolemy's theorem, i.e.  $PR \cdot QS = RS \cdot PQ + QR \cdot PS$

$$d \cos B \cdot d \sin A = d \sin(A - B) \cdot d + d \sin B \cdot d \cos A$$

$$\Rightarrow \sin(A - B) = \sin A \cos B - \cos A \sin B$$

### Method 2 New Trend Additional Mathematics Volume One (2002) p. 142

Consider a triangle  $KLN$  with a right angle at  $N$ , as shown in the figure.

Let  $\angle LKN = A$ ,  $\angle MKN = B$ , then  $\angle LKM = A - B$ .

$$\text{Area of } \Delta LKM = \frac{1}{2} qr \sin(A - B)$$

$$\text{Area of } \Delta MKN = \frac{1}{2} pq \sin B$$


$$\text{Area of } \Delta LKN = \frac{1}{2} pr \sin A$$

Area of  $\Delta LKM$  + Area of  $\Delta MKN$  = Area of  $\Delta LKN$


$$\therefore \frac{1}{2} qr \sin(A - B) + \frac{1}{2} pq \sin B = \frac{1}{2} pr \sin A$$

$$\sin(A - B) = \frac{p}{q} \sin A - \frac{p}{r} \sin B$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$



### The formula $\cos(A - B) = \cos A \cos B + \sin A \sin B$



Draw a unit circle with centre  $O$  and radius 1. Suppose  $P(x_1, y_1), Q(x_2, y_2)$  are two points on the circumference. Suppose  $OP$  makes an angle  $A$  with positive  $x$ -axis.  $OQ$  makes an angle  $B$  with positive axis.  $\angle QOP = A - B$ .

$$x_1 = \cos A, y_1 = \sin A; x_2 = \cos B, y_2 = \sin B$$

$$\text{By cosine rule, } PQ^2 = OP^2 + OQ^2 - 2OP \cdot OQ \cos \angle POQ.$$

$$(x_1 - x_2)^2 + (y_1 - y_2)^2 = 1 + 1 - 2 \cos(A - B)$$

$$(\cos A - \cos B)^2 + (\sin A - \sin B)^2 = 1 + 1 - 2 \cos(A - B)$$

$$\cos^2 A - 2 \cos A \cos B + \cos^2 B + \sin^2 A - 2 \sin A \sin B + \sin^2 B = 2 - 2 \cos(A - B)$$

$$-2(\cos A \cos B + \sin A \sin B) = -2 \cos(A - B)$$

$$\therefore \cos(A - B) = \cos A \cos B + \sin A \sin B \dots\dots (1)$$

**Variation** We have already proved the formula  $\cos(A - B) = \cos A \cos B + \sin A \sin B$ , where  $A$  and  $B$  can be positive or negative real numbers.

Replace  $B$  by  $-B$ , then  $\cos(A + B) = \cos A \cos(-B) + \sin A \sin(-B)$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B \dots\dots (2)$$

$$\begin{aligned} \text{Replace } A \text{ by } 90^\circ - A, \text{ then } \sin(A - B) &= \cos[90^\circ - (A - B)] = \cos[(90^\circ - A) + B] \\ &= \cos(90^\circ - A) \cos B - \sin(90^\circ - A) \sin B \end{aligned}$$

Hence,

$$\sin(A - B) = \sin A \cos B - \cos A \sin B \dots\dots (3)$$

Replace  $B$  by  $-B$  again, then  $\sin(A + B) = \sin A \cos(-B) - \cos A \sin(-B)$

$$\sin(A + B) = \sin A \cos B + \cos A \sin B \dots\dots (4)$$

$$\begin{aligned} \text{Now, } \tan(A + B) &= \frac{\sin(A + B)}{\cos(A + B)} = \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B} \\ &= \frac{\sin A \cos B + \cos A \sin B}{\cos A \cos B - \sin A \sin B} \\ &= \frac{\cos A \cos B}{\cos A \cos B - \sin A \sin B} \\ &\therefore \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B} \dots\dots (5) \end{aligned}$$

$$\text{Replace } B \text{ by } -B \text{ again, then } \tan(A - B) = \frac{\tan A + \tan(-B)}{1 - \tan A \tan(-B)} = \frac{\tan A - \tan B}{1 + \tan A + \tan B} \dots\dots (6)$$

**Example 1** Find the value of  $\sin 15^\circ$ ,  $\cos 15^\circ$  and  $\tan 15^\circ$ .

$$\sin 15^\circ = \sin(60^\circ - 45^\circ) = \sin 60^\circ \cos 45^\circ - \cos 60^\circ \sin 45^\circ = \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} - \frac{1}{2} \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{6} - \sqrt{2}}{4}$$

$$\cos 15^\circ = \cos(60^\circ - 45^\circ) = \cos 60^\circ \cos 45^\circ + \sin 60^\circ \sin 45^\circ = \frac{1}{2} \cdot \frac{\sqrt{2}}{2} + \frac{\sqrt{3}}{2} \cdot \frac{\sqrt{2}}{2} = \frac{\sqrt{6} + \sqrt{2}}{4}$$

$$\tan 15^\circ = \tan(60^\circ - 45^\circ) = \frac{\tan 60^\circ - \tan 45^\circ}{1 + \tan 60^\circ \tan 45^\circ} = \frac{\sqrt{3} - 1}{1 + \sqrt{3}} = \frac{(\sqrt{3} - 1)(\sqrt{3} + 1)}{(\sqrt{3} + 1)(\sqrt{3} - 1)} = \frac{3 - 2\sqrt{3} + 1}{3 - 1} = 2 - \sqrt{3}$$

**Example 2** Find the value of  $\sin 75^\circ$ ,  $\cos 75^\circ$  and  $\tan 75^\circ$ .

$$\sin 75^\circ = \cos 15^\circ = \frac{\sqrt{6} + \sqrt{2}}{4} \quad (\because \cos(90^\circ - \theta) = \sin \theta)$$

$$\cos 75^\circ = \sin 15^\circ = \frac{\sqrt{6} - \sqrt{2}}{4}$$

$$\tan 75^\circ = \frac{1}{\tan 15^\circ} = \frac{1}{2 - \sqrt{3}} = \frac{1}{2 - \sqrt{3}} \cdot \frac{2 + \sqrt{3}}{2 + \sqrt{3}} = 2 + \sqrt{3}$$

**Example 3** Find the value of  $\sin 165^\circ$ ,  $\cos 165^\circ$  and  $\tan 165^\circ$ .

$$\sin 165^\circ = \sin(180^\circ - 15^\circ) = -\sin 15^\circ = -\frac{\sqrt{6} - \sqrt{2}}{4}$$

$$\cos 165^\circ = \cos(180^\circ - 15^\circ) = -\cos 15^\circ = -\frac{\sqrt{6} + \sqrt{2}}{4}$$

$$\tan 165^\circ = \tan(180^\circ - 15^\circ) = -\tan 15^\circ = -(2 - \sqrt{3})$$

Using these results, we can deduce the values of the sines, cosines and tangents of  $15^\circ$ ,  $75^\circ$ ,  $105^\circ$ ,  $165^\circ$ ,  $195^\circ$ ,  $225^\circ$ ,  $285^\circ$  and  $345^\circ$ , or equivalently in radians  $\frac{\pi}{12}, \frac{5\pi}{12}, \frac{7\pi}{12}, \frac{11\pi}{12}, \frac{13\pi}{12}, \frac{17\pi}{12}, \frac{19\pi}{12}, \frac{23\pi}{12}$ .

**Example 4** If  $A + B = 45^\circ$ , prove that  $(1 + \tan A)(1 + \tan B) = 2$ .

Hence find the value of  $(1 + \tan 1^\circ)(1 + \tan 2^\circ) \cdots (1 + \tan 44^\circ)(1 + \tan 45^\circ)$ .

$$\tan(A + B) = \tan 45^\circ = 1$$

$$\frac{\tan A + \tan B}{1 - \tan A \tan B} = 1$$

$$\tan A + \tan B = 1 - \tan A \tan B$$

$$1 + \tan A + \tan B + \tan A \tan B = 2$$

$$(1 + \tan A)(1 + \tan B) = 2$$

$$(1 + \tan 1^\circ)(1 + \tan 44^\circ) = 2$$

$$(1 + \tan 2^\circ)(1 + \tan 43^\circ) = 2$$

..... (there are 22 pairs of  $(1 + \tan A)(1 + \tan B)$ )

$$1 + \tan 45^\circ = 2$$

$$(1 + \tan 1^\circ)(1 + \tan 2^\circ) \cdots (1 + \tan 44^\circ)(1 + \tan 45^\circ) = 2^{23}.$$

$$\cot(A + B) = \frac{1}{\tan(A + B)} = \frac{1 - \tan A \tan B}{\tan A + \tan B} = \frac{\cot A \cot B - 1}{\cot A + \cot B} \quad \dots \dots (7)$$

$$\cot(A - B) = \frac{\cot A \cot B + 1}{\cot B - \cot A} \quad \dots \dots (8)$$

**Example 5 (Identity)** If  $A + B + C = 180^\circ$ , prove that  $\cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2}$

$$\text{L.H.S.} = \cot \frac{A}{2} + \cot \frac{B}{2} + \cot \frac{C}{2}$$

$$= \cot \frac{A}{2} + \cot \frac{B}{2} + \tan(90^\circ - \frac{C}{2}) = \cot \frac{A}{2} + \cot \frac{B}{2} + \tan \frac{180^\circ - C}{2}$$

$$= \cot \frac{A}{2} + \cot \frac{B}{2} + \tan\left(\frac{A}{2} + \frac{B}{2}\right) = \cot \frac{A}{2} + \cot \frac{B}{2} + \frac{\tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}}$$

$$= \frac{\cot \frac{A}{2} + \cot \frac{B}{2} - \tan \frac{B}{2} - \tan \frac{A}{2} + \tan \frac{A}{2} + \tan \frac{B}{2}}{1 - \tan \frac{A}{2} \tan \frac{B}{2}} = \cot \frac{A}{2} \cot \frac{B}{2} \cdot \frac{\cot \frac{A}{2} + \cot \frac{B}{2}}{\cot \frac{A}{2} \cot \frac{B}{2} - 1}$$

$$= \cot \frac{A}{2} \cot \frac{B}{2} \cdot \frac{1}{\cot \frac{A+B}{2}} = \cot \frac{A}{2} \cot \frac{B}{2} \cdot \tan \frac{A+B}{2} = \cot \frac{A}{2} \cot \frac{B}{2} \cdot \tan\left(90^\circ - \frac{C}{2}\right)$$

$$= \cot \frac{A}{2} \cot \frac{B}{2} \cot \frac{C}{2} = \text{R.H.S.}$$

**Example 6** If  $\sin A + \cos B = p$ ,  $\cos A - \sin B = q$  and  $A - B = 30^\circ$ , prove that  $p^2 + q^2 - 3 = 0$

$$\sin A = p - \cos B$$

$$\cos A = q + \sin B$$

$$\therefore 1 = (p - \cos B)^2 + (q + \sin B)^2$$

$$p^2 + q^2 + 2q \sin B - 2p \cos B = 0$$

$$p^2 + q^2 + 2[(\cos A - \sin B) \sin B - (\sin A + \cos B) \cos B] = 0$$

$$p^2 + q^2 - 2 + 2(\sin B \cos A - \cos B \sin A) = 0$$

$$p^2 + q^2 - 2 + 2 \sin(B - A) = 0$$

$$p^2 + q^2 - 3 = 0 \quad (\because A - B = 30^\circ)$$

**Example 7** Evaluate  $\tan \theta \tan (\theta - 60^\circ) + \tan \theta \tan (\theta + 60^\circ) + \tan (\theta - 60^\circ) \tan (\theta + 60^\circ)$

Expression =  $\tan \theta [\tan (\theta - 60^\circ) + \tan (\theta + 60^\circ)] + \tan (\theta - 60^\circ) \tan (\theta + 60^\circ)$

$$\begin{aligned} &= \tan \theta \cdot \left( \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta} + \frac{\tan \theta + \sqrt{3}}{1 - \sqrt{3} \tan \theta} \right) + \frac{\tan \theta - \sqrt{3}}{1 + \sqrt{3} \tan \theta} \times \frac{\tan \theta + \sqrt{3}}{1 - \sqrt{3} \tan \theta} \\ &= \tan \theta \cdot \frac{\tan \theta - \sqrt{3} - \sqrt{3} \tan^2 \theta + 3 \tan \theta + \tan \theta + \sqrt{3} + \sqrt{3} \tan^2 \theta + 3 \tan \theta}{1 - 3 \tan^2 \theta} + \frac{\tan^2 \theta - 3}{1 - 3 \tan^2 \theta} \\ &= \tan \theta \cdot \frac{8 \tan \theta}{1 - 3 \tan^2 \theta} + \frac{\tan^2 \theta - 3}{1 - 3 \tan^2 \theta} = \frac{9 \tan^2 \theta - 3}{1 - 3 \tan^2 \theta} \\ &= \frac{3(3 \tan^2 \theta - 1)}{1 - 3 \tan^2 \theta} = -3 \end{aligned}$$

**Example 8** Prove that  $\tan n\theta - \tan(n-1)\theta = \tan \theta [ 1 + \tan n\theta \tan (n-1)\theta ]$

Use this formula to find  $\sum_{r=1}^n \tan r\theta \tan(r-1)\theta$ .

$$\tan \theta = \tan[n\theta - (n-1)\theta] = \frac{\tan n\theta - \tan(n-1)\theta}{1 + \tan n\theta \tan(n-1)\theta}$$

$$\therefore \tan n\theta - \tan(n-1)\theta = \tan \theta [ 1 + \tan n\theta \tan (n-1)\theta ]$$

$$\begin{aligned} \sum_{r=1}^n \tan r\theta \tan(r-1)\theta &= \frac{1}{\tan \theta} \sum_{r=1}^n \tan \theta [ 1 + \tan r\theta \tan(r-1)\theta ] - n \\ &= \frac{1}{\tan \theta} \sum_{r=1}^n [\tan r\theta - \tan(r-1)\theta] - n = \frac{\tan n\theta}{\tan \theta} - n \end{aligned}$$

**Example 9** Solve  $12 \sin \theta - 5 \cos \theta = 13$  for  $0^\circ \leq \theta \leq 360^\circ$ .

$$(12 \sin \theta - 5 \cos \theta)^2 = 169$$

$$144 \sin^2 \theta - 120 \sin \theta \cos \theta + 25 \cos^2 \theta = 169(\sin^2 \theta + \cos^2 \theta)$$

$$25 \sin^2 \theta + 120 \sin \theta \cos \theta + 144 \cos^2 \theta = 0$$

$$25 \tan^2 \theta + 120 \tan \theta + 144 = 0$$

$$(5 \tan \theta + 12)^2 = 0$$

$$\tan \theta = -\frac{12}{5}$$

$$\theta = 112.6^\circ, 292.6^\circ$$

Check: when  $\theta = 112.6^\circ$ ,

$$\text{LHS} = 12 \sin 112.6^\circ - 5 \cos 112.6^\circ = 13 = \text{RHS}$$

When  $\theta = 292.6^\circ$ ,

$$\text{LHS} = 12 \sin 292.6^\circ - 5 \cos 292.6^\circ = -13 \neq \text{RHS}$$

$$\therefore \theta = 112.6^\circ \text{ only}$$

**Classwork 1** Solve the equation  $3 \sin \theta - 2 \cos \theta = 1$  for  $0^\circ \leq \theta \leq 360^\circ$ .  $[\theta = 49.79^\circ \text{ or } 197.59^\circ]$

**Example 10** The point  $D$  divides the sides  $BC$  of  $\Delta ABC$  internally so that  $BD : DC = m : n$ . If  $\angle BAD = \alpha$ ,  $\angle CAD = \beta$  and  $\angle CDA = \theta$ , prove that  $m \cot \alpha - n \cot \beta = (m + n) \cot \theta = n \cot B - m \cot C$ .

$$\text{In } \Delta ADC, \frac{n}{\sin \beta} = \frac{AC}{\sin \theta} \dots (1)$$

$$\text{In } \Delta ABC, \frac{m+n}{\sin(\alpha+\beta)} = \frac{AC}{\sin B}$$

$\angle B = \theta - \alpha$  (ext.  $\angle$  of  $\Delta ABD$ )

$$\frac{m+n}{\sin(\alpha+\beta)} = \frac{AC}{\sin(\theta-\alpha)} \dots (2)$$

$$(1) \div (2) \quad \frac{n}{\sin \beta} \times \frac{\sin(\alpha+\beta)}{(m+n)} = \frac{AC}{\sin \theta} \times \frac{\sin(\theta-\alpha)}{AC}$$

$$m \sin \theta \sin(\alpha + \beta) = (m+n) \sin(\theta - \alpha) \sin \beta$$

$$m \sin \theta \sin(\alpha + \beta) = (m+n) \sin \beta (\sin \theta \cos \alpha - \cos \theta \sin \alpha)$$

$$(m+n) \sin \beta \cos \theta \sin \alpha = (m+n) \sin \beta \sin \theta \cos \alpha - m \sin \theta \sin(\alpha + \beta)$$

$$(m+n) \sin \alpha \sin \beta \cos \theta = [(m+n) \sin \beta \cos \alpha - m \sin(\alpha + \beta)] \sin \theta$$

$$(m+n) \sin \alpha \sin \beta \cos \theta = [(m+n) \cos \alpha \sin \beta - m \sin \alpha \cos \beta - m \cos \alpha \sin \beta] \sin \theta$$

$$(m+n) \sin \alpha \sin \beta \cos \theta = (n \cos \alpha \sin \beta - m \sin \alpha \cos \beta) \sin \theta$$

$$\therefore (m+n) \cot \theta = \frac{m \cos \alpha \sin \beta - n \sin \alpha \cos \beta}{\sin \alpha \sin \beta}$$

$$= m \cot \alpha - n \cot \beta$$

$$\text{In } \Delta ABD, \frac{m}{\sin \alpha} = \frac{AB}{\sin(180^\circ - \theta)}$$

$\alpha = \theta - B$  (ext.  $\angle$  of  $\Delta ABD$ )

$$\therefore \frac{m}{\sin(\theta-B)} = \frac{AB}{\sin \theta} \dots (3)$$

$$\text{In } \Delta ABC, \frac{m+n}{\sin A} = \frac{AB}{\sin C}$$

$$\frac{m+n}{\sin(180^\circ - (B+C))} = \frac{AB}{\sin C}$$

$$\frac{m+n}{\sin(B+C)} = \frac{AB}{\sin C} \dots (4)$$

$$(3) \div (4) \quad \frac{m}{\sin(\theta-B)} \times \frac{\sin(B+C)}{(m+n)} = \frac{AB}{\sin \theta} \times \frac{\sin C}{AB}$$

$$m \sin \theta \sin(B+C) = (m+n) \sin C \sin(\theta - B)$$

$$m \sin \theta \sin(B+C) = (m+n) \sin C (\sin \theta \cos B - \cos \theta \sin B)$$

$$(m+n) \sin C \cos \theta \sin B = (m+n) \sin C \sin \theta \cos B - m \sin \theta \sin(B+C)$$


$$(m+n) \sin B \sin C \cos \theta = [(m+n) \cos B \sin C - m \sin(B+C)] \sin \theta$$

$$(m+n) \sin B \sin C \cos \theta = [(m+n) \cos B \sin C - m \sin B \cos C - m \cos B \sin C] \sin \theta$$

$$(m+n) \sin B \sin C \cos \theta = (n \cos B \sin C - m \sin B \cos C) \sin \theta$$

$$\therefore (m+n) \cot \theta = \frac{n \cos B \sin C - m \sin B \cos C}{\sin B \sin C}$$

$$\therefore (m+n) \cot \theta = n \cot B - m \cot C$$



## Subsidiary angles

### Example 11

- (a) Rewrite the expression  $4 \cos \theta - 3 \sin \theta$  in the form  $r \cos(\theta + \alpha)$ , where  $r > 0$  and  $0 \leq \alpha < \frac{\pi}{2}$ .

Give your answer correct to 2 decimal places.

- (b) Find the maximum and minimum values of  $4 \cos \theta - 3 \sin \theta$ .

$$\begin{aligned} r \cos(\theta + \alpha) &= r (\cos \theta \cos \alpha - \sin \theta \sin \alpha) \\ &= (r \cos \alpha) \cos \theta - (r \sin \alpha) \sin \theta \end{aligned}$$

As  $4 \cos \theta - 3 \sin \theta = (r \cos \alpha) \cos \theta - (r \sin \alpha) \sin \theta$

$r \cos \alpha = 4$  and  $r \sin \alpha = 3$

$$r = \sqrt{3^2 + 4^2} = 5$$

$$\tan \alpha = \frac{3}{4}$$

$\alpha = 0.64$  radians (correct to 2 decimal places)

$$\therefore 4 \cos \theta - 3 \sin \theta = 5 \cos(\theta + 0.64)$$


$$\begin{aligned} (b) \text{ Let } y &= 4 \cos \theta - 3 \sin \theta \\ &= 5 \cos(\theta + \alpha), \end{aligned}$$

Since  $-1 \leq \cos(\theta + \alpha) \leq 1$ ,

$$\therefore -5 \leq 5 \cos(\theta + \alpha) \leq 5$$

$\therefore$  The maximum value of  $y$  is 5.

The minimum value of  $y$  is -5.



### Classwork 2

- (a) Express  $3 \sin \theta - 2 \cos \theta$  in the form  $R \sin(\theta - \alpha)$  where  $R > 0$  and  $0^\circ \leq \alpha \leq 90^\circ$ .

- (b) Hence, or otherwise, find the range of possible values of  $3 \sin \theta - 2 \cos \theta$ .

$$[\sqrt{13} \sin(\theta - 33.69^\circ), -\sqrt{13} \leq 3 \sin \theta - 2 \cos \theta \leq \sqrt{13}]$$

**Example 12** Solve the equation  $\sqrt{3} \cos \theta + \sin \theta = \sqrt{2}$ , where  $0^\circ \leq \theta \leq 360^\circ$

Consider a right-angled triangle as shown in the figure.

$$r^2 = 3 + 1$$

$$r = 2$$

$$\text{and } \tan \alpha = \sqrt{3}$$

$$\alpha = 60^\circ$$

$$2 \sin 60^\circ = \sqrt{3}, 2 \cos 60^\circ = 1$$

The equation becomes


$$2 \sin 60^\circ \cos \theta + 2 \cos 60^\circ \sin \theta = \sqrt{2}$$

$$2 \sin(60^\circ + \theta) = \sqrt{2}$$

$$\sin(60^\circ + \theta) = \frac{\sqrt{2}}{2}$$

$$60^\circ + \theta = 135^\circ \text{ or } 405^\circ$$

$$\theta = 75^\circ \text{ or } 345^\circ$$



**Classwork 1** Solve the equation  $3 \sin \theta - 2 \cos \theta = 1$  for  $0^\circ \leq \theta \leq 360^\circ$ .  $[\theta = 49.79^\circ \text{ or } 197.59^\circ]$

**Example 13** Show that if the equation  $\sin \theta + k \cos \theta = \sqrt{2} + 1$  has a solution then  $k^2 \geq 2\sqrt{2} + 2$

$$\sqrt{k^2 + 1} \cdot \left( \sin \theta \cdot \frac{1}{\sqrt{k^2 + 1}} + \cos \theta \cdot \frac{k}{\sqrt{k^2 + 1}} \right) = \sqrt{2} + 1$$

$$\sin \theta \cdot \cos \alpha + \cos \theta \cdot \sin \alpha = \frac{\sqrt{2} + 1}{\sqrt{k^2 + 1}}$$

$$\sin(\theta + \alpha) = \frac{\sqrt{2} + 1}{\sqrt{k^2 + 1}}$$

The trigonometric equation has a solution, so  $-1 \leq \frac{\sqrt{2} + 1}{\sqrt{k^2 + 1}} \leq 1$

$$(\sqrt{2} + 1)^2 \leq (\sqrt{k^2 + 1})^2$$

$$2 + 2\sqrt{2} + 1 \leq k^2 + 1$$

$$k^2 \geq 2\sqrt{2} + 2$$

**Example 14** Prove that, for all real  $x$ , the maximum value of  $c \sin(x + A) + d \sin(x + B)$  is  $\sqrt{c^2 + d^2 + 2cd \cos(A - B)}$ .

$$c \sin(x + A) + d \sin(x + B)$$

$$= c \sin x \cos A + c \cos x \sin A + d \sin x \cos B + d \cos x \sin B$$

$$= \sin x(c \cos A + d \cos B) + \cos x(c \sin A + d \sin B)$$

$$= u \cdot \left[ \sin x \cdot \frac{(c \cos A + d \cos B)}{u} + \cos x \cdot \frac{(c \sin A + d \sin B)}{u} \right] \dots\dots (*)$$


$$\text{where } u = \sqrt{(c \cos A + d \cos B)^2 + (c \sin A + d \sin B)^2}$$

$$= \sqrt{(c^2 \cos^2 A + 2cd \cos A \cos B + d^2 \cos^2 B) + (c^2 \sin^2 A + 2cd \sin A \sin B + d^2 \sin^2 B)}$$


$$= \sqrt{(c^2 \cos^2 A + c^2 \sin^2 A) + (d^2 \cos^2 B + d^2 \sin^2 B) + (2cd \cos A \cos B + 2cd \sin A \sin B)}$$

$$= \sqrt{c^2 + d^2 + 2cd \cos(A - B)}$$

$$(*) c \sin(x + A) + d \sin(x + B) = u \cdot (\sin x \cos \alpha + \cos x \sin \alpha) \\ = u \sin(x + \alpha)$$



$$\therefore \text{The maximum value of } c \sin(x + A) + d \sin(x + B) \text{ is } u = \sqrt{c^2 + d^2 + 2cd \cos(A - B)}.$$



## Double angle formulae

$$\therefore \sin(A + B) = \sin A \cos B + \cos A \sin B$$

Let  $B = A$

$$\sin(A + A) = \sin A \cos A + \cos A \sin A$$

$$\sin 2A = 2 \sin A \cos A \dots (9)$$

Note: The above formula can be proved by the following method:

Given an isosceles triangle  $ABC$  with  $AB = AC = 1$ , let  $\angle BAC = 2\theta$ .

Draw  $AD \perp BC$ , where  $D$  is the foot of perpendicular from  $A$  to  $BC$ . Then  $\Delta ABD \cong \Delta ACD$  (R.H.S.)

$$\angle BAD = \angle CAD = \theta. \text{ (corr. } \angle \text{s } \cong \Delta \text{'s)}$$

$$AD = 1 \times \cos \theta = \cos \theta$$

$$BD = CD = 1 \times \sin \theta = \sin \theta$$

By finding the area of triangle in two different ways,

$$\frac{1}{2} AB \times AC \sin 2\theta = \frac{1}{2} BC \times AD$$

$$\therefore \sin 2\theta = 2 \sin \theta \cos \theta$$

$$\therefore \cos(A + B) = \cos A \cos B - \sin A \sin B$$

Let  $B = A$

$$\cos(A + A) = \cos A \cos A - \sin A \sin A$$

$$\therefore \cos 2A = \cos^2 A - \sin^2 A \dots (10)$$

By using the identity  $\sin^2 A + \cos^2 A = 1$

$$\text{So } \sin^2 A = 1 - \cos^2 A$$

$$\begin{aligned} \text{Sub. into (10): } \cos 2A &= \cos^2 A - (1 - \cos^2 A) \\ &= \cos^2 A - 1 + \cos^2 A \\ &= 2 \cos^2 A - 1 \end{aligned}$$

$$\therefore \cos 2A = 2 \cos^2 A - 1 \dots (11)$$

$$\text{Also, } \cos^2 A = 1 - \sin^2 A$$

$$\begin{aligned} \text{Sub. into (10): } \cos 2A &= (1 - \sin^2 A) - \sin^2 A \\ &= 1 - 2 \sin^2 A \end{aligned}$$

$$\therefore \cos 2A = 1 - 2 \sin^2 A \dots (12)$$


$$\therefore \tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

Let  $B = A$

$$\tan(A + A) = \frac{\tan A + \tan A}{1 - \tan A \tan A}$$

$$\therefore \tan 2A = \frac{2 \tan A}{1 - \tan^2 A} \dots (13)$$

$$\cot 2A = \frac{1}{\tan 2A} = \frac{1 - \tan^2 A}{2 \tan A} = \frac{1}{2} \left( \cot A - \frac{1}{\cot A} \right) \dots (14)$$



**Example 15** Without using calculators, evaluate  $\cos 36^\circ \cos 72^\circ$ .

$$\begin{aligned}\cos 36^\circ \cos 72^\circ &= \frac{2 \sin 36^\circ \cos 36^\circ \cos 72^\circ}{2 \sin 36^\circ} \\&= \frac{2 \sin 72^\circ \cos 72^\circ}{4 \sin 36^\circ} \\&= \frac{\sin 144^\circ}{4 \sin 36^\circ} = \frac{1}{4}\end{aligned}$$

**Example 16** Without using calculators, evaluate  $\cos 12^\circ \cos 24^\circ \cos 48^\circ \cos 96^\circ$ .

$$\begin{aligned}\cos 12^\circ \cos 24^\circ \cos 48^\circ \cos 96^\circ &= \frac{2 \sin 12^\circ \cos 12^\circ \cos 24^\circ \cos 48^\circ \cos 96^\circ}{2 \sin 12^\circ} \\&= \frac{2 \sin 24^\circ \cos 24^\circ \cos 48^\circ \cos 96^\circ}{4 \sin 12^\circ} \\&= \frac{2 \sin 48^\circ \cos 48^\circ \cos 96^\circ}{8 \sin 12^\circ} \\&= \frac{2 \sin 96^\circ \cos 96^\circ}{16 \sin 12^\circ} \\&= \frac{\sin 192^\circ}{16 \sin 12^\circ} = -\frac{1}{16}\end{aligned}$$

### Classwork 3

(a) By mathematical induction, prove that for all positive integer  $n$ ,

$$\cos \theta \cos 2\theta \cos 4\theta \cdots \cos 2^{n-1}\theta = \frac{\sin 2^n \theta}{2^n \sin \theta}.$$

(b) Hence evaluate  $\cos 20^\circ \cos 40^\circ \cos 80^\circ$  without using calculators.

**Example 17** Prove that  $\frac{\tan 2\theta + \sec 2\theta - 1}{\tan 2\theta - \sec 2\theta + 1} = \tan\left(\theta + \frac{\pi}{4}\right)$ .

$$\begin{aligned} \text{L.H.S.} &= \frac{\tan 2\theta + \sec 2\theta - 1}{\tan 2\theta - \sec 2\theta + 1} \\ &= \frac{\left(\frac{\sin 2\theta}{\cos 2\theta} + \frac{1}{\cos 2\theta} - 1\right)}{\left(\frac{\sin 2\theta}{\cos 2\theta} - \frac{1}{\cos 2\theta} + 1\right)} \cdot \frac{\cos 2\theta}{\cos 2\theta} \\ &= \frac{\sin 2\theta + 1 - \cos 2\theta}{\sin 2\theta - 1 + \cos 2\theta} \\ &= \frac{2\sin \theta \cos \theta + 1 - (1 - 2\sin^2 \theta)}{2\sin \theta \cos \theta - 1 + (1 - 2\sin^2 \theta)} \\ &= \frac{2\sin \theta \cos \theta + 2\sin^2 \theta}{2\sin \theta \cos \theta - 2\sin^2 \theta} \\ &= \frac{2\sin \theta (\cos \theta + \sin \theta)}{2\sin \theta (\cos \theta - \sin \theta)} \\ &= \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} \end{aligned}$$

$$\begin{aligned} \text{R.H.S.} &= \tan\left(\theta + \frac{\pi}{4}\right) \\ &= \frac{\tan \theta + \tan \frac{\pi}{4}}{1 - \tan \theta \cdot \tan \frac{\pi}{4}} \\ &= \frac{\tan \theta + 1}{1 - \tan \theta \cdot 1} \\ &= \frac{\left(\frac{\sin \theta}{\cos \theta} + 1\right)}{\left(1 - \frac{\sin \theta}{\cos \theta}\right)} \cdot \frac{\cos \theta}{\cos \theta} \\ &= \frac{\cos \theta + \sin \theta}{\cos \theta - \sin \theta} \end{aligned}$$

$\therefore \text{L.H.S.} = \text{R. H.S.}$

**Example 18** In the given figure,  $\Delta ABC$  is isosceles with  $AC = BA = a$ ,  $AB = 1$ .  $P$  is a point on  $AB$  such that  $\angle ACP = \alpha$ ,  $\angle BCP = 2\alpha$ , show that  $AP = \frac{1}{1+2\cos\alpha}$ . Hence deduce that  $\frac{1}{3} < AP < \frac{1}{2}$ .

Let  $AP = x$ , then  $BP = 1 - x$ .

$$\text{In } \Delta APC, \frac{x}{\sin \alpha} = \frac{CP}{\sin A} \dots (1)$$

$$\text{In } \Delta BPC, \frac{1-x}{\sin 2\alpha} = \frac{CP}{\sin B} \dots (2)$$

$\angle A = \angle B$  (base  $\angle$ s isos.  $\Delta$ )

$$\therefore (1) = (2): \frac{x}{\sin \alpha} = \frac{1-x}{\sin 2\alpha}$$

$$\frac{x}{\sin \alpha} = \frac{1-x}{2 \sin \alpha \cos \alpha}$$

$$2x \cos \alpha = 1 - x$$

$$x = AP = \frac{1}{1+2\cos\alpha}$$

$$0^\circ < \angle C = 3\alpha < 180^\circ$$

$$\therefore 0^\circ < \alpha < 60^\circ$$


$$\frac{1}{2} < \cos \alpha < 1$$

$$1 < 2 \cos \alpha < 2$$

$$2 < 1 + 2 \cos \alpha < 3$$

$$\frac{1}{2} > \frac{1}{1+2\cos\alpha} > \frac{1}{3}$$

$$\frac{1}{3} < AP < \frac{1}{2}$$



**Example 19** Prove that for all real values of  $x$ ,

$$-7 \leq \sin^2 x - 24 \sin x \cos x + 11 \cos^2 x \leq 19.$$

$$\sin^2 x - 24 \sin x \cos x + 11 \cos^2 x$$

$$= \frac{1-\cos 2x}{2} - 12 \sin 2x + 11 \left( \frac{1+\cos 2x}{2} \right)$$

$$= 6 + 5 \cos 2x - 12 \sin 2x$$

$$= 6 + 13 \left( \cos 2x \cdot \frac{5}{13} - \sin 2x \cdot \frac{12}{13} \right)$$

$$= 6 + 13(\cos 2x \cos \alpha - \sin 2x \sin \alpha)$$


$$= 6 + 13 \cos(2x + \alpha)$$

$$-1 \leq \cos(2x + \alpha) \leq 1$$

$$-13 \leq 13 \cos(2x + \alpha) \leq 13$$

$$-7 \leq 6 + 13 \cos(2x + \alpha) \leq 19$$

$$-7 \leq \sin^2 x - 24 \sin x \cos x + 11 \cos^2 x \leq 19.$$



**Half angle formulae**

$$\cos 2A = 2 \cos^2 A - 1 \dots (11)$$

$$\cos 2A = 1 - 2 \sin^2 A \dots (12)$$

$$\tan 2A = \frac{2 \tan A}{1 - \tan^2 A} \dots (13)$$

Formulae (11) and (12) can be changed into:

$$\cos^2 A = \frac{1 + \cos 2A}{2} \dots (15) \text{ or } \cos A = \pm \sqrt{\frac{1 + \cos 2A}{2}} \text{ or } \cos \frac{A}{2} = \pm \sqrt{\frac{1 + \cos A}{2}}$$

$$\sin^2 A = \frac{1 - \cos 2A}{2} \dots (16) \text{ or } \sin A = \pm \sqrt{\frac{1 - \cos 2A}{2}} \text{ or } \sin \frac{A}{2} = \pm \sqrt{\frac{1 - \cos A}{2}}$$

$$\tan \frac{A}{2} = \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} = \pm \sqrt{\frac{1 - \cos A}{1 + \cos A}} \dots (17)$$

Using formulae (15), (16) and (17), we can find the trigonometric ratios of  $22.5^\circ, 67.5^\circ, 112.5^\circ, 157.5^\circ, 202.5^\circ, 247.5^\circ, 292.5^\circ, 337.5^\circ$ . Equivalently, in radians:  $\frac{\pi}{8}, \frac{3\pi}{8}, \frac{5\pi}{8}, \frac{7\pi}{8}, \frac{9\pi}{8}, \frac{11\pi}{8}, \frac{13\pi}{8}, \frac{15\pi}{8}$ .

$$\sin 22.5^\circ = \sqrt{\frac{1 - \cos 2 \times 22.5^\circ}{2}} = \sqrt{\frac{1 - \cos 45^\circ}{2}} = \sqrt{\frac{1 - \frac{\sqrt{2}}{2}}{2}} = \frac{\sqrt{2 - \sqrt{2}}}{2}$$

$$\cos 22.5^\circ = \sqrt{\frac{1 + \cos 2 \times 22.5^\circ}{2}} = \sqrt{\frac{1 + \cos 45^\circ}{2}} = \sqrt{\frac{1 + \frac{\sqrt{2}}{2}}{2}} = \frac{\sqrt{2 + \sqrt{2}}}{2}$$

$$1 = \tan(2 \times 22.5^\circ) = \frac{2 \tan 22.5^\circ}{1 - \tan^2 22.5^\circ}$$

Cross multiplying:  $1 - \tan^2 22.5^\circ = 2 \tan 22.5^\circ$

$$\tan^2 22.5^\circ + 2 \tan 22.5^\circ - 1 = 0$$

This is a quadratic equation in  $\tan 22.5^\circ$ .

$$\tan 22.5^\circ = -1 \pm \sqrt{2}$$

$$\therefore 0 < \tan 22.5^\circ < \tan 45^\circ = 1$$

$$\therefore \tan 22.5^\circ = -1 + \sqrt{2}$$

$$\sin 292.5^\circ = \sin(270^\circ + 22.5^\circ) = -\cos 22.5^\circ = -\frac{\sqrt{2 + \sqrt{2}}}{2}$$

$$\cos 292.5^\circ = \cos(270^\circ + 22.5^\circ) = \sin 22.5^\circ = \frac{\sqrt{2 - \sqrt{2}}}{2}$$

$$\tan 292.5^\circ = \tan(270^\circ + 22.5^\circ) = -\cot 22.5^\circ = -\frac{1}{-1 + \sqrt{2}} = -(1 + \sqrt{2})$$

The other trigonometric ratios are similarly found.

**Triple angle formulae (三倍角公式)**

$$\sin 3A = \sin(2A + A)$$

$$\begin{aligned} &= \sin 2A \cos A + \cos 2A \sin A \\ &= 2 \sin A \cos A \cos A + (1 - 2 \sin^2 A) \sin A \\ &= 2 \sin A (1 - \sin^2 A) + \sin A - 2 \sin^3 A \\ &= 2 \sin A - 2 \sin^3 A + \sin A - 2 \sin^3 A \end{aligned}$$

$\sin 3A = 3 \sin A - 4 \sin^3 A \cdots (18)$        $\sin 3A = 3 \times ? - 4 \times ??$  Express  $\sin 3A$  in terms of  $\sin A$  only

$$\cos 3A = \cos(2A + A)$$

$$\begin{aligned} &= \cos 2A \cos A - \sin 2A \sin A \\ &= (2 \cos^2 A - 1) \cos A - 2 \sin A \cos A \sin A \\ &= 2 \cos^3 A - \cos A - 2 \cos A (1 - \cos^2 A) \\ &= 2 \cos^3 A - \cos A - 2 \cos A + 2 \cos^3 A \end{aligned}$$

$$\cos 3A = 4 \cos^3 A - 3 \cos A \cdots (19)$$

Express  $\cos 3A$  in terms of  $\cos A$  only

$$\tan 3A = \tan(2A + A) = \frac{\tan 2A + \tan A}{1 - \tan 2A \tan A}$$

$$\begin{aligned} &= \frac{\left(\frac{2 \tan A}{1 - \tan^2 A} + \tan A\right)}{\left(1 - \frac{2 \tan A}{1 - \tan^2 A} \cdot \tan A\right)} \cdot \frac{(1 - \tan^2 A)}{(1 - \tan^2 A)} \\ &= \frac{2 \tan A + \tan A (1 - \tan^2 A)}{(1 - \tan^2 A) - 2 \tan^2 A} \end{aligned}$$

$$\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A} \cdots (20) \quad \text{Express } \tan 3A \text{ in terms of } \tan A \text{ only.}$$

**Classwork 4**

$$\text{Prove that } \tan 4A = \frac{4 \tan A - 4 \tan^3 A}{1 - 6 \tan^2 A + \tan^4 A} \cdots (21)$$

$$\text{and } \tan 5A = \frac{5 \tan A - 10 \tan^3 A + \tan^5 A}{1 - 10 \tan^2 A + 5 \tan^4 A} \cdots (22)$$

**Example 20** Without using calculators, express  $\sin 18^\circ$ ,  $\cos 36^\circ$  and  $\tan 18^\circ$  in surd form.

Let  $\theta = 18^\circ$ , then  $5\theta = 90^\circ$ ,  $3\theta = 90^\circ - 2\theta$

$$\cos 3\theta = \cos(90^\circ - 2\theta) = \sin 2\theta$$

$$4\cos^3 \theta - 3\cos \theta = 2\sin \theta \cos \theta$$

$\therefore \cos \theta = \cos 18^\circ \neq 0$ , divide both sides by  $\cos \theta$

$$4\cos^2 \theta - 3 = 2\sin \theta$$

$$4(1 - \sin^2 \theta) - 3 = 2\sin \theta$$

$$4 - 4\sin^2 \theta - 3 = 2\sin \theta$$

$$4\sin^2 \theta + 2\sin \theta - 1 = 0$$

This is a quadratic equation in  $\sin \theta$ .

$$\sin 18^\circ = \frac{-1 \pm \sqrt{5}}{4}$$

$\therefore 0^\circ < \theta < 90^\circ \therefore \sin 18^\circ > 0$

$$\sin 18^\circ = \frac{-1 + \sqrt{5}}{4}$$

Let  $\theta = 36^\circ$ , then  $5\theta = 180^\circ$ ,  $3\theta = 180^\circ - 2\theta$

$$\sin 3\theta = \sin(180^\circ - 2\theta) = \sin 2\theta$$

$$3\sin \theta - 4\sin^3 \theta = 2\sin \theta \cos \theta$$

$\therefore \sin \theta = \sin 36^\circ \neq 0$ , divide both sides by  $\sin \theta$

$$3 - 4\sin^2 \theta = 2\cos \theta$$

$$3 - 4(1 - \cos^2 \theta) = 2\cos \theta$$

$$3 - 4 + 4\cos^2 \theta = 2\cos \theta$$

$$4\cos^2 \theta - 2\cos \theta - 1 = 0$$

This is a quadratic equation in  $\cos \theta$ .

$$\cos 36^\circ = \frac{1 \pm \sqrt{5}}{4}$$

$\therefore 0^\circ < \theta < 90^\circ \therefore \cos 36^\circ > 0$

$$\cos 36^\circ = \frac{1 + \sqrt{5}}{4}$$

Let  $\beta = 18^\circ$ , then  $5\beta = 90^\circ$ ,  $3\beta = 90^\circ - 2\beta$ , let  $\tan \beta = t$

$$\tan 3\beta = \tan(90^\circ - 2\beta) = \cot 2\beta$$

$$\frac{3t - t^3}{1 - 3t^2} = \frac{1 - t^2}{2t}$$

$$6t^2 - 2t^4 = 1 - t^2 - 3t^2 + 3t^4$$

$$5t^4 - 10t^2 + 1 = 0$$

This is a quadratic equation in  $t^2$ .

$$t^2 = \frac{5 \pm \sqrt{20}}{5} \Rightarrow t = \sqrt{\frac{5 \pm \sqrt{20}}{5}}$$

$$\therefore 0^\circ < 18^\circ < 45^\circ \therefore 0 < \tan 18^\circ < \tan 45^\circ = 1 \text{ and } \sqrt{\frac{5 + \sqrt{20}}{5}} > 1$$

$$t^2 = \frac{5 - \sqrt{20}}{5} = \frac{5 - 2\sqrt{5}}{5}$$

$$\tan 18^\circ = \sqrt{\frac{5 - 2\sqrt{5}}{5}}$$

**Example 21** Find  $\cos 36^\circ$  without using triple angle formula.

Consider the following triangle  $\Delta ABC$ .

Given  $AB = AC = 1$ .  $D$  is a point lying on  $AC$  such that  $AD = BD = BC = x$ .

Let  $\angle A = \theta$ ,  $CD = 1 - x$ .

Then  $\angle ABD = \theta$ , (base  $\angle$ s isos.  $\Delta$ )

$\angle BDC = 2\theta$  (ext.  $\angle$  of  $\Delta$ )

$\angle ACB = 2\theta$  (base  $\angle$ s isos.  $\Delta$ )

$\angle ABC = 2\theta$  (base  $\angle$ s isos.  $\Delta$ )

$\angle CBD = 2\theta - \theta = \theta$

$\Delta ABC \sim \Delta BCD$  (equiangular)

In  $\Delta ABC$ ,  $\theta + 2\theta + 2\theta = 180^\circ$  ( $\angle$  sum of  $\Delta$ )

$\theta = 36^\circ$

$$\frac{AB}{BC} = \frac{BC}{CD} \quad (\text{corr. sides, } \sim \Delta \text{s})$$

$$\frac{1}{x} = \frac{x}{1-x}$$

$$1 - x = x^2$$

$$x^2 + x - 1 = 0$$

$$x = \frac{-1 + \sqrt{5}}{2} \quad \text{or} \quad \frac{-1 - \sqrt{5}}{2} \quad (< 0, \text{ rejected})$$

Draw  $DE \perp AB$  as shown. Then  $\Delta ADE \cong \Delta BDE$  (R.H.S.)

$$AE = ED = \frac{1}{2} \quad (\text{corr. sides, } \cong \Delta \text{s})$$

$$\cos 36^\circ = \frac{AE}{AD} = \frac{\frac{1}{2}}{x} = \frac{\frac{1}{2}}{\frac{-1+\sqrt{5}}{2}} = \frac{1}{-1 + \sqrt{5}} = \frac{1}{-1 + \sqrt{5}} \cdot \frac{\sqrt{5} + 1}{\sqrt{5} + 1} = \frac{1 + \sqrt{5}}{4}$$

**Example 22** Solve the cubic equation  $x^3 - 3x - 1 = 0$  and hence prove that  $\cos 20^\circ$  is irrational.

Let  $x = 2 \cos \theta$ , then the equation becomes  $(2 \cos \theta)^3 - 3(2 \cos \theta) - 1 = 0$

$$8 \cos^3 \theta - 6 \cos \theta = 1$$

$$4 \cos^3 \theta - 3 \cos \theta = \frac{1}{2}$$

$$\cos 3\theta = \frac{1}{2}$$

$$3\theta = 60^\circ, 300^\circ, 420^\circ, 660^\circ, 780^\circ, 1020^\circ$$

$$\theta = 20^\circ, 100^\circ, 140^\circ, 220^\circ, 260^\circ, 340^\circ$$

$$x = 2 \cos \theta = 2 \cos 20^\circ, 2 \cos 100^\circ, 2 \cos 140^\circ, 2 \cos 220^\circ, 2 \cos 260^\circ, 2 \cos 340^\circ$$

$$\therefore \cos 340^\circ = \cos 20^\circ, \cos 260^\circ = \cos 100^\circ, \cos 220^\circ = \cos 140^\circ$$

$\therefore$  The 3 roots of  $x^3 - 3x - 1 = 0$  are  $2 \cos 20^\circ, 2 \cos 100^\circ, 2 \cos 140^\circ = 1.879, -1.532, -0.347$ .

To prove  $\cos 20^\circ$  is irrational, we use the method of contradiction.

If  $\cos 20^\circ$  is rational, then  $2 \cos 20^\circ$  is also rational  $\therefore$  One root of the cubic equation is rational

Suppose  $2 \cos 20^\circ = \frac{b}{a}$ , then by factor theorem  $ax - b$  is a factor of  $x^3 - 3x - 1$


$$a = 1, b = \pm 1$$

$$\text{Let } f(x) = x^3 - 3x - 1$$

$$f(1) = 1^3 - 3 - 1 \neq 0 \text{ and } f(-1) = -1 + 3 - 1 \neq 0$$

$$\therefore x^3 - 3x - 1 = 0 \text{ has no rational root}$$

Our supposition is false. i.e.  $\cos 20^\circ$  is irrational.



Compound angle formulae


Circular function of  $t = \tan \frac{\theta}{2}$

**Circular function of  $t = \tan \frac{\theta}{2}$ .**

$$\sin \theta = 2 \sin \frac{\theta}{2} \cos \frac{\theta}{2} = \frac{2 \sin \frac{\theta}{2}}{\cos \frac{\theta}{2}} \cdot \cos^2 \frac{\theta}{2} = \frac{2 \tan \frac{\theta}{2}}{\sec^2 \frac{\theta}{2}} = \frac{2 \tan \frac{\theta}{2}}{1 + \tan^2 \frac{\theta}{2}} = \frac{2t}{1+t^2} \quad \dots (23)$$

$$\tan \theta = \tan 2\left(\frac{\theta}{2}\right) = \frac{2 \tan \frac{\theta}{2}}{1 - \tan^2 \frac{\theta}{2}} = \frac{2t}{1-t^2} \quad \dots (24)$$

$$\cos \theta = \frac{\sin \theta}{\tan \theta} = \frac{2t}{1+t^2} \times \frac{1-t^2}{2t} = \frac{1-t^2}{1+t^2} \quad \dots (25)$$



**Example 23** Solve  $2 \sin \theta - 3 \cos \theta = 1$  by using circular function of  $t = \tan \frac{\theta}{2}$ .

$$\frac{2(2t)}{1+t^2} - \frac{3(1-t^2)}{1+t^2} = 1$$

$$4t - 3 + 3t^2 = 1 + t^2$$

$$2t^2 + 4t - 4 = 0$$

$$t^2 + 2t - 2 = 0$$

$$t = \tan \frac{\theta}{2} = -1 \pm \sqrt{3}$$

$$\frac{\theta}{2} = 36.206^\circ \text{ or } -69.896^\circ \text{ (rejected) or } 216.206^\circ \text{ (rejected) or } 110.104^\circ$$

$$\theta = 72.4^\circ \text{ or } 220.2^\circ \text{ (correct to 1 decimal place)}$$

$$\frac{\sin x}{1 + \cos x} = \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{1 + 2 \cos^2 \frac{x}{2} - 1} = \frac{2 \sin \frac{x}{2} \cos \frac{x}{2}}{2 \cos^2 \frac{x}{2}} = \frac{\sin \frac{x}{2}}{\cos \frac{x}{2}} = \tan \frac{x}{2} \quad \dots (26)$$

$$\frac{\sin x}{1 + \cos x} = \frac{\sin x}{1 + \cos x} \cdot \frac{1 - \cos x}{1 - \cos x} = \frac{\sin x(1 - \cos x)}{1 - \cos^2 x} = \frac{1 - \cos x}{\sin x} \quad \dots (27)$$

$$\therefore \tan \frac{x}{2} = \frac{\sin x}{1 + \cos x} = \frac{1 - \cos x}{\sin x}$$

**Example 24** If  $\sin x = -\frac{3}{5}$  and  $270^\circ < x < 360^\circ$ , find  $\tan \frac{x}{2}$ .

$$\cos x = \sqrt{1 - \sin^2 x} = \frac{4}{5}$$

$$\tan \frac{x}{2} = \frac{\sin x}{1 + \cos x} = \frac{-\frac{3}{5}}{1 + \frac{4}{5}} = -\frac{1}{3}$$

**Example 25** Given  $\theta$  is in third quadrant and  $\tan \theta = \frac{4}{3}$ , evaluate  $\tan \frac{\theta}{2}$  without using calculators.

$$\frac{4}{3} = \frac{2t}{1-t^2}, \text{ where } t = \tan \frac{\theta}{2}, 180^\circ < \theta < 270^\circ, 90^\circ < \frac{\theta}{2} < 135^\circ, \tan \frac{\theta}{2} < 0$$

$$2t^2 + 3t - 2 = 0$$

$$t = \frac{1}{2} \text{ (rejected, } \because 90^\circ < \frac{\theta}{2} < 135^\circ, \tan \frac{\theta}{2} < 0) \text{ or } -2 \Rightarrow \tan \frac{\theta}{2} = -2$$

**Example 26** Given  $8 \sin \theta + 15 \cos \theta = -15$ , where  $0^\circ \leq \theta \leq 360^\circ$ .

(a) Use subsidiary angle to solve the equation.

(b) Use circular function of  $t = \tan \frac{\theta}{2}$  to solve the equation.

(c) Which method is correct? Explain briefly.

(a)  $\sqrt{8^2 + 15^2} = 17$ ,  $\cos \alpha = \frac{8}{17}$ ,  $\sin \alpha = \frac{15}{17}$ ,  $\alpha = 61.9275^\circ$

$$17 \cdot \left( \sin \theta \cdot \frac{8}{17} + \cos \theta \cdot \frac{15}{17} \right) = -15$$

$$\sin \theta \cos \alpha + \cos \theta \sin \alpha = -\frac{15}{17}$$

$$\sin(\theta + \alpha) = -\frac{15}{17}$$

$$\theta + 61.9275^\circ = -61.9275^\circ, 241.9275^\circ \text{ or } 298.0725^\circ$$

$$\theta = -123.8^\circ \text{ (rejected), } 180^\circ \text{ or } 236.1450^\circ$$

(b) Let  $t = \tan \frac{\theta}{2}$ ,  $\sin \theta = \frac{2t}{1+t^2}$ ,  $\cos \theta = \frac{1-t^2}{1+t^2}$ .

Then the equation  $8 \sin \theta + 15 \cos \theta = -15$  becomes

$$\frac{8 \cdot 2t}{1+t^2} + \frac{15(1-t^2)}{1+t^2} = -15$$

$$16t + 15 - 15t^2 = -15 - 15t^2$$

$$t = \tan \frac{\theta}{2} = -\frac{15}{8}$$


$$\frac{\theta}{2} = 118.0725^\circ \text{ or } 298.0725^\circ$$

$$\theta = 236.1450^\circ \text{ only}$$

(c) Note that  $t = \tan \frac{\theta}{2}$  is undefined for  $\frac{\theta}{2} = 90^\circ$

i.e.  $\theta = 180^\circ$  cannot be found in method 2.

Hence method 1 is correct.



**Half angle formula in terms of the sides of a triangle (outside syllabus)**

Given a triangle  $ABC$ . Let  $BC = a$ ,  $CA = b$ ,  $AB = c$ .

$$\text{Let } s = \frac{1}{2}(a+b+c), 2s = a+b+c, 2s - 2a = b+c-a, 2s - 2b = c+a-b, 2s - 2c = a+b-c$$

By Heron's formula, area of  $\Delta ABC = \sqrt{s(s-a)(s-b)(s-c)}$ . ... (28)

$$\text{On the other hand, area of triangle} = \frac{1}{2}bc \sin A$$

$$\frac{1}{2}bc \sin A = \sqrt{s(s-a)(s-b)(s-c)}$$

$$\therefore \sin A = \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)} \quad \dots (29)$$

$$\text{Cosine formula gives } \cos A = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\therefore 2 \cos^2 \frac{A}{2} - 1 = \frac{b^2 + c^2 - a^2}{2bc}$$

$$\cos^2 \frac{A}{2} = \frac{b^2 + c^2 - a^2 + 2bc}{4bc}$$

$$\cos \frac{A}{2} = \pm \frac{1}{2} \sqrt{\frac{(b+c)^2 - a^2}{bc}} = \pm \frac{1}{2} \sqrt{\frac{(b+c+a)(b+c-a)}{bc}} = \pm \sqrt{\frac{s(s-a)}{bc}}$$

$$\text{For } 0^\circ < A < 180^\circ, 0^\circ < \frac{A}{2} < 90^\circ, \cos \frac{A}{2} > 0$$

$$\therefore \cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}} \quad \dots (30)$$

$$\therefore \sin A = 2 \sin \frac{A}{2} \cos \frac{A}{2}$$

$$\text{By formula (29) and (30), } \frac{2}{bc} \sqrt{s(s-a)(s-b)(s-c)} = 2 \sin \frac{A}{2} \sqrt{\frac{s(s-a)}{bc}}$$

$$\therefore \sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}} \quad \dots (31)$$

$$\tan \frac{A}{2} = \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} = \sqrt{\frac{(s-b)(s-c)}{bc}} \times \sqrt{\frac{bc}{s(s-a)}} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}} \quad \dots (32)$$

The formulae for  $\sin \frac{B}{2}$ ,  $\sin \frac{C}{2}$ ,  $\cos \frac{B}{2}$ ,  $\cos \frac{C}{2}$ ,  $\tan \frac{B}{2}$  and  $\tan \frac{C}{2}$  are symmetric.

**Example 27**

Given  $\theta$  is an acute angle. If, in  $\Delta ABC$ ,  $(b + c) \cos \theta = 2\sqrt{bc} \cos \frac{A}{2}$ , prove that  $a = (b + c) \sin \theta$ .

Hence, find  $a$ , given  $b = 18.7$ ,  $c = 16.4$  and  $A = 57^\circ$ .

$$\begin{aligned}(b + c) \cos \theta &= 2\sqrt{bc} \cos \frac{A}{2} \\ &= 2\sqrt{bc} \cdot \sqrt{\frac{s(s-a)}{bc}} \\ &= 2\sqrt{s(s-a)} \\ &= \sqrt{(a+b+c)(b+c-a)}\end{aligned}$$

$$\begin{aligned}\cos \theta &= \sqrt{\left(\frac{a+b+c}{b+c}\right)\left(\frac{b+c-a}{b+c}\right)} \\ &= \sqrt{\left(1+\frac{a}{b+c}\right)\left(1-\frac{a}{b+c}\right)} \\ &= \sqrt{1-\left(\frac{a}{b+c}\right)^2}\end{aligned}$$

$$\cos^2 \theta = 1 - \left(\frac{a}{b+c}\right)^2$$

$$\sin^2 \theta = 1 - \cos^2 \theta = \left(\frac{a}{b+c}\right)^2$$

$$\sin \theta = \frac{a}{b+c} \quad (\because \theta \text{ is acute, } \sin \theta > 0)$$

$$\therefore a = (b + c) \sin \theta$$

$$b = 18.7, c = 16.4, A = 57^\circ$$

$$\begin{aligned}\cos \theta &= \frac{2\sqrt{bc} \cos \frac{A}{2}}{b+c} \\ &= \frac{2\sqrt{18.7 \times 16.4} \cos \frac{57^\circ}{2}}{18.7 + 16.4} \\ &= 0.876928352\end{aligned}$$

$$\theta = 28.72597648^\circ$$

$$a = (b + c) \sin \theta = (18.7 + 16.4) \sin 28.72597648^\circ = 16.87 \text{ (correct to 4 sig. fig.)}$$

**Example 28** In  $\Delta ABC$ , prove that  $a^2 - (b-c)^2 \cos^2 \frac{A}{2} = (b+c)^2 \sin^2 \frac{A}{2}$ .

$$\begin{aligned}
 \text{L.H.S.} &= a^2 - (b-c)^2 \cos^2 \frac{A}{2} \\
 &= a^2 - (b-c)^2 \cdot \frac{s(s-a)}{bc} \\
 &= \frac{a^2 bc - (b-c)^2 \cdot \frac{1}{4}(a+b+c)(b+c-a)}{bc} \\
 &= \frac{a^2 bc - (b-c)^2 \cdot \frac{1}{4}[(b+c)^2 - a^2]}{bc} \\
 &= \frac{a^2 bc + \frac{1}{4}a^2(b^2 - 2bc + c^2) - \frac{1}{4}(b-c)^2(b+c)^2}{bc} \\
 &= \frac{\frac{1}{4}a^2(b^2 + 2bc + c^2) - \frac{1}{4}(b-c)^2(b+c)^2}{bc} \\
 &= \frac{\frac{1}{4}(b+c)^2[a^2 - (b-c)^2]}{bc} \\
 &= \frac{\frac{1}{4}(b+c)^2(a-b+c)(a+b-c)}{bc} \\
 &= \frac{(b+c)^2(s-b)(s-c)}{bc} \\
 &= (b+c)^2 \sin^2 \frac{A}{2} = \text{R.H.S.}
 \end{aligned}$$

**Sum and Product formulae**

$$\sin(A+B) = \sin A \cos B + \cos A \sin B$$

$$\sin(A-B) = \sin A \cos B - \cos A \sin B$$

$$\sin(A+B) + \sin(A-B) = 2 \sin A \cos B$$

$$\therefore \sin A \cos B = \frac{1}{2} [\sin(A+B) + \sin(A-B)] \quad \dots\dots (33) \text{ Product to sum formula}$$

$$\sin(A+B) - \sin(A-B) = 2 \cos A \sin B$$

$$\therefore \cos A \sin B = \frac{1}{2} [\sin(A+B) - \sin(A-B)] \quad \dots\dots (34)$$

$$\cos(A+B) = \cos A \cos B - \sin A \sin B$$

$$\cos(A-B) = \cos A \cos B + \sin A \sin B$$

$$\cos(A+B) + \cos(A-B) = 2 \cos A \cos B$$

$$\therefore \cos A \cos B = \frac{1}{2} [\cos(A+B) + \cos(A-B)] \quad \dots\dots (35)$$

$$\cos(A+B) - \cos(A-B) = -2 \sin A \sin B$$

$$\therefore \sin A \sin B = -\frac{1}{2} [\cos(A+B) - \cos(A-B)] \quad \dots\dots (36)$$

Formulae (33) to (36) are called product to sum formulae.

$$\text{Let } x = A+B, y = A-B, \text{ then } A = \frac{x+y}{2}, B = \frac{x-y}{2},$$

$$(33) \text{ becomes } \sin \frac{x+y}{2} \cos \frac{x-y}{2} = \frac{1}{2} (\sin x + \sin y)$$

$$\therefore \sin x + \sin y = 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2} \quad \dots\dots (37)$$

$$(34) \text{ becomes } \cos \frac{x+y}{2} \sin \frac{x-y}{2} = \frac{1}{2} (\sin x - \sin y)$$

$$\therefore \sin x - \sin y = 2 \cos \frac{x+y}{2} \sin \frac{x-y}{2} \quad \dots\dots (38)$$

$$(35) \text{ becomes } \cos \frac{x+y}{2} \cos \frac{x-y}{2} = \frac{1}{2} (\cos x + \cos y)$$

$$\therefore \cos x + \cos y = 2 \cos \frac{x+y}{2} \cos \frac{x-y}{2} \quad \dots\dots (39)$$

$$(36) \text{ becomes } \sin \frac{x+y}{2} \sin \frac{x-y}{2} = -\frac{1}{2} (\cos x - \cos y)$$

$$\therefore \cos x - \cos y = -2 \sin \frac{x+y}{2} \sin \frac{x-y}{2} \quad \dots\dots (40)$$

Formulae (37) to (40) are called sum to product formulae.

Memorize: S + S  $\rightarrow$  2SC

S - S  $\rightarrow$  2CS

C + C  $\rightarrow$  2CC

C - C  $\rightarrow$  -2SS

There are no formulae for S + C, S - C, T + T or T - T !

**Example 29** If  $\cos 16^\circ = \sin 14^\circ + \sin d^\circ$  and  $0 < d < 90$ , find  $d$  without using calculators.

$$\sin d^\circ = \cos 16^\circ - \sin 14^\circ = \sin 74^\circ - \sin 14^\circ$$

$$\sin d^\circ = 2 \cos \frac{74^\circ + 14^\circ}{2} \sin \frac{74^\circ - 14^\circ}{2}$$

$$\sin d^\circ = \cos 44^\circ = \sin 46^\circ$$

$$d = 46$$

**Example 30**

Express  $\frac{\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ}{\sin 1^\circ + \sin 2^\circ + \dots + \sin 44^\circ}$  in the form  $a + b\sqrt{c}$ , where  $a, b$  and  $c$  are integers.

$$\begin{aligned}\cos n^\circ + \sin n^\circ &= \sqrt{2}(\cos 45^\circ \cos n^\circ + \sin 45^\circ \sin n^\circ) \quad (\text{Subsidiary angles}) \\ &= \sqrt{2} \cos(45^\circ - n^\circ)\end{aligned}$$

$$(\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ) + (\sin 1^\circ + \sin 2^\circ + \dots + \sin 44^\circ)$$

$$= \sqrt{2}(\cos 44^\circ + \cos 43^\circ + \dots + \cos 1^\circ)$$

$$\sin 1^\circ + \sin 2^\circ + \dots + \sin 44^\circ = (\sqrt{2} - 1)(\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ)$$

$$\frac{\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ}{\sin 1^\circ + \sin 2^\circ + \dots + \sin 44^\circ} = \frac{1}{\sqrt{2} - 1} = 1 + \sqrt{2}$$

**Method 2** Recall  $\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$  and  $\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$ .

Take  $A = 45^\circ - n^\circ$  and  $B = n^\circ$  for  $n = 1, 2, \dots, 22$ , we have

$$\begin{aligned}\frac{\cos 1^\circ + \cos 2^\circ + \dots + \cos 44^\circ}{\sin 1^\circ + \sin 2^\circ + \dots + \sin 44^\circ} &= \frac{2 \cos \frac{45^\circ}{2} \left( \cos \frac{43^\circ}{2} + \cos \frac{41^\circ}{2} + \dots + \cos \frac{1^\circ}{2} \right)}{2 \sin \frac{45^\circ}{2} \left( \cos \frac{43^\circ}{2} + \cos \frac{41^\circ}{2} + \dots + \cos \frac{1^\circ}{2} \right)} \\ &= \cot 22.5^\circ = \frac{1}{\tan 22.5^\circ} = \frac{1}{\sqrt{2} - 1} \quad (\text{found on page 14}) \\ &= 1 + \sqrt{2}\end{aligned}$$

**Example 31** Find the value of  $\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ$  without using calculators.

$$\sin 20^\circ \sin 40^\circ \sin 60^\circ \sin 80^\circ$$

$$= (\sin 80^\circ \sin 20^\circ)(\sin 40^\circ)(\sin 60^\circ)$$

$$= -\frac{1}{2}(\cos 100^\circ - \cos 60^\circ) \cdot (\sin 40^\circ) \left( \frac{\sqrt{3}}{2} \right) \quad (\because \sin 60^\circ = \frac{\sqrt{3}}{2})$$

$$= -\frac{\sqrt{3}}{4} \left( \cos 100^\circ \sin 40^\circ - \frac{1}{2} \sin 40^\circ \right) \quad (\because \cos 60^\circ = \frac{1}{2})$$

$$= -\frac{\sqrt{3}}{4} \left[ \frac{1}{2} (\sin 140^\circ - \sin 60^\circ) - \frac{1}{2} \sin 40^\circ \right]$$

$$= -\frac{\sqrt{3}}{8} (\sin 140^\circ - \sin 60^\circ - \sin 40^\circ)$$

$$= -\frac{\sqrt{3}}{8} \left( \sin 40^\circ - \frac{\sqrt{3}}{2} - \sin 40^\circ \right)$$

$$= \frac{3}{16}$$

**Classwork 5** Without using calculators, prove that

$$(a) \quad \sin 23^\circ \sin 40^\circ - \cos 43^\circ \cos 74^\circ + \cos 24^\circ \sin 83^\circ = \cos 17^\circ;$$

$$(b) \quad \cos 17^\circ \cos 81^\circ + \sin 43^\circ \cos 69^\circ + \frac{1}{2} \cos 142^\circ = 0.$$

**Example 32** Without using calculators, find the value of  $\sin 12^\circ \sin 24^\circ \sin 48^\circ \sin 96^\circ$ .

$$\sin 12^\circ \sin 24^\circ \sin 48^\circ \sin 96^\circ$$

$$= (\sin 12^\circ \sin 48^\circ) (\sin 24^\circ \sin 96^\circ)$$

$$= \frac{1}{4} (\cos 36^\circ - \cos 60^\circ) (\cos 72^\circ - \cos 120^\circ)$$

$$= \frac{1}{4} \left( \cos 36^\circ - \frac{1}{2} \right) \left( \cos 72^\circ + \frac{1}{2} \right)$$

$$= \frac{1}{16} (4 \cos 36^\circ \cos 72^\circ + 2 \cos 36^\circ - 2 \cos 72^\circ - 1)$$

$$= \frac{1}{16} \left( 1 + \frac{2 \sin 36^\circ \cos 36^\circ - 2 \sin 36^\circ \cos 72^\circ}{\sin 36^\circ} - 1 \right)$$

$$= \frac{1}{16} \left( \frac{\sin 72^\circ - \sin 108^\circ + \sin 36^\circ}{\sin 36^\circ} \right) = \frac{1}{16}$$

**Example 33** In  $\Delta ABC$ , prove that  $\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \leq \frac{1}{8}$ .

$$\text{LHS} = \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$$

$$= -\frac{1}{2} \left( \cos \frac{A+B}{2} - \cos \frac{A-B}{2} \right) \sin \frac{C}{2}$$

$$= \frac{1}{2} \left( \cos \frac{A-B}{2} - \sin \frac{C}{2} \right) \sin \frac{C}{2}$$

$$\leq \frac{1}{2} \left( 1 - \sin \frac{C}{2} \right) \sin \frac{C}{2} \quad (\text{equality holds when } A = B)$$

$$\leq \frac{1}{2} \left( \frac{1 - \sin \frac{C}{2} + \sin \frac{C}{2}}{2} \right)^2 \quad (\text{G.M.} \leq \text{A.M.: } ab \leq \left( \frac{a+b}{2} \right)^2 \text{ for } a \geq 0, b \geq 0)$$

$$= \frac{1}{8} = \text{R.H.S.} \quad (\text{equality holds when } 1 - \sin \frac{C}{2} = \sin \frac{C}{2}, \text{ i.e. } C = \frac{\pi}{3})$$

**Example 34** If  $A + B + C = 180^\circ$ , prove that  $\cos^2 A + (\cos B + \cos C)^2 - 1 = 4 \sin^2 \frac{A}{2} \cos B \cos C$

Hence, if also  $\cos B + \cos C - \cos A = 1$ , prove that  $\sec A - \sec B - \sec C = 1$ .

$$\begin{aligned}
 \text{L.H.S.} &= \cos^2 A + (\cos B + \cos C)^2 - 1 = \cos^2 A + \left( 2 \cos \frac{B+C}{2} \cos \frac{B-C}{2} \right)^2 - 1 \\
 &= \cos^2 A + 4 \left( \cos \frac{180^\circ - A}{2} \cos \frac{B-C}{2} \right)^2 - 1 = 4 \sin^2 \frac{A}{2} \cos^2 \frac{B-C}{2} - \sin^2 A \\
 &= 4 \sin^2 \frac{A}{2} \cos^2 \frac{B-C}{2} - \left( 2 \sin \frac{A}{2} \cos \frac{A}{2} \right)^2 = 4 \sin^2 \frac{A}{2} \left( \cos^2 \frac{B-C}{2} - \cos^2 \frac{A}{2} \right) \\
 &= 4 \sin^2 \frac{A}{2} \left[ \cos^2 \frac{B-C}{2} - \cos \frac{180^\circ - (B+C)}{2} \right] = 4 \sin^2 \frac{A}{2} \left( \cos^2 \frac{B-C}{2} - \sin^2 \frac{B+C}{2} \right) \\
 &= 4 \sin^2 \frac{A}{2} \left[ \frac{1 + \cos(B+C)}{2} - \frac{1 - \cos(B+C)}{2} \right] = 2 \sin^2 \frac{A}{2} [\cos(B+C) + \cos(B-C)] \\
 &= 4 \sin^2 \frac{A}{2} \cos B \cos C = \text{R.H.S.}
 \end{aligned}$$

If also  $\cos B + \cos C - \cos A = 1$ , then  $\cos B + \cos C = 1 + \cos A \dots\dots (*)$

$$\cos^2 A + (\cos B + \cos C)^2 - 1 = 4 \sin^2 \frac{A}{2} \cos B \cos C \quad (\text{proved})$$

$$\cos^2 A + (1 + \cos A)^2 - 1 = 4 \sin^2 \frac{A}{2} \cos B \cos C \quad \text{by } (*)$$

$$2 \cos^2 A + 2 \cos A = 4 \left( \frac{1 - \cos A}{2} \right) \cos B \cos C \Rightarrow \cos A (\cos A + 1) = (1 - \cos A) \cos B \cos C$$

$$\cos A (\cos B + \cos C) = (1 - \cos A) \cos B \cos C \quad \text{by } (*)$$

$$\cos B \cos C - \cos C \cos A - \cos A \cos B = \cos A \cos B \cos C$$

$$\frac{\cos B \cos C}{\cos A \cos B \cos C} - \frac{\cos C \cos A}{\cos A \cos B \cos C} - \frac{\cos A \cos B}{\cos A \cos B \cos C} = \frac{\cos A + \cos B \cos C}{\cos A + \cos B \cos C}$$

$$\sec A - \sec B - \sec C = 1$$

**Example 35** Prove that  $\frac{1}{2\sin\theta}(\operatorname{cosec} 2\theta - \operatorname{cosec} 4\theta) = \frac{\cos 3\theta}{\sin 2\theta \sin 4\theta}$ .

Hence find the sum to  $n$  terms of the series  $\frac{\cos 3\theta}{\sin 2\theta \sin 4\theta} + \frac{\cos 5\theta}{\sin 4\theta \sin 6\theta} + \frac{\cos 7\theta}{\sin 6\theta \sin 8\theta} + \dots$ .

$$\begin{aligned}\text{L.H.S.} &= \frac{1}{2\sin\theta}(\operatorname{cosec} 2\theta - \operatorname{cosec} 4\theta) \\ &= \frac{1}{2\sin\theta} \left( \frac{1}{\sin 2\theta} - \frac{1}{\sin 4\theta} \right) \\ &= \frac{\sin 4\theta - \sin 2\theta}{2\sin\theta \sin 2\theta \sin 4\theta} \\ &= \frac{2\sin 2\theta \cos 2\theta - \sin 2\theta}{2\sin\theta \sin 2\theta \sin 4\theta} \\ &= \frac{\sin 2\theta(2\cos 2\theta - 1)}{2\sin\theta \sin 2\theta \sin 4\theta} \\ &= \frac{2\sin\theta \cos\theta [2(2\cos^2\theta - 1) - 1]}{2\sin\theta \sin 2\theta \sin 4\theta} \\ &= \frac{4\cos^3\theta - 3\cos\theta}{\sin 2\theta \sin 4\theta} \\ &= \frac{\cos 3\theta}{\sin 2\theta \sin 4\theta} = \text{R.H.S.}\end{aligned}$$

$$\begin{aligned}\frac{1}{2\sin\theta}(\operatorname{cosec} 2r\theta - \operatorname{cosec}(2r+2)\theta) &= \frac{1}{2\sin\theta} \left( \frac{1}{\sin 2r\theta} - \frac{1}{\sin 2(r+1)\theta} \right) \\ &= \frac{\sin(2r\theta + 2\theta) - \sin 2r\theta}{2\sin\theta \sin 2r\theta \sin 2(r+1)\theta} \\ &= \frac{2\cos(2r\theta + \theta)\sin\theta}{2\sin\theta \sin 2r\theta \sin 2(r+1)\theta} \\ &= \frac{\cos 2(r+1)\theta}{\sin 2r\theta \sin 2(r+1)\theta}\end{aligned}$$

$$\begin{aligned}\frac{\cos 3\theta}{\sin 2\theta \sin 4\theta} + \frac{\cos 5\theta}{\sin 4\theta \sin 6\theta} + \frac{\cos 7\theta}{\sin 6\theta \sin 8\theta} + \dots + \frac{\cos(2n+1)\theta}{\sin 2n\theta \sin(2n+2)\theta} \\ &= \sum_{r=1}^n \frac{\cos(2r+1)\theta}{\sin 2r\theta \sin(2r+2)\theta} \\ &= \sum_{r=1}^n \frac{1}{2\sin\theta}(\operatorname{cosec} 2r\theta - \operatorname{cosec}(2r+2)\theta) \\ &= \frac{1}{2\sin\theta} \left( \frac{1}{\sin 2\theta} - \frac{1}{\sin(2n+2)\theta} \right)\end{aligned}$$

**Example 36** In  $\Delta ABC$ , if  $2 \sin B = \sin A + \sin C$ , then  $2 \cot \frac{B}{2} = \cot \frac{A}{2} + \cot \frac{C}{2}$ .

$$2 \sin B = \sin A + \sin C \text{ (Given)}$$

$$2 \sin B = 2 \sin \frac{A+C}{2} \cos \frac{A-C}{2}$$

$$2 \sin \frac{B}{2} \cos \frac{B}{2} = \cos \frac{B}{2} \cos \frac{A-C}{2} \quad (\because \frac{A+C}{2} = 90^\circ - \frac{B}{2})$$

$$2 \sin \frac{B}{2} = \cos \frac{A-C}{2} \quad (\text{clearly } \cos \frac{B}{2} \neq 0)$$

$$2 \sin \frac{180^\circ - (A+C)}{2} = \cos \frac{A-C}{2}$$

$$2 \cos \frac{A+C}{2} = \cos \frac{A-C}{2}$$

$$2 \cos \frac{A}{2} \cos \frac{C}{2} - 2 \sin \frac{A}{2} \sin \frac{C}{2} = \cos \frac{A}{2} \cos \frac{C}{2} + \sin \frac{A}{2} \sin \frac{C}{2}$$

$$\cos \frac{A}{2} \cos \frac{C}{2} = 3 \sin \frac{A}{2} \sin \frac{C}{2}$$

$$\cot \frac{A}{2} \cot \frac{C}{2} = 3 \Rightarrow \cot \frac{A}{2} = \frac{3}{\cot \frac{C}{2}} \quad \dots (*) \text{ or } \tan \frac{A}{2} = \frac{1}{3 \tan \frac{C}{2}}$$

$$2 \cot \frac{B}{2} = 2 \cot \frac{180^\circ - (A+C)}{2}$$

$$= 2 \tan \frac{A+C}{2}$$

$$= 2 \cdot \frac{\tan \frac{A}{2} + \tan \frac{C}{2}}{1 - \tan \frac{A}{2} \cdot \tan \frac{C}{2}}$$

$$= 2 \cdot \frac{\frac{1}{3 \tan \frac{C}{2}} + \tan \frac{C}{2}}{1 - \frac{1}{3 \tan \frac{C}{2}} \cdot \tan \frac{C}{2}}$$

$$= 2 \cdot \frac{\frac{1}{3 \tan \frac{C}{2}} + \tan \frac{C}{2}}{1 - \frac{1}{3 \tan \frac{C}{2}} \cdot \tan \frac{C}{2}} \quad \text{by } (*)$$

$$= \frac{1}{\tan \frac{C}{2}} + 3 \tan \frac{C}{2}$$

$$= \cot \frac{C}{2} + \frac{3}{\cot \frac{C}{2}}$$

$$= \cot \frac{A}{2} + \cot \frac{C}{2} \quad \text{by } (*)$$

**Example 37** If  $A + B + C = 180^\circ$ , prove that

$$\sin^3 A \sin(B - C) + \sin^3 B \sin(C - A) + \sin^3 C \sin(A - B) = 0$$

But  $\sin^2 A \sin(B - C) + \sin^2 B \sin(C - A) + \sin^2 C \sin(A - B) = 0$  only if  $\Delta ABC$  is isosceles.

$$\begin{aligned}\sin^3 A \sin(B - C) &= \sin^2 A \cdot \frac{1}{2} [\cos(A + C - B) - \cos(A + B - C)]\end{aligned}$$

$$= \frac{1}{2} \sin^2 A [\cos(180^\circ - 2B) - \cos(180^\circ - 2C)]$$

$$= \frac{1}{2} \sin^2 A (\cos 2C - \cos 2B)$$

$$= \frac{1}{2} \sin^2 A (1 - 2 \sin^2 C - 1 + 2 \sin^2 B)$$

$$= \frac{1}{2} \sin^2 A (2 \sin^2 B - 2 \sin^2 C)$$

$$= \sin^2 A (\sin^2 B - \sin^2 C)$$

Similarly,  $\sin^3 B \sin(C - A) = \sin^2 B (\sin^2 C - \sin^2 A)$ ,  $\sin^3 C \sin(A - B) = \sin^2 C (\sin^2 A - \sin^2 B)$

$$\therefore \sin^3 A \sin(B - C) + \sin^3 B \sin(C - A) + \sin^3 C \sin(A - B)$$

$$= \sin^2 A (\sin^2 B - \sin^2 C) + \sin^2 B (\sin^2 C - \sin^2 A) + \sin^2 C (\sin^2 A - \sin^2 B) = 0$$

If  $\sin^2 A \sin(B - C) + \sin^2 B \sin(C - A) + \sin^2 C \sin(A - B) = 0$ , then

$$\sin A (\sin^2 B - \sin^2 C) + \sin B (\sin^2 C - \sin^2 A) + \sin C (\sin^2 A - \sin^2 B) = 0$$

$$\sin A (\sin B - \sin C) (\sin B + \sin C) + \sin B (\sin C - \sin B) (\sin C + \sin B) + \sin^2 A (\sin C - \sin B) = 0$$

$$(\sin B - \sin C)[\sin A (\sin B + \sin C) - \sin B \sin C - \sin^2 A] = 0$$

$$(\sin B - \sin C)(\sin A - \sin B)(\sin C - \sin A) = 0$$

$$A = B \text{ or } B = C \text{ or } C = A$$

$\therefore \Delta ABC$  is isosceles

**Example 38** If  $n$  is an odd integer, and  $A, B, C$  are the angles of a triangle, prove that

$$\sin nA + \sin nB + \sin nC = 4 \sin \frac{n\pi}{2} \cos \frac{nA}{2} \cos \frac{nB}{2} \cos \frac{nC}{2}.$$

$$\text{L.H.S.} = (\sin nA + \sin nB) + \sin nC$$

$$= 2 \sin \frac{n(A+B)}{2} \cos \frac{n(A-B)}{2} + \sin nC = 2 \sin \frac{n(\pi-C)}{2} \cos \frac{n(A-B)}{2} + 2 \sin \frac{nC}{2} \cos \frac{nC}{2}$$

$$= 2 \sin \left[ k\pi + \left( \frac{\pi}{2} - \frac{nC}{2} \right) \right] \cos \frac{n(A-B)}{2} + 2 \sin \frac{nC}{2} \cos \frac{nC}{2}, \text{ where } n = 2k + 1$$

$$= 2(-1)^k \cos \frac{nC}{2} \cos \frac{n(A-B)}{2} + 2 \sin \frac{nC}{2} \cos \frac{nC}{2}$$

$$= 2 \sin \left( k\pi + \frac{\pi}{2} \right) \cos \frac{nC}{2} \cos \frac{n(A-B)}{2} + 2 \sin \frac{n[\pi-(A+B)]}{2} \cos \frac{nC}{2}$$

$$= 2 \cos \frac{nC}{2} \left\{ \sin \frac{n\pi}{2} \cos \frac{n(A-B)}{2} + \sin \left[ k\pi + \frac{\pi}{2} - \frac{n(A+B)}{2} \right] \right\}$$

$$= 2 \cos \frac{nC}{2} \left[ \sin \frac{n\pi}{2} \cos \frac{n(A-B)}{2} + (-1)^k \cos \frac{n(A+B)}{2} \right]$$

$$= 2 \cos \frac{nC}{2} \left[ \sin \frac{n\pi}{2} \cos \frac{n(A-B)}{2} + \sin \frac{n\pi}{2} \cos \frac{n(A+B)}{2} \right]$$

$$= 2 \sin \frac{n\pi}{2} \cos \frac{nC}{2} \left[ \cos \frac{n(A+B)}{2} + \cos \frac{n(A-B)}{2} \right] = 4 \sin \frac{n\pi}{2} \cos \frac{nA}{2} \cos \frac{nB}{2} \cos \frac{nC}{2} = \text{R.H.S.}$$

**Remark** If the question is changed into: If  $n$  is an even integer and  $A, B, C$  are the angles of a triangle, find a necessary condition that  $\sin nA + \sin nB + \sin nC = 0$ , is this condition sufficient?

$$\text{"Necessity"} \sin nA + \sin nB + \sin nC = 0$$

$$\Rightarrow \sin 2kA + \sin 2kB + \sin 2kC = 0, \text{ where } n = 2k$$

$$\Rightarrow 2 \sin k(A+B) \cos k(A-B) + 2 \sin k[\pi - (A+B)] \cos k[\pi - (A+B)] = 0$$

$$\Rightarrow \sin k(A+B) \cos k(A-B) + (-1)^{k+1} \sin k(A+B) (-1)^k \cos k(A+B) = 0$$

$$\Rightarrow \sin k(A+B) [\cos k(A-B) - \cos k(A+B)] = 0$$

$$\Rightarrow 2 \sin k(A+B) \sin kA \sin kB = 0$$

$$\Rightarrow k(A+B) = m\pi \quad \text{or} \quad kA = m\pi \quad \text{or} \quad kB = m\pi$$

$\therefore$  Necessary condition is: any one angle is  $\frac{m\pi}{k}$  or the sum of any two angles is  $\frac{m\pi}{k}$ .

The condition is "sufficient": If  $A = \frac{m\pi}{k}$  or  $A+B = \frac{m\pi}{k}$ , trace back the " $\Rightarrow$ " sign.

**Example 39**

If  $\sin \theta + \sin \phi = a$ ,  $\tan \theta + \tan \phi = b$ ,  $\sec \theta + \sec \phi = c$ , prove that  $8bc = a[4b^2 + (b^2 - c^2)^2]$ .

$$b^2 - c^2 = (\tan \theta + \tan \phi)^2 - (\sec \theta + \sec \phi)^2$$

$$= -2 + 2 \tan \theta \tan \phi - 2 \sec \theta \sec \phi, \text{ using the identity } \sec^2 A - \tan^2 A = 1$$

$$= \frac{-2 \cos \theta \cos \phi}{\cos \theta \cos \phi} + \frac{2 \sin \theta \sin \phi}{\cos \theta \cos \phi} - \frac{2}{\cos \theta \cos \phi}$$

$$= \frac{-2 \cos(\theta + \phi)}{\cos \theta \cos \phi} - \frac{2}{\cos \theta \cos \phi} = -2 \cdot \frac{1 + \cos(\theta + \phi)}{\cos \theta \cos \phi}$$

$$4b^2 + (b^2 - c^2)^2 = 4(\tan \theta + \tan \phi)^2 + 4 \cdot \left[ \frac{1 + \cos(\theta + \phi)}{\cos \theta \cos \phi} \right]^2$$

$$= 4 \cdot \left[ \left( \frac{\sin \theta}{\cos \theta} + \frac{\sin \phi}{\cos \phi} \right)^2 + \frac{1 + 2 \cos(\theta + \phi) + \cos^2(\theta + \phi)}{\cos^2 \theta \cos^2 \phi} \right]$$

$$= 4 \cdot \left[ \frac{(\sin \theta \cos \phi + \cos \theta \sin \phi)^2}{\cos^2 \theta \cos^2 \phi} + \frac{1 + 2 \cos(\theta + \phi) + \cos^2(\theta + \phi)}{\cos^2 \theta \cos^2 \phi} \right]$$

$$= 4 \cdot \frac{\sin^2(\theta + \phi) + 1 + 2 \cos(\theta + \phi) + \cos^2(\theta + \phi)}{\cos^2 \theta \cos^2 \phi}$$

$$= 4 \cdot \frac{1 + 1 + 2 \cos(\theta + \phi)}{\cos^2 \theta \cos^2 \phi}$$

$$= 8 \cdot \frac{1 + \cos(\theta + \phi)}{\cos^2 \theta \cos^2 \phi}$$

$$a[4b^2 + (b^2 - c^2)^2] = (\sin \theta + \sin \phi) \cdot 8 \cdot \frac{1 + \cos(\theta + \phi)}{\cos^2 \theta \cos^2 \phi}$$

$$= \frac{8}{\cos^2 \theta \cos^2 \phi} \cdot \left( 2 \sin \frac{\theta + \phi}{2} \cos \frac{\theta - \phi}{2} \right) \left( 2 \cos^2 \frac{\theta + \phi}{2} \right)$$

$$= \frac{32}{\cos^2 \theta \cos^2 \phi} \cdot \left( \sin \frac{\theta + \phi}{2} \cos \frac{\theta - \phi}{2} \right) \left( \cos^2 \frac{\theta + \phi}{2} \right)$$

$$8bc = 8(\tan \theta + \tan \phi)(\sec \theta + \sec \phi)$$

$$= 8 \cdot \frac{\sin(\theta + \phi)}{\cos \theta \cos \phi} \cdot \frac{\cos \theta + \cos \phi}{\cos \theta \cos \phi}$$

$$= \frac{8}{\cos^2 \theta \cos^2 \phi} \cdot \left( 2 \sin \frac{\theta + \phi}{2} \cos \frac{\theta + \phi}{2} \right) \cdot \left( 2 \cos \frac{\theta + \phi}{2} \cos \frac{\theta - \phi}{2} \right)$$

$$\therefore 8bc = a[4b^2 + (b^2 - c^2)^2].$$

**Theorem**

Let  $A, B, C$  be three angles such that  $0 \leq A, B, C \leq \frac{\pi}{2}$  and  $A + B + C = \frac{\pi}{2}$

Then  $\sin A + \sin B + \sin C \leq \frac{3}{2}$ , equality holds when  $A = B = C = \frac{\pi}{6}$ .

**Proof:**  $C = \frac{\pi}{2} - (A + B)$ ,  $\sin C = \cos(A + B)$

$$\sin A + \sin B + \sin C = \sin A + \sin B + \cos(A + B)$$

$$\begin{aligned} &= 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2} + 1 - 2 \sin^2 \frac{A+B}{2} \\ &= 2 \sin \frac{A+B}{2} \left( \cos \frac{A-B}{2} - \sin \frac{A+B}{2} \right) + 1 \\ &\leq 2 \left[ \frac{\sin \frac{A+B}{2} + \cos \frac{A-B}{2} - \sin \frac{A+B}{2}}{2} \right]^2 + 1 \quad \because ab \leq \left( \frac{a+b}{2} \right)^2 \end{aligned}$$

$$\text{equality holds when } a = b; \text{ i.e. } \cos \frac{A-B}{2} = 2 \sin \frac{A+B}{2} \dots (1)$$

$$\begin{aligned} &= \frac{1}{2} \cos^2 \frac{A-B}{2} + 1 \\ &\leq \frac{1}{2} + 1 = \frac{3}{2}, \text{ equality holds when } \frac{A-B}{2} = 0, \text{ i.e. } A = B \dots \dots (2) \end{aligned}$$

Combine (1) and (2), equality holds when  $A = B = C = \frac{\pi}{6}$

**Example 40** If  $0 < x < y < \frac{\pi}{2}$ , prove that  $\sin x + \cos y - \sin(x-y) \leq \frac{\pi}{2}$

Let  $A = x, B = \frac{\pi}{2} - y, C = y - x$ , then  $0 \leq A, B, C \leq \frac{\pi}{2}$  and  $A + B + C = \frac{\pi}{2}$

$$\begin{aligned} \sin x + \cos y - \sin(x-y) &= \sin x + \sin\left(\frac{\pi}{2} - y\right) + \sin(y-x) \\ &= \sin A + \sin B + \sin C \\ &\leq \frac{3}{2} \quad \text{by the above theorem} \\ &\leq \frac{\pi}{2} \end{aligned}$$

The question is proved.

**Example 41** Given  $\sin^2 x + \sin^2 y = \sin(x + y)$ , where  $x$  and  $y$  are acute angles. Prove that  $x + y = 90^\circ$

Proof: If  $x = y$ , then the equation becomes  $2 \sin^2 x = \sin 2x$

$$2 \sin^2 x = 2 \sin x \cos x$$

$$\because 0 < x < 90^\circ \therefore \sin x \neq 0$$

$$\sin x = \cos x$$

$$\tan x = 1 \Rightarrow x = y = 45^\circ$$

If  $x \neq y$ , without loss of generality let  $x > y > 0^\circ$ .

Since  $\sin^2 x + \sin^2 y = (\sin x + \sin y)^2 - 2 \sin x \sin y$

$$\begin{aligned} &= \left( 2 \sin \frac{x+y}{2} \cos \frac{x-y}{2} \right)^2 - 2 \sin x \sin y \\ &= 4 \sin^2 \frac{x+y}{2} \cos^2 \frac{x-y}{2} + \cos(x+y) - \cos(x-y) \\ &= [1 - \cos(x+y)][1 + \cos(x-y)] + \cos(x+y) - \cos(x-y) \\ &= 1 - \cos(x+y)\cos(x-y) \end{aligned}$$

Hence  $\sin^2 x + \sin^2 y = \sin(x + y)$  becomes

$$1 - \cos(x+y)\cos(x-y) = \sin(x+y)$$

$$\cos(x+y)\cos(x-y) = 1 - \sin(x+y) \quad \dots\dots\dots (*)$$

Suppose  $x + y \neq 90^\circ$

If  $x + y > 90^\circ$ , by (\*): LHS < 0, RHS > 0 impossible.

If  $x + y < 90^\circ$ , since  $x > y > 0^\circ$ ,

then  $\cos(x-y) > \cos(x+y) > 0$

By (\*)  $1 - \sin(x+y) = \cos(x+y) \cos(x-y) > \cos^2(x+y)$

$$1 - \sin(x+y) > 1 - \sin^2(x+y)$$

$$\sin^2(x+y) - \sin(x+y) > 0$$

$$\sin(x+y)[\sin(x+y) - 1] > 0$$

$\sin(x+y) < 0$  or  $\sin(x+y) > 1$  which is impossible.

Therefore  $x + y = 90^\circ$

The question is proved.

**Example 42** If  $\sin 5^\circ + \sin 10^\circ + \sin 15^\circ + \dots + \sin 170^\circ + \sin 175^\circ = \tan x^\circ$ , find  $x$ .

$$2(\sin 5^\circ + \sin 10^\circ + \sin 15^\circ + \dots + \sin 170^\circ + \sin 175^\circ) \sin 2.5^\circ$$

$$= \cos 2.5^\circ - \cos 7.5^\circ + \cos 7.5^\circ - \cos 12.5^\circ + \cos 12.5^\circ - \cos 17.5^\circ + \dots + \cos 172.5^\circ - \cos 177.5^\circ$$

$$= \cos 2.5^\circ - \cos 177.5^\circ$$

$$= 2 \sin 90^\circ \sin 87.5^\circ$$

$$= 2 \sin 87.5^\circ$$

$$\sin 5^\circ + \sin 10^\circ + \sin 15^\circ + \dots + \sin 170^\circ + \sin 175^\circ = \tan 87.5^\circ$$

$$x = 87.5$$

**Example 43**

(a) Prove that  $\sin(\theta + \frac{k-1}{2}\alpha) \sin \frac{k\alpha}{2} + \sin \frac{\alpha}{2} \sin(\theta + k\alpha) = \frac{1}{2} [\cos(\theta - \frac{\alpha}{2}) - \cos(\theta + \frac{2k+1}{2}\alpha)]$

Hence prove that  $\sin(\theta + \frac{k-1}{2}\alpha) \sin \frac{k\alpha}{2} + \sin \frac{\alpha}{2} \sin(\theta + k\alpha) = \sin(\theta + \frac{k\alpha}{2}) \sin \frac{k+1}{2}\alpha$

(b) Prove by mathematical induction that

$$\sin \theta + \sin(\theta + \alpha) + \dots + \sin[\theta + (n-1)\alpha] = \frac{\sin[\theta + (\frac{n-1}{2})\alpha] \sin \frac{n\alpha}{2}}{\sin \frac{\alpha}{2}}$$

$$\begin{aligned} (a) \quad & \sin(\theta + \frac{k-1}{2}\alpha) \sin \frac{k\alpha}{2} + \sin \frac{\alpha}{2} \sin(\theta + k\alpha) \\ &= -\frac{1}{2} [\cos(\theta + \frac{2k-1}{2}\alpha) - \cos(\theta - \frac{\alpha}{2})] - \frac{1}{2} [\cos(\theta + \frac{2k+1}{2}\alpha) - \cos(\theta + \frac{2k-1}{2}\alpha)] \\ &= \frac{1}{2} [\cos(\theta - \frac{\alpha}{2}) - \cos(\theta + \frac{2k+1}{2}\alpha)] \\ &\quad \sin(\theta + \frac{k-1}{2}\alpha) \sin \frac{k\alpha}{2} + \sin \frac{\alpha}{2} \sin(\theta + k\alpha) \\ &= -\frac{1}{2} [\cos(\theta + \frac{2k+1}{2}\alpha) - \cos(\theta - \frac{\alpha}{2})] \\ &= -\frac{1}{2} (-2) [\sin(\theta + \frac{k\alpha}{2}) \sin(\theta + \frac{k+1}{2}\alpha)] \\ &= \sin(\theta + \frac{k\alpha}{2}) \sin \frac{k+1}{2}\alpha \end{aligned}$$

$$(b) \quad n = 1, \text{ LHS} = \sin \theta, \text{ RHS} = \frac{\sin \theta \sin \frac{\alpha}{2}}{\sin \frac{\alpha}{2}} = \sin \theta$$

It is true for  $n = 1$ .

Suppose it is true for  $n = k$ .

$$\text{i.e. } \sin \theta + \sin(\theta + \alpha) + \dots + \sin[\theta + (k-1)\alpha] = \frac{\sin[\theta + (\frac{k-1}{2})\alpha] \sin \frac{k\alpha}{2}}{\sin \frac{\alpha}{2}} \dots \dots \dots (*)$$

When  $n = k + 1$ ,

$$\begin{aligned} \text{L.H.S.} &= \sin \theta + \sin(\theta + \alpha) + \dots + \sin[\theta + (k-1)\alpha] + \sin(\theta + k\alpha) \\ &= \frac{\sin[\theta + (\frac{k-1}{2})\alpha] \sin \frac{k\alpha}{2}}{\sin \frac{\alpha}{2}} + \sin(\theta + k\alpha) \quad \text{by } (*) \\ &= \frac{\sin[\theta + (\frac{k-1}{2})\alpha] \sin \frac{k\alpha}{2} + \sin \frac{\alpha}{2} \sin(\theta + k\alpha)}{\sin \frac{\alpha}{2}} \\ &= \frac{\sin(\theta + \frac{k\alpha}{2}) \sin \frac{k+1}{2}\alpha}{\sin \frac{\alpha}{2}} \quad \text{by (a)} \\ &= \text{R.H.S.} \end{aligned}$$

Hence it is also true for  $n = k + 1$

By the principle of induction, the statement is true for all positive integer  $n$ .

**Example 44**

In the figure,  $ABCDE$  is a regular pentagon of side = 1. Prove that  $\cos \frac{\pi}{5} + \cos \frac{3\pi}{5} = \frac{1}{2}$ .

$$\angle A = \angle B = \angle C = \angle D = \angle E = \frac{3\pi}{5} \quad (\angle \text{ sum of polygon})$$

Join  $BE$ . Draw  $AG \perp BE$ ,  $CH \perp BE$ ,  $DF \perp BE$ .

Then  $\Delta ABG \cong \Delta AEG$  (R.H.S.)

$$\begin{aligned}\angle ABG &= \angle AEG \quad (\text{corr. } \angle \text{s } \cong \Delta \text{s}) \\ &= \left( \pi - \frac{3\pi}{5} \right) \div 2 = \frac{\pi}{5} \quad (\angle \text{ sum of isos. } \Delta)\end{aligned}$$

$$\angle CBH = \angle B - \angle ABG = \frac{3\pi}{5} - \frac{\pi}{5} = \frac{2\pi}{5}$$

$$\angle DEF = \angle E - \angle AEG = \frac{3\pi}{5} - \frac{\pi}{5} = \frac{2\pi}{5}$$

$$\angle BHC = \frac{\pi}{2} = \angle DFE \quad (\text{by construction})$$

$$BC = DE = 1 \quad (\text{given})$$

$$\Delta BCH \cong \Delta EDF \quad (\text{A.A.S.})$$

$$CH = DF \quad (\text{corr. sides } \cong \Delta \text{s})$$

$$CH \parallel DF \quad (\text{int. } \angle \text{s supp.})$$

$CDFH$  is a rectangle (opp. sides are equal and  $\parallel$ )


$$BE = BG + GE = 2 \cos \frac{\pi}{5} \quad \dots \dots (1)$$

$$BE = BH + HF + FE = 2 \cos \frac{2\pi}{5} + 1 \quad \dots \dots (2)$$

$$(1) = (2): 2 \cos \frac{\pi}{5} = 2 \cos \frac{2\pi}{5} + 1$$

$$\cos \frac{\pi}{5} - \cos \frac{2\pi}{5} = \frac{1}{2}$$

$$\cos \frac{\pi}{5} + \cos \frac{3\pi}{5} = \frac{1}{2}$$



**Example 45**

In the figure,  $ABCDEFG$  is a regular heptagon of side = 1. Prove that  $\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} = \frac{1}{2}$ .

$$\angle A = \angle B = \angle C = \angle D = \angle E = \angle F = \angle G = \frac{5\pi}{7} \quad (\angle \text{ sum of polygon})$$

Join  $BG, CF$ . Draw  $AH \perp BG, BK \perp CF$ ,

$GJ \perp CF, DN \perp CF, EM \perp CF$ .

Then  $\Delta ABH \cong \Delta AGH$  (R.H.S.)

$\angle ABH = \angle AGH$  (corr.  $\angle$ s  $\cong$   $\Delta$ s)

$$= \left( \pi - \frac{5\pi}{7} \right) \div 2 = \frac{\pi}{7} \quad (\angle \text{ sum of isos. } \Delta)$$

Join  $AC, AF$ . Then  $\Delta ABC \cong \Delta AGF$  (S.A.S.)

$$\angle ACB = \angle AFG = \frac{\pi}{7} \quad \dots \dots (1) \quad (\text{corr. } \angle \text{s} \cong \Delta \text{s})$$

$AC = AF$  (corr. sides  $\cong$   $\Delta$ s)

$$\angle ACF = \angle AFC \quad \dots \dots (2) \quad (\text{base } \angle \text{s isos. } \Delta \text{s})$$

$\therefore \angle BCK = \angle GFJ$  by (1) and (2)

$$\angle BKC = \frac{\pi}{2} = \angle GJF \quad (\text{by construction})$$

$BC = GF = 1$  (given)

$\therefore \Delta BCK \cong \Delta GFJ$  (A.A.S.)

$BK = GJ$  (corr. sides  $\cong$   $\Delta$ s)

$BK \parallel GJ$  (int.  $\angle$ s supp.)

$BGJK$  is a rectangle (opp. sides are equal and parallel)

$$BG = KJ = 2BH = 2 \cos \frac{\pi}{7} \quad (\text{opp. sides of rectangle})$$

$$\angle GBK = \frac{\pi}{2} = \angle BGJ \quad (\text{int. } \angle \text{s } BG \parallel KJ)$$

$$\angle CBK = \angle B - \angle GBK - \angle ABH = \frac{5\pi}{7} - \frac{\pi}{2} - \frac{\pi}{7} = \frac{\pi}{7}$$

$$\angle FGJ = \angle G - \angle BGJ - \angle AGH = \frac{5\pi}{7} - \frac{\pi}{2} - \frac{\pi}{7} = \frac{\pi}{7}$$

$$\angle BCK = \pi - \frac{\pi}{2} - \frac{\pi}{7} = \frac{3\pi}{7} \quad (\angle \text{ sum of } \Delta)$$

$$\angle GFJ = \pi - \frac{\pi}{2} - \frac{\pi}{7} = \frac{3\pi}{7} \quad (\angle \text{ sum of } \Delta)$$

$$\angle DCN = \angle C - \angle BCK = \frac{5\pi}{7} - \frac{3\pi}{7} = \frac{2\pi}{7}$$

$$\angle EFM = \angle F - \angle GFJ = \frac{5\pi}{7} - \frac{3\pi}{7} = \frac{2\pi}{7}$$

$CD = EF = 1$  (given)

$$\angle DNM = \frac{\pi}{2} = \angle EMN \quad (\text{by construction})$$


$\therefore \Delta CDN \cong \Delta FEM$  (A.A.S.)

$CN = FM, DN = EM$  (corr. sides  $\cong$   $\Delta$ s)

$DN \parallel EM$  (int.  $\angle$ s supp.)

$DEM$  is a rectangle (opp. sides are equal and parallel)

$MN = 1$  (opp. sides of rectangle)



$$CF = CN + NM + MF = 2 \cos \frac{2\pi}{7} + 1 \quad \dots\dots (1)$$

$$CF = CK + KJ + JF = 2 \cos \frac{3\pi}{7} + 2 \cos \frac{\pi}{7} \dots\dots (2)$$

$$(1) = (2): 2 \cos \frac{3\pi}{7} + 2 \cos \frac{\pi}{7} = 2 \cos \frac{2\pi}{7} + 1$$

$$\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} - \cos \frac{2\pi}{7} = \frac{1}{2}$$

$$\cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} = \frac{1}{2} \quad (\because \cos(\pi - \theta) = -\cos \theta)$$

### Example 46

In general  $\cos \frac{\pi}{2n-1} + \cos \frac{3\pi}{2n-1} + \dots + \cos \frac{(2n-3)\pi}{2n-1} = \frac{1}{2}$  for  $n \geq 2$ .

Let  $y = \cos \frac{\pi}{2n-1} + \cos \frac{3\pi}{2n-1} + \dots + \cos \frac{(2n-3)\pi}{2n-1}$

$$\begin{aligned} 2y \sin \frac{\pi}{2n-1} &= \sin \frac{2\pi}{2n-1} + \left( \sin \frac{4\pi}{2n-1} - \sin \frac{2\pi}{2n-1} \right) + \dots + \left[ \sin \frac{(2n-2)\pi}{2n-1} - \sin \frac{(2n-4)\pi}{2n-1} \right] \\ &= \sin \frac{(2n-2)\pi}{2n-1} \\ &= \sin \left( \pi - \frac{\pi}{2n-1} \right) \\ &= \sin \frac{\pi}{2n-1} \end{aligned}$$

$$\therefore 0 < \frac{\pi}{2n-1} < \frac{\pi}{2}$$

$$\therefore \sin \frac{\pi}{2n-1} \neq 0$$

$$y = \cos \frac{\pi}{2n-1} + \cos \frac{3\pi}{2n-1} + \dots + \cos \frac{(2n-3)\pi}{2n-1} = \frac{1}{2}$$

$$\text{In particular, } n = 3, \cos \frac{\pi}{5} + \cos \frac{3\pi}{5} = \frac{1}{2}$$

$$n = 4, \cos \frac{\pi}{7} + \cos \frac{3\pi}{7} + \cos \frac{5\pi}{7} = \frac{1}{2}$$

Find the value of  $\cos 1^\circ + \cos 2^\circ + \cos 3^\circ + \dots + \cos 90^\circ$ .

$$y = \cos 1^\circ + \cos 2^\circ + \cos 3^\circ + \dots + \cos 90^\circ$$

$$\begin{aligned} 2y \sin 0.5^\circ &= (\sin 1.5^\circ - \sin 0.5^\circ) + (\sin 2.5^\circ - \sin 1.5^\circ) + \dots + (\sin 90.5^\circ - \sin 89.5^\circ) \\ &= \sin 90.5^\circ - \sin 0.5^\circ \\ &= 2 \cos 45.5^\circ \sin 45^\circ \end{aligned}$$

$$\cos 1^\circ + \cos 2^\circ + \cos 3^\circ + \dots + \cos 90^\circ = \frac{\cos 45.5^\circ \sin 45^\circ}{\sin \frac{1^\circ}{2}} = 56.794$$

# Formulae for the Trigonometric functions

Created by Mr. Francis Hung on 23 June 2008

Last updated: March 19, 2023

## I The magic hexagon:

Along each diagonal,

$$\csc \theta = \frac{1}{\sin \theta}$$

$$\sin \theta = \frac{1}{\csc \theta}$$

$$\sec \theta = \frac{1}{\cos \theta}$$

$$\cos \theta = \frac{1}{\sec \theta}$$

$$\cot \theta = \frac{1}{\tan \theta}$$


$$\tan \theta = \frac{1}{\cot \theta}$$

In each shaded triangle,

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$



The S family

The C family

In any three adjacent vertices,

$$\sin \theta = \cos \theta \cdot \tan \theta$$

$$\cot \theta = \cos \theta \cdot \csc \theta$$

$$\sec \theta = \tan \theta \cdot \csc \theta$$

$$\cos \theta = \sin \theta \cdot \cot \theta$$

$$\csc \theta = \cot \theta \cdot \sec \theta$$

$$\tan \theta = \sin \theta \cdot \sec \theta$$

## II General Solutions

$$\sin \theta = \sin \alpha, \quad \theta = 180^\circ n + (-1)^n \alpha$$

$$\theta = n\pi + (-1)^n \alpha, \text{ where } n \text{ is an integer.}$$

$$\cos \theta = \cos \alpha, \quad \theta = 360^\circ n \pm \alpha$$

$$\theta = 2n\pi \pm \alpha, \text{ where } n \text{ is an integer.}$$

$$\tan \theta = \tan \alpha, \quad \theta = 180^\circ n + \alpha$$

$$\theta = n\pi + \alpha, \text{ where } n \text{ is an integer.}$$

## III Compound Angle Formulae

$$\sin(A + B) = \sin A \cos B + \cos A \sin B$$

$$\text{SC} + \text{CS}$$

$$\sin(A - B) = \sin A \cos B - \cos A \sin B$$

$$\text{SC} - \text{CS}$$

$$\cos(A + B) = \cos A \cos B - \sin A \sin B$$

$$\text{CC} - \text{SS}$$

$$\cos(A - B) = \cos A \cos B + \sin A \sin B$$

$$\text{CC} + \text{SS}$$

$$\tan(A + B) = \frac{\tan A + \tan B}{1 - \tan A \tan B}$$

$$\tan(A - B) = \frac{\tan A - \tan B}{1 + \tan A \tan B}$$

$$\tan(A + B + C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan A \tan C}$$

## IV Multiple angles

$$\sin 2\theta = 2 \sin \theta \cdot \cos \theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta = 2 \cos^2 \theta - 1 = 1 - 2 \sin^2 \theta$$

$$\tan 2\theta = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

$$\sin 3\theta = 3 \sin \theta - 4 \sin^3 \theta$$

$$\cos 3\theta = 4 \cos^3 \theta - 3 \cos \theta$$

$$\tan 3\theta = \frac{3 \tan \theta - \tan^3 \theta}{1 - 3 \tan^2 \theta}$$


$$\tan 4\theta = \frac{4 \tan \theta - 4 \tan^3 \theta}{1 - 6 \tan^2 \theta + \tan^4 \theta}$$

**V Half angles**

Let  $t = \tan \frac{\theta}{2}$ , then  $\sin \theta = \frac{2t}{1+t^2}$

$$\cos \theta = \frac{1-t^2}{1+t^2}$$

$$\tan \theta = \frac{2t}{1-t^2}$$



$$\sin \frac{x}{2} = \pm \sqrt{\frac{1-\cos x}{2}}$$

$$\cos \frac{x}{2} = \pm \sqrt{\frac{1+\cos x}{2}}$$

$$\tan \frac{x}{2} = \pm \sqrt{\frac{1-\cos x}{1+\cos x}} = \frac{\sin x}{1+\cos x} = \frac{1-\cos x}{\sin x}$$

**VI Sum and Product**

**Sum**  $\sin A + \sin B = 2 \sin \frac{A+B}{2} \cos \frac{A-B}{2}$

$$\sin A - \sin B = 2 \cos \frac{A+B}{2} \sin \frac{A-B}{2}$$

$$\cos A + \cos B = 2 \cos \frac{A+B}{2} \cos \frac{A-B}{2}$$

$$\cos A - \cos B = -2 \sin \frac{A+B}{2} \sin \frac{A-B}{2}$$

**Product**  $\sin X \cos Y = \frac{1}{2} [\sin(X+Y) + \sin(X-Y)]$

$$\cos X \sin Y = \frac{1}{2} [\sin(X+Y) - \sin(X-Y)]$$

$$\cos X \cos Y = \frac{1}{2} [\cos(X+Y) + \cos(X-Y)]$$

$$\sin X \sin Y = -\frac{1}{2} [\cos(X+Y) - \cos(X-Y)]$$

**VII Differentiation:** In each shaded triangle's edge,

$$DS = + \frac{d \sin x}{dx} = \cos x$$

$$DC = - \frac{d \cos x}{dx} = -\sin x$$

$$\frac{d \tan x}{dx} = \sec^2 x$$

$$\frac{d \cot x}{dx} = -\csc^2 x$$

$$\frac{d \sec x}{dx} = \sec x \tan x$$

$$\frac{d \csc x}{dx} = -\csc x \cot x$$

**Integration:** the inverse process of differentiation

$$\int \cos x dx = \sin x + C$$

$$\int \sin x dx = -\cos x + C$$

$$\int \sec^2 x dx = \tan x + C$$

$$\int \csc^2 x dx = -\cot x + C$$

$$\int \sec x \tan x dx = \sec x + C$$

$$\int \csc x \cot x dx = -\csc x + C$$

## Supplementary Exercise on Trigonometry

Created by Mr. Francis Hung

Last updated: March 19, 2023

1. Prove that  $\cos^2(A + \theta) + \cos^2(B + \theta) + 2 \cos(A - B) \sin(A + \theta)\sin(B + \theta)$  is independent of  $\theta$ .
2. Prove that, if  $\sin(\alpha + \beta) = k \sin(\alpha - \beta)$ , then  $(k + 1) \cot \alpha = (k - 1) \cot \beta$ .
3. If  $x \sin \theta + y \cos \theta = \sin \varphi$  and  $x \cos \theta - y \sin \theta = \cos \varphi$ , express  $x$  in terms of  $\theta$  and  $\varphi$  as simple as possible.
4. Without using tables or calculators, find the values of  $\tan 15^\circ$  and  $\tan 22\frac{1}{2}^\circ$ .
5. If  $\cos \theta + \cos \varphi = x$  and  $\sin \theta + \sin \varphi = y$ , prove that  $\cos \frac{1}{2}(\theta - \varphi) = \pm \frac{1}{2} \sqrt{x^2 + y^2}$ .
6. Prove the identities:  $\sin^2 A + \sin^2 B - \sin^2(A - B) = 2 \sin A \sin B \cos(A - B)$ , and  

$$\frac{\tan 3A - 2 \tan 2A + \tan A}{4(\tan 3A - \tan 2A)} = \sin^2 A.$$
7. In any triangle  $ABC$ , prove that
  - (a)  $b^2 \sin(C - A) = (c^2 - a^2) \sin B$ ,
  - (b)  $a^2 - (b - c)^2 \cos^2 \frac{A}{2} = (b + c)^2 \sin^2 \frac{A}{2}$ ,
  - (c)  $\tan(\frac{A}{2} + B) = \frac{c+b}{c-b} \tan \frac{A}{2}$ .
8. Solve completely the triangle  $ABC$  in which  $a = 2.718$ ,  $b = 3.142$ ,  $A = 54^\circ 18'$ . Show that there are two possible triangles and find their areas.
9. If  $A, B, C$  are the angles of a triangle, prove that  

$$(\sin B - \cos B)^2 + (\sin C - \cos C)^2 - (\sin A - \cos A)^2 = 1 - 4 \sin A \cos B \cos C$$
.
10. Given that  $(1 + \cos A)(1 + \cos B)(1 + \cos C)(1 + \cos D) = p \sin A \sin B \sin C \sin D$ ,  
 prove that  $(1 - \cos A)(1 - \cos B)(1 - \cos C)(1 - \cos D) = \frac{1}{p} \sin A \sin B \sin C \sin D$ .
11. If  $A, B, C$  are the angles of a triangle, using sine rule to prove
 
$$\begin{cases} a = b \cos C + c \cos B \\ b = c \cos A + a \cos C \\ c = a \cos B + b \cos A \end{cases} \cdots (*)$$

Hence, solve the system (\*) and express  $\cos A, \cos B, \cos C$  in terms of  $a, b$  and  $c$ .
12. If  $\theta + \varphi = \frac{1}{4}\pi$ , prove that  $(1 + \tan \theta)(1 + \tan \varphi) = 2$ . Deduce the value of  $\tan \frac{1}{8}\pi$ .
13. Establish the identity  $\sin \theta(\cos 2\theta + \cos 4\theta + \cos 6\theta) = \sin 3\theta \cos 4\theta$ .  
 Prove that, if  $x = \cos 3\theta + \sin 3\theta$  and  $y = \cos \theta - \sin \theta$ , then  $x - y = 2y \sin 2\theta$ .
14. By expressing  $(3 + \cos \theta) \operatorname{cosec} \theta$  in terms of  $\tan \frac{1}{2}\theta$  ( $= t$ ), show that this expression cannot have any value between  $-2\sqrt{2}$  and  $2\sqrt{2}$ .
15. By projection of the sides of an equilateral triangle onto a certain line, or otherwise, prove that  

$$\cos \theta + \cos(\theta + \frac{2}{3}\pi) + \cos(\theta + \frac{4}{3}\pi) = 0$$
,  
 and find the value of  $\sin \theta + \sin(\theta + \frac{2}{3}\pi) + \sin(\theta + \frac{4}{3}\pi)$ .
16. If  $\tan \alpha = k \tan \beta$ , show that  $(k - 1) \sin(\alpha + \beta) = (k + 1) \sin(\alpha - \beta)$ .  
 Show that, if the equation  $\tan x = k \tan(x - \alpha)$  has real solutions (in  $x$ ),  $(k - 1)^2$  is not less than  $(k + 1)^2 \sin^2 \alpha$ .  
 Solve the equation when  $k = -2$  and  $\alpha = 30^\circ$ , give your answer in general solution.
17. (a) If  $\sin \alpha + \cos \alpha = 2a$ , form the quadratic equation whose roots are  $\sin \alpha$  and  $\cos \alpha$ .  
 (b) Solve the equation and find the general solution of  $x$ :  

$$\cos^2 x + \cos x - \sin x - \sin^2 x = 0$$
.

18. Let  $y = \sin \theta (3 \sin \theta - \sin 2\alpha) + \cos \theta (3 \cos \theta - \cos 2\alpha)$  ( $0^\circ < \alpha < 90^\circ$ ).  
 (a) Find the general solution of  $\theta$  such that  $y$  has a minimum value and find this value.  
 (b) Find the general solution of  $\theta$  such that  $y$  has a maximum value and find this value.  
 (c) Find also the maximum and minimum value of the expression  

$$\sin \alpha (3 \sin \alpha - \sin 2\theta) + \cos \alpha (3 \cos \alpha - \cos 2\theta)$$
 ( $0^\circ < \alpha < 90^\circ$ ).
19. If  $\alpha, \beta$  are two distinct roots of the equation  $a \cos \theta + b \sin \theta = c$ , prove that  

$$\frac{a}{b} \sin(\alpha + \beta) - \cos(\alpha + \beta) = 1.$$
20. Prove the identity  $\frac{\cos(2\theta + \varphi) + \cos(2\varphi + \theta)}{2 \cos(\theta + \varphi) - 1} = \frac{\cos(2\theta - \varphi) + \cos(2\varphi - \theta)}{2 \cos(\theta - \varphi) - 1}.$
21. Prove the identities  
 (a)  $\sin^2(2\theta + \varphi) + \sin^2(2\varphi + \theta) - \sin^2(\theta - \varphi) = 2 \cos(\theta - \varphi) \sin(2\theta + \varphi) \sin(2\varphi + \theta),$   
 (b)  $\tan(3A - \frac{3}{4}\pi) \tan(A + \frac{1}{4}\pi) = \frac{1 + 2 \sin 2A}{1 - 2 \sin 2A},$   
 (c)  $(\cos A + \cos B + \cos C)^2 + (\sin A + \sin B + \sin C)^2 = 1 + 8 \cos \frac{1}{2}(B-C) \cos \frac{1}{2}(C-A) \cos \frac{1}{2}(A-B),$   
 (d)  $\frac{\tan 3A}{\tan A} = \frac{2 \cos 2A + 1}{2 \cos 2A - 1},$   
 (e)  $\sin^2(A + \theta) + \sin^2(B + \theta) = 1 - \cos(A - B) \cos(A + B + 2\theta).$

**End of Exercise**

$$\begin{aligned}
 1. \quad & \cos^2(A + \theta) + \cos^2(B + \theta) + 2 \cos(A - B) \sin(A + \theta) \sin(B + \theta) \\
 &= \frac{1}{2} [1 + \cos 2(A + \theta)] + \frac{1}{2} [1 + 2 \cos 2(B + \theta)] - \cos(A - B) [\cos(A + B + 2\theta) - \cos(A - B)] \\
 &= 1 + \frac{1}{2} [\cos 2(A + \theta) + \cos 2(B + \theta)] - \cos(A - B) \cos(A + B + 2\theta) + \cos^2(A - B) \\
 &= 1 + \cos(A + B + 2\theta) \cos(A - B) - \cos(A - B) \cos(A + B + 2\theta) + \cos^2(A - B) \\
 &= 1 + \cos^2(A - B)
 \end{aligned}$$

After simplification, the expression does not carry  $\theta$ , which is independent of  $\theta$ .

$$2. \quad \sin(\alpha + \beta) = k \sin(\alpha - \beta), \text{ then } (k+1) \cot \alpha = (k-1) \cot \beta.$$

$$\sin \alpha \cos \beta + \cos \alpha \sin \beta = k \sin \alpha \cos \beta - k \cos \alpha \sin \beta$$

$$k \cos \alpha \sin \beta + \cos \alpha \sin \beta = k \sin \alpha \cos \beta - \sin \alpha \cos \beta$$

$$(k+1) \cos \alpha \sin \beta = (k-1) \sin \alpha \cos \beta$$

$$(k+1) \cot \alpha = (k-1) \cot \beta$$

$$3. \quad x \sin \theta + y \cos \theta = \sin \varphi \dots\dots(1)$$

$$x \cos \theta - y \sin \theta = \cos \varphi \dots\dots(2)$$

$$(1) \times \sin \theta: x \sin^2 \theta + y \sin \theta \cos \theta = \sin \theta \sin \varphi$$

$$+ (2) \times \cos \theta: x \cos^2 \theta - y \sin \theta \cos \theta = \cos \theta \cos \varphi$$

$$\begin{aligned}
 x &= \sin \theta \sin \varphi + \cos \theta \cos \varphi \\
 &= \cos(\varphi - \theta)
 \end{aligned}$$

$$4. \quad \text{Using the formula } \tan x = \frac{\sin 2x}{1 + \cos 2x} \quad (\text{proof: } \frac{\sin 2x}{1 + \cos 2x} = \frac{2 \sin x \cos x}{1 + 2 \cos^2 x - 1} = \tan x)$$

$$\tan 15^\circ = \frac{\sin 30^\circ}{1 + \cos 30^\circ} = \frac{\frac{1}{2}}{1 + \frac{\sqrt{3}}{2}} = \frac{1}{2 + \sqrt{3}} = \frac{1}{2 + \sqrt{3}} \cdot \frac{2 - \sqrt{3}}{2 - \sqrt{3}} = 2 - \sqrt{3}$$

$$\tan 22\frac{1}{2}^\circ = \frac{\sin 45^\circ}{1 + \cos 45^\circ} = \frac{\frac{1}{\sqrt{2}}}{1 + \frac{1}{\sqrt{2}}} = \frac{1}{\sqrt{2} + 1} = \frac{1}{\sqrt{2} + 1} \cdot \frac{\sqrt{2} - 1}{\sqrt{2} - 1} = \sqrt{2} - 1$$

$$\text{Alternatively, using the formula } \tan 2x = \frac{2 \tan x}{1 - \tan^2 x}$$

$$\tan 30^\circ = \frac{2 \tan 15^\circ}{1 - \tan^2 15^\circ}$$

$$\frac{1}{\sqrt{3}} = \frac{2 \tan 15^\circ}{1 - \tan^2 15^\circ}$$

$$1 - \tan^2 15^\circ = 2\sqrt{3} \tan 15^\circ$$

$$\tan^2 15^\circ + 2\sqrt{3} \tan 15^\circ - 1 = 0$$

$$\begin{aligned}
 \tan 15^\circ &= -\sqrt{3} \pm \sqrt{(\sqrt{3})^2 + 1} \\
 &= -\sqrt{3} + 2 \text{ or } -\sqrt{3} - 2
 \end{aligned}$$

$$\therefore 0^\circ < 15^\circ < 90^\circ \therefore \tan 15^\circ > 0 \Rightarrow -\sqrt{3} - 2 \text{ is rejected}$$

$$\tan 15^\circ = 2 - \sqrt{3}$$

$$\tan 22\frac{1}{2}^\circ \text{ can be found in a similar way.}$$

5.  $\cos \theta + \cos \varphi = x \dots\dots\dots(1)$

$\sin \theta + \sin \varphi = y \dots\dots\dots(2)$

$$\begin{aligned} & (1)^2 \cos^2 \theta + \cos^2 \varphi + 2 \cos \theta \cos \varphi = x^2 \\ & +) \quad (2)^2 \sin^2 \theta + \sin^2 \varphi + 2 \sin \theta \sin \varphi = y^2 \end{aligned}$$

$$\begin{aligned} & 1 + 1 + 2 \cos(\theta - \varphi) = x^2 + y^2 \\ & 2[1 + \cos(\theta - \varphi)] = x^2 + y^2 \end{aligned}$$

$$2[1 + 2 \cos^2 \frac{1}{2}(\theta - \varphi) - 1] = x^2 + y^2$$

$$4 \cos^2 \frac{1}{2}(\theta - \varphi) = x^2 + y^2$$

$$\cos \frac{1}{2}(\theta - \varphi) = \pm \frac{1}{2} \sqrt{x^2 + y^2}$$

6. RHS =  $2 \sin A \sin B \cos(A - B)$

$$= -[\cos(A + B) - \cos(A - B)] \cos(A - B)$$

$$= -\cos(A + B) \cos(A - B) + \cos^2(A - B)$$

$$= -\frac{1}{2} (\cos 2A + \cos 2B) + 1 - \sin^2(A - B)$$

$$= -\frac{1}{2} (1 - 2 \sin^2 A + 1 - 2 \sin^2 B) + 1 - \sin^2(A - B)$$

$$= \sin^2 A + \sin^2 B - \sin^2(A - B) = \text{LHS}$$

You may try to prove the identity from the left side.

$$\begin{aligned} \text{LHS} &= \frac{\tan 3A - 2 \tan 2A + \tan A}{4(\tan 3A - \tan 2A)} \\ &= \frac{1}{4} - \frac{\tan 2A - \tan A}{4(\tan 3A - \tan 2A)} \\ &= \frac{1}{4} - \frac{\sin(2A - A)}{4 \sin(3A - 2A)} \quad (\text{note that } \tan \alpha - \tan \beta = \frac{\sin \alpha}{\cos \alpha} - \frac{\sin \beta}{\cos \beta} = \frac{\sin \alpha \cos \beta - \cos \alpha \sin \beta}{\cos \alpha \cos \beta}) \\ &= \frac{1}{4} - \frac{\cos 2A \cos A}{4 \sin 3A \cos 2A} \\ &= \frac{1}{4} - \frac{\sin A \cos 3A}{4 \sin A \cos A} \\ &= \frac{1}{4} - \frac{4 \cos^3 A - 3 \cos A}{4 \cos A} \\ &= \frac{1}{4}[1 - (4 \cos^2 A - 3)] \\ &= \frac{1}{4}(4 - 4 \cos^2 A) \\ &= \sin^2 A = \text{RHS} \end{aligned}$$

7. In  $\Delta ABC$ ,  $A + B + C = 180^\circ$ ,  $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C} = k \Rightarrow a = k \sin A$ ,  $b = k \sin B$ ,  $c = k \sin C$ .

$$\begin{aligned} (a) \quad \frac{b^2}{c^2 - a^2} &= \frac{(k \sin B)^2}{(k \sin C)^2 - (k \sin A)^2} \\ &= \frac{\sin^2 B}{\sin^2 C - \sin^2 A} \\ &= \frac{\sin^2 B}{(\sin C - \sin A)(\sin C + \sin A)} \end{aligned}$$

$$\begin{aligned}
&= \frac{\sin^2 B}{2 \cos \frac{C+A}{2} \sin \frac{C-A}{2} \cdot 2 \sin \frac{A+C}{2} \cos \frac{A-C}{2}} \\
&= \frac{\sin^2 B}{2 \sin \frac{A+C}{2} \cos \frac{A+C}{2} \cdot 2 \sin \frac{C-A}{2} \cos \frac{C-A}{2}} \\
&= \frac{\sin^2 B}{\sin(A+C) \sin(C-A)} \\
&= \frac{\sin^2 B}{\sin(180^\circ - B) \sin(C-A)} \\
&= \frac{\sin^2 B}{\sin B \sin(C-A)} \\
&= \frac{\sin B}{\sin(C-A)}. \text{ Hence result follows.}
\end{aligned}$$

$$\begin{aligned}
(b) \quad &a^2 - (b-c)^2 \cos^2 \frac{1}{2} A \\
&= a^2 - (b-c)^2 \cdot \frac{1+\cos A}{2} \\
&= \frac{1}{2} [2b^2 + 2c^2 - 4bc \cos A - (b^2 - 2bc + c^2)(1 + \cos A)], \text{ by cosine rule} \\
&= \frac{1}{2} [2b^2 + 2c^2 - 4bc \cos A - (b^2 - 2bc + c^2 + b^2 \cos A - 2bc \cos A + c^2 \cos A)] \\
&= \frac{1}{2} [b^2 + 2bc + c^2 - (b^2 \cos A + 2bc \cos A + c^2 \cos A)] \\
&= \frac{1}{2} [(b+c)^2 - (b+c)^2 \cos A] \\
&= (b+c)^2 \frac{1-\cos A}{2} \\
&= (b+c)^2 \sin^2 \frac{1}{2} A \\
(c) \quad &\frac{c+b}{c-b} \tan \frac{A}{2} \\
&= \frac{k \sin C + k \sin B}{k \sin C - k \sin B} \cdot \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} \\
&= \frac{2 \sin \frac{B+C}{2} \cos \frac{C-B}{2}}{2 \cos \frac{B+C}{2} \sin \frac{C-B}{2}} \cdot \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}} \\
&= \frac{2 \cos \frac{A}{2} \cos \frac{C-B}{2}}{2 \sin \frac{A}{2} \sin \frac{C-B}{2}} \cdot \frac{\sin \frac{A}{2}}{\cos \frac{A}{2}}, \text{ note that } \sin \frac{B+C}{2} = \sin \frac{180^\circ - A}{2} = \cos \frac{A}{2},
\end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{\tan \frac{C-B}{2}} \quad \cos \frac{B+C}{2} \text{ can be simplified similarly.} \\
 &= \frac{1}{\tan \frac{180^\circ - A - B - B}{2}} \\
 &= \frac{1}{\tan \left( 90^\circ - \frac{A}{2} - B \right)} \\
 &= \tan \left( \frac{1}{2} A + B \right), \text{ this is known as tangent rule.}
 \end{aligned}$$

8.  $a = 2.718, b = 3.142, A = 54^\circ 18'$  : SSA

By cosine rule  $a^2 = b^2 + c^2 - 2bc \cos A$

$$2.718^2 = 3.142^2 + c^2 - 2 \times 3.142 c \cos 54^\circ 18'$$


$$c^2 - 3.666973 c + 2.48464 = 0$$

$$c = 2.77 \text{ or } 0.90 \quad (B = 69.8^\circ \text{ or } 110.2^\circ, C = 55.9^\circ \text{ or } 15.6^\circ)$$

So, there are two possible triangles.

$$c = 2.77, \text{ area} = \frac{1}{2} bc \sin A = \frac{1}{2} \times 3.142 \times 2.77 \sin 54.3^\circ = 3.53$$

$$c = 0.90, \text{ area} = \frac{1}{2} \times 3.142 \times 0.90 \sin 54.3^\circ = 1.14$$



9.  $(\sin B - \cos B)^2 + (\sin C - \cos C)^2 - (\sin A - \cos A)^2$

$$= \sin^2 B - 2 \sin B \cos B + \cos^2 B + \sin^2 C - 2 \sin C \cos C + \cos^2 C - \sin^2 A + 2 \sin A \cos A - \cos^2 A$$

$$= 1 - \sin 2B + 1 - 2 \sin C \cos C - (1 - \sin 2A)$$

$$= 1 + \sin 2A - \sin 2B - 2 \sin C \cos C$$

$$= 1 + 2 \cos(A+B) \sin(A-B) - 2 \sin C \cos C$$

$$= 1 - 2 \cos C \sin(A-B) - 2 \sin C \cos C$$

$$= 1 - 2 \cos C [\sin(A-B) + \sin C]$$

$$= 1 - 4 \cos C \sin \frac{A-B+C}{2} \cos \frac{A-B-C}{2}$$

$$= 1 - 4 \cos C \sin(90^\circ - B) \cos(A - 90^\circ)$$

$$= 1 - 4 \sin A \cos B \cos C.$$

10. Given that  $(1 + \cos A)(1 + \cos B)(1 + \cos C)(1 + \cos D) = p \sin A \sin B \sin C \sin D$

$$\begin{aligned}
 &= \frac{(1 - \cos A)(1 - \cos B)(1 - \cos C)(1 - \cos D)}{(1 + \cos A)(1 + \cos B)(1 + \cos C)(1 + \cos D)} \\
 &= \frac{(1 - \cos^2 A)(1 - \cos^2 B)(1 - \cos^2 C)(1 - \cos^2 D)}{p \sin A \sin B \sin C \sin D} \\
 &= \frac{\sin^2 A \sin^2 B \sin^2 C \sin^2 D}{p \sin A \sin B \sin C \sin D} \\
 &= \frac{1}{p} \sin A \sin B \sin C \sin D.
 \end{aligned}$$

11. In the triangle on the right,

$$a = BC = BD + DC = c \cos B + b \cos C \dots\dots (1)$$

$$b = AC = AE + EC = c \cos A + a \cos C \dots\dots (2)$$

Similarly  $c = a \cos B + b \cos A \dots\dots (3)$  (do this part yourself)

These equations are known as projection formulae.

$$\text{In (2) } \cos C = \frac{b - c \cos A}{a} \dots\dots (4)$$

$$\text{In (3) } \cos B = \frac{c - b \cos A}{a} \dots\dots (5)$$

Put (4) and (5) into (1)

$$a = c \cdot \frac{c - b \cos A}{a} + b \cdot \frac{b - c \cos A}{a}$$

$$\Rightarrow a^2 = c^2 + b^2 - 2bc \cos A$$

$$\text{Similarly } b^2 = a^2 + c^2 - 2ac \cos B$$

$$c^2 = a^2 + b^2 - 2ab \cos C$$

(Do the substitution yourself.)

12.  $\theta + \varphi = \frac{1}{4}\pi$ ,  $\tan(\theta + \varphi) = \tan \frac{1}{4}\pi = 1$

$$\frac{\tan \theta + \tan \varphi}{1 - \tan \theta \tan \varphi} = 1$$

$$\tan \theta + \tan \varphi = 1 - \tan \theta \tan \varphi$$

$$1 + \tan \theta + \tan \varphi + \tan \theta \tan \varphi = 2$$

$$(1 + \tan \theta)(1 + \tan \varphi) = 2$$

$$\text{Let } \theta = \varphi = \frac{1}{8}\pi, \text{ then } \theta + \varphi = \frac{1}{4}\pi$$

$$\text{By the above result, } (1 + \tan \theta)(1 + \tan \varphi) = 2$$

$$\Rightarrow (1 + \tan \theta)^2 = 2$$

$$1 + \tan \theta = \pm \sqrt{2}$$

$$\tan \frac{1}{8}\pi = \sqrt{2} - 1 \text{ (reject } -\sqrt{2} - 1)$$

13.  $\sin \theta (\cos 2\theta + \cos 4\theta + \cos 6\theta)$

$$= \sin \theta (\cos 4\theta + 2\cos 4\theta \cos 2\theta)$$

$$= \cos 4\theta \sin \theta (2 \cos 2\theta + 1)$$

$$= \cos 4\theta [2(1 - 2\sin^2 \theta) + 1] \sin \theta$$

$$= \cos 4\theta (3 \sin \theta - 4 \sin^3 \theta)$$

$$= \sin 3\theta \cos 4\theta$$

If  $x = \cos 3\theta + \sin 3\theta$  and  $y = \cos \theta - \sin \theta$ ,

$$x - y = \cos 3\theta + \sin 3\theta - \cos \theta + \sin \theta$$

$$= 4 \cos^3 \theta - 3 \cos \theta + 3 \sin \theta - 4 \sin^3 \theta - \cos \theta + \sin \theta$$



$$= 4 \cos^3 \theta - 4 \cos \theta + 4 \sin \theta - 4 \sin^3 \theta$$

$$= 4 \cos \theta (\cos^2 \theta - 1) + 4 \sin^2 \theta (1 - \sin^2 \theta)$$

$$= -4 \cos \theta \sin^2 \theta + 4 \sin \theta \cos^2 \theta$$

$$= 4 \sin \theta \cos \theta (\cos \theta - \sin \theta)$$

$$= 2y \sin 2\theta.$$



14. Let  $E = (3 + \cos \theta) \cosec \theta$

$$\begin{aligned} &= (3 + \frac{1-t^2}{1+t^2}) \frac{1+t^2}{2t} \\ &= \frac{3+3t^2+1-t^2}{2t} \\ &= \frac{4+2t^2}{2t} \\ &= \frac{2+t^2}{t} \end{aligned}$$

$$Et = 2 + t^2$$

$$t^2 - Et + 2 = 0$$

$\therefore t = \tan \frac{\theta}{2}$  can be any real number,  $\Delta \geq 0$

$$E^2 - 4 \times 2 \geq 0$$

$$(E + 2\sqrt{2})(E - 2\sqrt{2}) \geq 0$$

$$\Rightarrow E \leq -2\sqrt{2} \text{ or } E \geq 2\sqrt{2}$$

This expression cannot have any value between  $-2\sqrt{2}$  and  $2\sqrt{2}$ .

15.  $\Delta PQR$  is an equilateral with  $QR$  inclined to the horizontal at an angle  $\theta$ .

Suppose  $PQ = QR = RP = 1$  unit.

The projection of  $QR$  on horizontal is  $MT$ , the projection of  $PQ$  on horizontal is  $MN$ , the projection of  $PR$  on horizontal is  $NT$ .

$$MT = QR \cos \theta = \cos \theta$$

$$MN = PQ \cos(\frac{1}{3}\pi + \theta) = \cos(\frac{1}{3}\pi + \theta)$$

$$NT = PR \cos(\frac{1}{3}\pi - \theta)$$

$$\therefore MT = MN + NT$$

$$\cos \theta = \cos(\frac{1}{3}\pi + \theta) + \cos(\frac{1}{3}\pi - \theta)$$

$$\cos \theta - \cos(\frac{1}{3}\pi + \theta) - \cos(\frac{1}{3}\pi - \theta) = 0$$

$$\cos \theta + \cos(\frac{1}{3}\pi + \theta + \pi) + \cos(\frac{1}{3}\pi - \theta - \pi) = 0$$

$$\cos \theta + \cos(\frac{4}{3}\pi + \theta) + \cos(-\frac{2}{3}\pi - \theta) = 0$$

$$\cos \theta + \cos(\theta + \frac{2}{3}\pi) + \cos(\theta + \frac{4}{3}\pi) = 0$$

Using the projection of  $\Delta PQR$  onto the vertical,

$$AB = PR \sin(\frac{1}{3}\pi - \theta) = \sin(\frac{1}{3}\pi - \theta)$$

$$BC = QR \sin \theta = \sin \theta$$

$$AC = PQ \sin(\frac{1}{3}\pi + \theta) = \sin(\frac{1}{3}\pi + \theta)$$

$$AB + BC = AC$$

$$\sin(\frac{1}{3}\pi - \theta) + \sin \theta = \sin(\frac{1}{3}\pi + \theta)$$

$$\sin(\frac{1}{3}\pi - \theta) + \sin \theta - \sin(\frac{1}{3}\pi + \theta) = 0$$

$$\sin \theta + \sin[\pi - (\frac{1}{3}\pi - \theta)] + \sin(\frac{1}{3}\pi + \theta + \pi) = 0$$

$$\sin \theta + \sin(\theta + \frac{2}{3}\pi) + \sin(\theta + \frac{4}{3}\pi) = 0$$

16.  $\tan \alpha = k \tan \beta$



$$\frac{\sin \alpha}{\cos \alpha} = \frac{k \sin \beta}{\cos \beta}$$

$$\sin \alpha \cos \beta = k \sin \beta \cos \alpha$$

$$\frac{1}{2} [\sin(\alpha + \beta) + \sin(\alpha - \beta)] = \frac{k}{2} [\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

$$\sin(\alpha - \beta) + k \sin(\alpha - \beta) = k \sin(\alpha + \beta) - \sin(\alpha + \beta)$$

$$(k-1) \sin(\alpha + \beta) = (k+1) \sin(\alpha - \beta)$$



Given  $\tan x = k \tan(x - \alpha)$

$$(k-1) \sin(2x - \alpha) = (k+1) \sin \alpha$$

$$\sin(2x - \alpha) = \frac{(k+1)}{(k-1)} \sin \alpha$$

$$\text{It has real solutions (in } x\text{)} \Rightarrow -1 \leq \frac{(k+1)}{(k-1)} \sin \alpha \leq 1$$

$$\frac{(k+1)^2}{(k-1)^2} \sin^2 \alpha \leq 1$$

$$(k+1)^2 \sin^2 \alpha \leq (k-1)^2$$

$(k-1)^2$  is not less than  $(k+1)^2 \sin^2 \alpha$ .

When  $k = -2$  and  $\alpha = 30^\circ$ ,

$$(-2-1) \sin(2x - 30^\circ) = (-2+1) \sin 30^\circ$$

$$\sin(2x - 30^\circ) = \frac{1}{6}$$

$$2x - 30^\circ = 180^\circ n + (-1)^n 9.594^\circ$$

$$2x = 180^\circ n + (-1)^n 9.594^\circ + 30^\circ$$

$$x = 90^\circ n + (-1)^n 4.797^\circ + 15^\circ, n = 0, \pm 1, \pm 2, \pm 3, \dots$$

17. (a) If  $\sin \alpha + \cos \alpha = 2a$

$$\text{Squaring: } \sin^2 \alpha + \cos^2 \alpha + 2 \sin \alpha \cos \alpha = 4a^2$$

$$2 \sin \alpha \cos \alpha = 4a^2 - 1$$

$$\sin \alpha \cos \alpha = 2a^2 - \frac{1}{2}$$

$$\text{sum of roots} = 2a, \text{ product of roots} = 2a^2 - \frac{1}{2}$$

$$\text{Quadratic equation whose roots are } \sin \alpha, \cos \alpha \text{ is } x^2 - 2ax + 2a^2 - \frac{1}{2} = 0$$

$$2x^2 - 4ax + 4a^2 - 1 = 0$$

$$(b) \cos^2 x + \cos x - \sin x - \sin^2 x = 0$$

$$\cos^2 x - \sin^2 x + \cos x - \sin x = 0$$

$$(\cos x + \sin x)(\cos x - \sin x) + \cos x - \sin x = 0$$

$$(\cos x - \sin x)(\cos x + \sin x + 1) = 0$$

$$\cos x - \sin x = 0 \text{ or } \cos x + \sin x = -1$$

$$\tan x = 1 \text{ or } \frac{1}{\sqrt{2}} \cos x + \frac{1}{\sqrt{2}} \sin x = -\frac{1}{\sqrt{2}}$$

$$x = 180^\circ n + 45^\circ \text{ or } \cos(x - 45^\circ) = \cos 135^\circ$$

$$x = 180^\circ n + 45^\circ \text{ or } x - 45^\circ = 360^\circ n \pm 135^\circ$$

$$x = 180^\circ n + 45^\circ \text{ or } x = 360^\circ n + 180^\circ \text{ or } 360^\circ n - 90^\circ, n = 0, \pm 1, \pm 2, \pm 3, \dots$$

18. Let  $y = \sin \theta (3 \sin \theta - \sin 2\alpha) + \cos \theta (3 \cos \theta - \cos 2\alpha)$  ( $0^\circ < \alpha < 90^\circ$ ).

$$= 3 \sin^2 \theta - \sin \theta \sin 2\alpha + 3 \cos^2 \theta - \cos \theta \cos 2\alpha$$

$$= 3 - (\sin \theta \sin 2\alpha + \cos \theta \cos 2\alpha)$$

$$= 3 - \cos(\theta - 2\alpha)$$

- (a) When  $y$  has a minimum value,  $\cos(\theta - 2\alpha) = 1$

$$\theta - 2\alpha = 360^\circ n$$

$$\theta = 360^\circ n + 2\alpha, n = 0, \pm 1, \pm 2, \pm 3, \dots$$

- (b) When  $y$  has a maximum value,  $\cos(\theta - 2\alpha) = -1$

$$\theta - 2\alpha = 360^\circ n + 180^\circ$$

$$\theta = 360^\circ n + 180^\circ + 2\alpha, n = 0, \pm 1, \pm 2, \pm 3, \dots$$

(c) Interchange the role of  $\alpha$  and  $\theta$ , we have

$$\sin \alpha (3 \sin \alpha - \sin 2\theta) + \cos \alpha (3 \cos \alpha - \cos 2\theta) = 3 - \cos(\alpha - 2\theta)$$

Maximum value is  $3 - (-1) = 4$

Minimum value is  $3 - 1 = 2$

19.  $a \cos \theta + b \sin \theta = c$ ,  $\alpha, \beta$  are two distinct roots.

$$a \cos \alpha + b \sin \alpha = c \quad \dots \dots (1)$$

$$a \cos \beta + b \sin \beta = c \quad \dots \dots (2)$$

$$(1) = (2) \quad a \cos \alpha + b \sin \alpha = a \cos \beta + b \sin \beta$$

$$a(\cos \alpha - \cos \beta) = b(\sin \beta - \sin \alpha)$$

$$-2a \sin \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2} = -2b \cos \frac{\alpha + \beta}{2} \sin \frac{\alpha - \beta}{2}$$

$$\therefore \alpha, \beta \text{ are distinct} \therefore \sin \frac{\alpha - \beta}{2} \neq 0$$

$$a \sin \frac{\alpha + \beta}{2} = b \cos \frac{\alpha + \beta}{2}$$

$$\tan \frac{\alpha + \beta}{2} = \frac{b}{a}$$

Using the formula for half angle:  $t = \tan \frac{\alpha + \beta}{2}$

$$\begin{aligned} \frac{a}{b} \sin(\alpha + \beta) - \cos(\alpha + \beta) &= \frac{a}{b} \cdot \frac{2t}{1+t^2} - \frac{1-t^2}{1+t^2} \\ &= \frac{a}{b} \cdot \left[ \frac{2\left(\frac{b}{a}\right)}{1+\left(\frac{b}{a}\right)^2} \right] - \frac{1-\left(\frac{b}{a}\right)^2}{1+\left(\frac{b}{a}\right)^2} \\ &= \frac{a}{b} \cdot \frac{2ab}{a^2+b^2} - \frac{a^2-b^2}{a^2+b^2} \\ &= \frac{2a^2-a^2+b^2}{a^2+b^2} = 1 \end{aligned}$$

20.  $\frac{\cos(2\theta + \varphi) + \cos(2\varphi + \theta)}{2 \cos(\theta + \varphi) - 1} = \frac{\cos(2\theta - \varphi) + \cos(2\varphi - \theta)}{2 \cos(\theta - \varphi) - 1}$  is equivalent to

$$[2 \cos(\theta - \varphi) - 1][\cos(2\theta + \varphi) + \cos(2\varphi + \theta)] = [2 \cos(\theta + \varphi) - 1][\cos(2\theta - \varphi) + \cos(2\varphi - \theta)]$$

$$\text{LHS} = 2 \cos(2\theta + \varphi) \cos(\theta - \varphi) + 2 \cos(2\varphi + \theta) \cos(\theta - \varphi) - [\cos(2\theta + \varphi) + \cos(2\varphi + \theta)]$$

$$= \cos 3\theta + \cos(\theta + 2\varphi) + \cos(2\theta + \varphi) + \cos 3\varphi - \cos(2\theta + \varphi) - \cos(2\varphi + \theta)$$

$$= \cos 3\theta + \cos 3\varphi$$

$$\text{RHS} = 2 \cos(2\theta - \varphi) \cos(\theta + \varphi) + 2 \cos(2\varphi - \theta) \cos(\theta + \varphi) - [\cos(2\theta - \varphi) + \cos(2\varphi - \theta)]$$

$$= \cos 3\theta + \cos(\theta - 2\varphi) + \cos 3\varphi + \cos(\varphi - 2\theta) - \cos(2\theta - \varphi) - \cos(2\varphi - \theta)$$

$$= \cos 3\theta + \cos 3\varphi$$

$\therefore \text{LHS} = \text{RHS}$

21. (a)  $2 \cos(\theta - \varphi) \sin(2\theta + \varphi) \sin(2\varphi + \theta)$


$$= -\cos(\theta - \varphi)[\cos(3\theta + 3\varphi) - \cos(\theta - \varphi)]$$

$$= -\cos(\theta - \varphi)\cos(3\theta + 3\varphi) + \cos^2(\theta - \varphi)$$

$$= -\frac{1}{2} [\cos(4\theta + 2\varphi) + \cos(2\theta + 4\varphi)] + 1 - \sin^2(\theta - \varphi)$$

$$= -\frac{1}{2} [1 - 2 \sin^2(2\theta + \varphi) + 1 - 2 \sin^2(\theta + 2\varphi)] + 1 - \sin^2(\theta - \varphi)$$

$$= \sin^2(2\theta + \varphi) + \sin^2(2\varphi + \theta) - \sin^2(\theta - \varphi) \text{ (similar to Q6(a))}$$



$$\begin{aligned}
 (b) & \quad \tan(3A - \frac{3}{4}\pi) \tan(A + \frac{1}{4}\pi) \\
 &= \frac{\sin(3A - \frac{3\pi}{4}) \sin(A + \frac{\pi}{4})}{\cos(3A - \frac{3\pi}{4}) \cos(A + \frac{\pi}{4})} \\
 &= \frac{-\frac{1}{2}[\cos(4A - \frac{\pi}{2}) - \cos(2A - \pi)]}{\frac{1}{2}[\cos(4A - \frac{\pi}{2}) + \cos(2A - \pi)]} \\
 &= \frac{-(\sin 4A + \cos 2A)}{\sin 4A - \cos 2A} \\
 &= -\frac{2\sin 2A \cos 2A + \cos 2A}{2\sin 2A \cos 2A - \cos 2A} \\
 &= -\frac{2\sin 2A + 1}{2\sin 2A - 1} \\
 &= \frac{1 + 2\sin 2A}{1 - 2\sin 2A}
 \end{aligned}$$

$$\begin{aligned}
 (c) & \quad (\cos A + \cos B + \cos C)^2 + (\sin A + \sin B + \sin C)^2 \\
 &= \cos^2 A + \cos^2 B + \cos^2 C + \sin^2 A + \sin^2 B + \sin^2 C + 2(\cos A \cos B + \sin A \sin B \\
 &\quad + \cos B \cos C + \sin B \sin C + \cos C \cos A + \sin A \sin C) \\
 &= 3 + 2[\cos(A - B) + \cos(B - C) + \cos(C - A)] \\
 &= 3 + 4 \cos \frac{A-C}{2} \cos \frac{A-2B+C}{2} + 4 \cos^2 \frac{C-A}{2} - 2 \\
 &= 1 + 4 \cos \frac{C-A}{2} [\cos \frac{A-2B+C}{2} + \cos \frac{C-A}{2}] \\
 &= 1 + 8 \cos \frac{C-A}{2} \cos \frac{C-B}{2} \cos \frac{A-B}{2} = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 (d) & \quad \frac{\tan 3A}{\tan A} \\
 &= \frac{\sin 3A}{\cos 3A} \cdot \frac{\cos A}{\sin A} \\
 &= \frac{3\sin A - 4\sin^3 A}{4\cos^3 A - 3\cos A} \cdot \frac{\cos A}{\sin A} \\
 &= \frac{3 - 4\sin^2 A}{4\cos^2 A - 3} \\
 &= \frac{3 - 2(1 - \cos 2A)}{2(1 + \cos 2A) - 3} \\
 &= \frac{2\cos 2A + 1}{2\cos 2A - 1},
 \end{aligned}$$

$$\begin{aligned}
 (e) & \quad \sin^2(A + \theta) + \sin^2(B + \theta) \\
 &= \frac{1}{2}[1 - \cos(2A + 2\theta)] + \frac{1}{2}[1 - \cos(2B + 2\theta)] \\
 &= 1 - \frac{1}{2}[\cos(2A + 2\theta) + \cos(2B + 2\theta)] \\
 &= 1 - \cos(A - B) \cos(A + B + 2\theta)
 \end{aligned}$$