


## Polar area example 2

### 1993 HKAL Pure Mathematics Paper 2 Q10 (c)

Created by Mr. Francis Hung on 20181027

Last updated: February 26, 2023

Given  $x^3 + y^3 = 3axy$ , find the area of the loop.



$$x = r \cos \theta, y = r \sin \theta, \text{ sub. into } x^3 + y^3 = 3axy$$

$$(r \cos \theta)^3 + (r \sin \theta)^3 = 3a(r \cos \theta)(r \sin \theta)$$

$$\therefore \text{The polar equation of } \Gamma \text{ is } r = \frac{3a \sin \theta \cos \theta}{\sin^3 \theta + \cos^3 \theta}$$

$$\frac{3a \sin(\pi + \theta) \cos(\pi + \theta)}{\sin^3(\pi + \theta) + \cos^3(\pi + \theta)} = \frac{-3a \sin \theta \cos \theta}{\sin^3 \theta + \cos^3 \theta} \Rightarrow \text{the curve repeats for every multiple of } \pi.$$

$$r = \frac{3a \sin \theta \cos \theta}{\sin^3 \theta + \cos^3 \theta} = \frac{3a \sin \theta \cos \theta}{(\sin \theta + \cos \theta)(\sin^2 \theta - \sin \theta \cos \theta + \cos^2 \theta)} = \frac{\frac{3a}{2} \sin 2\theta}{\sqrt{2} \sin(\theta + \frac{\pi}{4})(1 - \frac{1}{2} \sin 2\theta)}$$

$$1 - \frac{1}{2} \sin 2\theta \geq 1 - \frac{1}{2} = \frac{1}{2} > 0$$

$$r \text{ is undefined when } \theta = \frac{3\pi}{4}.$$

When  $\theta \rightarrow \left(\frac{3\pi}{4}\right)^-$ ,  $r \rightarrow -\infty$ ; when  $\theta \rightarrow \left(\frac{3\pi}{4}\right)^+$ ,  $r \rightarrow \infty$ .

$\theta$	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{5\pi}{12}$	$\frac{\pi}{2}$	$\frac{7\pi}{12}$	$\frac{2\pi}{3}$	$\left(\frac{3\pi}{4}\right)^-$	$\left(\frac{3\pi}{4}\right)^+$	$\frac{5\pi}{6}$	$\frac{11\pi}{12}$	$\pi$
	$0^\circ$	$15^\circ$	$30^\circ$	$45^\circ$	$60^\circ$	$75^\circ$	$90^\circ$	$105^\circ$	$120^\circ$	$134^\circ$	$136^\circ$	$150^\circ$	$165^\circ$	$180^\circ$
$r$	0	$0.8a$	$1.7a$	$2.1a$	$1.7a$	$0.8a$	0	$-0.8a$	$-2.5a$	$-40a$	$40a$	$2.5a$	$0.8a$	0

From the table, the small loop corresponds to  $0 \leq \theta \leq \frac{\pi}{2}$ .

$$\text{Area} = \frac{1}{2} \int_0^{\frac{\pi}{2}} r^2 d\theta = \int_0^{\frac{\pi}{4}} \left( \frac{3a \sin \theta \cos \theta}{\sin^3 \theta + \cos^3 \theta} \right)^2 d\theta = 9a^2 \int_0^{\frac{\pi}{4}} \left( \frac{\sin \theta \cos \theta}{\sin^3 \theta + \cos^3 \theta} \right)^2 d\theta$$

$$= 9a^2 \int_0^{\frac{\pi}{4}} \frac{\tan \theta \sec \theta}{(\tan^3 \theta + 1)^2} d\theta = 3a^2 \int_1^2 \frac{dw}{w^2}, w = 1 + \tan^3 \theta, dw = 3 \tan^2 \theta \sec^2 \theta d\theta$$

$$= 3a^2 \left( -\frac{1}{w} \right)_1^2 = 3a^2 \left( -\frac{1}{2} + 1 \right) = \frac{3a^2}{2} \text{ sq. units}$$

**Method 2**

Perform the rotation of axis (in anti-clockwise direction by  $45^\circ$ ):

$$\begin{cases} x = \frac{1}{\sqrt{2}}(x_1 - y_1) \\ y = \frac{1}{\sqrt{2}}(x_1 + y_1) \end{cases}$$

$$\left[ \frac{1}{\sqrt{2}}(x_1 - y_1) \right]^3 + \left[ \frac{1}{\sqrt{2}}(x_1 + y_1) \right]^3 = 3a \frac{1}{\sqrt{2}}(x_1 - y_1) \frac{1}{\sqrt{2}}(x_1 + y_1)$$

$$\frac{1}{2\sqrt{2}}(2x_1^3 + 6x_1y_1^2) = \frac{3a}{2}(x_1^2 - y_1^2)$$

$$2x_1^3 + 6x_1y_1^2 = 3a\sqrt{2}(x_1^2 - y_1^2)$$


$$(6x_1 + 3a\sqrt{2})y_1^2 = 3a\sqrt{2}x_1^2 - 2x_1^3$$

$$y_1^2 = \frac{3a\sqrt{2}x_1^2 - 2x_1^3}{6x_1 + 3a\sqrt{2}}$$

$$y_1^2 = \frac{(3a\sqrt{2} - 2x_1)x_1^2}{3(2x_1 + a\sqrt{2})}$$

$$y_1^2 = \frac{(3a - \sqrt{2}x_1)x_1^2}{3(\sqrt{2}x_1 + a)}$$

$$y_1 = \pm \frac{x_1}{\sqrt{3}} \cdot \sqrt{\frac{3a - \sqrt{2}x_1}{\sqrt{2}x_1 + a}}$$



After rotation, the area is  $2 \times \int_0^{\frac{3a}{\sqrt{2}}} y_1 dx_1$

$$\begin{aligned}
 &= 2 \int_0^{\frac{3a}{\sqrt{2}}} \frac{x}{\sqrt{3}} \cdot \sqrt{\frac{3a - \sqrt{2}x}{\sqrt{2}x + a}} dx \\
 &= \sqrt{2} \int_0^{\frac{3a}{\sqrt{2}}} \frac{x}{\sqrt{3}} \cdot \sqrt{\frac{3a - \sqrt{2}x}{\sqrt{2}x + a}} d(\sqrt{2}x) \\
 &= \sqrt{2} \int_0^{3a} \frac{t}{\sqrt{2}\sqrt{3}} \cdot \sqrt{\frac{3a - t}{t + a}} dt, t = \sqrt{2}x \\
 &= \frac{1}{\sqrt{3}} \int_0^{3a} t \cdot \sqrt{\frac{3a - t}{t + a}} dt
 \end{aligned}$$

Let  $u = \sqrt{t+a}$ ;  $t=0, u=\sqrt{a}$ ;  $t=3a, u=2\sqrt{a}$

$$du = \frac{dt}{2\sqrt{t+a}}; t=u^2-a$$

$$\begin{aligned}
 \text{Area} &= \frac{1}{\sqrt{3}} \int_{\sqrt{a}}^{2\sqrt{a}} (u^2 - a) \cdot \sqrt{3a - u^2 + a} (2du) \\
 &= \frac{2}{\sqrt{3}} \int_{\sqrt{a}}^{2\sqrt{a}} (u^2 - a) \cdot \sqrt{4a - u^2} du
 \end{aligned}$$

Let  $u = 2\sqrt{a} \sin \theta$ ,  $\sqrt{4a - u^2} = 2\sqrt{a} \cos \theta$ ;  $du = 2\sqrt{a} \cos \theta d\theta$

$$u = \sqrt{a}, \theta = \frac{\pi}{6}; u = 2\sqrt{a}, \theta = \frac{\pi}{2}$$

$$\begin{aligned}
 \text{Area} &= \frac{2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} (4a \sin^2 \theta - a) \cdot (2\sqrt{a} \cos \theta)^2 d\theta \\
 &= \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} (4 \sin^2 \theta - 1) \cdot \cos^2 \theta d\theta \\
 &= \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} 4 \sin^2 \theta \cos^2 \theta d\theta - \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \cos^2 \theta d\theta \\
 &= \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \sin^2 2\theta d\theta - \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{1 + \cos 2\theta}{2} d\theta \\
 &= \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{1 - \cos 4\theta}{2} d\theta - \frac{8a^2}{\sqrt{3}} \int_{\frac{\pi}{6}}^{\frac{\pi}{2}} \frac{1 + \cos 2\theta}{2} d\theta \\
 &= \frac{4a^2}{\sqrt{3}} \left( \theta - \frac{\sin 4\theta}{4} \right) \Big|_{\frac{\pi}{6}}^{\frac{\pi}{2}} - \frac{4a^2}{\sqrt{3}} \left( \theta + \frac{\sin 2\theta}{2} \right) \Big|_{\frac{\pi}{6}}^{\frac{\pi}{2}} \\
 &= \frac{4a^2}{\sqrt{3}} \left( -\frac{\sin 4\theta}{4} - \frac{\sin 2\theta}{2} \right) \Big|_{\frac{\pi}{6}}^{\frac{\pi}{2}} \\
 &= \frac{4a^2}{\sqrt{3}} \left( \frac{\sin \frac{2\pi}{3}}{4} + \frac{\sin \frac{\pi}{3}}{2} \right) = \frac{a^2}{\sqrt{3}} \left( \sin \frac{2\pi}{3} + 2 \sin \frac{\pi}{3} \right) = \frac{a^2}{\sqrt{3}} \left( \frac{3\sqrt{3}}{2} \right) = \frac{3a^2}{2}
 \end{aligned}$$