

Fixed Point Theorem Example

Created by Mr. Francis Hung on 20111124.

Last updated: 12 February 2022

Supplementary Problem on Root Approximation

In this problem you may assumed that if $-1 < M < 1$, then $\lim_{n \rightarrow \infty} M^n = 0$.

- (a) If $a < b$, let $g(x)$ be a continuous differentiable function defined on $[a, b]$. Suppose $y = g(x)$ satisfies the following 2 conditions:

- (1) $a \leq g(x) \leq b$ for all $x: a \leq x \leq b$,
- (2) $|g'(x)| \leq M < 1$ for all $x: a \leq x \leq b$.

If $x_{n+1} = g(x_n)$, prove that $\lim_{n \rightarrow \infty} g(x_n) = \lim_{n \rightarrow \infty} x_n$ and $a \leq \lim_{n \rightarrow \infty} g(x_n) \leq b$.

- (b) Let $g(x) = \frac{2}{x}$, $1 \leq x \leq 2$. Show that fixed-point method fails.
- (c) Let $g(x) = \frac{1}{x^2} - \frac{10}{x}$, $3.1 \leq x \leq 3.2$. Show that fixed-point method also fails.
- (d) Given that a root of the equation $x + \ln x = 0$ is α . Show that $0.4 \leq \alpha \leq 0.8$. Here are three iterative formulae:

(i) $x_{r+1} = -\ln x_r$,

(ii) $x_{r+1} = e^{-x_r}$

(iii) $x_{r+1} = \frac{x_r + e^{-x_r}}{2}$

(α) Which of the formula can be used?

(β) Which formula is better? Use it to find the root correct to 3 decimal places.

Solution

- (a) Let $h(x) = g(x) - x$

then $h(a) = g(a) - a \geq 0$

$h(b) = g(b) - b \geq 0$

\therefore there is a root $\alpha: a < \alpha < b$ such that $h(\alpha) = 0$

$\Rightarrow g(\alpha) = \alpha$

By mean value theorem, $g(x_n) - g(\alpha) = (x_n - \alpha)g'(c)$, $\alpha < c < x_n$

$|x_{n+1} - \alpha| = |g(x_n) - g(\alpha)|$

$= |x_n - \alpha||g'(c)|$

$|x_{n+1} - \alpha| \leq |x_n - \alpha|M$

$\therefore |x_n - \alpha| \leq |x_{n-1} - \alpha|M$

$|x_{n-1} - \alpha| \leq |x_{n-2} - \alpha|M$

.....

$\times) |x_1 - \alpha| \leq |x_0 - \alpha|M$

$|x_n - \alpha| \leq |x_0 - \alpha|M^n$

Take limit $n \rightarrow \infty$

$\lim_{n \rightarrow \infty} |x_n - \alpha| \leq |x_0 - \alpha| \lim_{n \rightarrow \infty} M^n$

$\left| \lim_{n \rightarrow \infty} (x_n - \alpha) \right| \leq |x_0 - \alpha| \cdot 0 = 0$

$\Rightarrow \lim_{n \rightarrow \infty} x_n = \alpha = g(\alpha) = g(\lim_{n \rightarrow \infty} x_n)$

$\Rightarrow \lim_{n \rightarrow \infty} g(x_n) = \alpha$ ($\because g$ is continuous.)

$\Rightarrow \lim_{n \rightarrow \infty} g(x_n) = \lim_{n \rightarrow \infty} x_n$

$\therefore a \leq \alpha \leq b$

$\therefore a \leq \lim_{n \rightarrow \infty} g(x_n) \leq b$.

(b) $g(x) = \frac{2}{x} \quad 1 \leq x \leq 2$

$$g'(x) = -\frac{2}{x^2} < 0 \Rightarrow g \text{ is strictly decreasing on } [1, 2]$$

$$g(1) = 2, g(2) = 1, \therefore 1 \leq g(x) \leq 2$$

\therefore condition (1) is satisfied.

$$|g'(x)| = \frac{2}{x^2}$$

$$\frac{2}{4} \leq |g'(x)| \leq \frac{2}{1}$$

$$\Rightarrow \frac{1}{2} \leq |g'(x)| \leq 2, \text{ In particular, } |g'(1.25)| = 1.28 > 1$$

condition (2) fails.

(c) $g(x) = \frac{1}{x^2} - \frac{10}{x}, 3.1 \leq x \leq 3.2$

$$g(3.1) = -3.12 \notin [3.1, 3.2]$$

$$g(3.2) = -3.027 \notin [3.1, 3.2]$$

condition (1) fails

$$g'(x) = -\frac{2}{x^3} + \frac{10}{x^2}$$

$$g''(x) = \frac{6}{x^4} - \frac{20}{x^3}$$

$$\text{Let } g''(x) = 0$$

$$\Rightarrow \frac{6}{x^4} - \frac{20}{x^3} = 0$$

$$\Rightarrow 6x^3 = 20x^4$$

$$\Rightarrow 6 = 20x$$

$$\Rightarrow x = 0.3$$

$$\therefore 3.1 \leq x \leq 3.2$$

$$\therefore g''(x) \neq 0 \quad \forall x \in [3.1, 3.2]$$

$$g''(3.1) = -0.61 < 0$$

$\therefore g'(x)$ is strictly decreasing

$$g'(3.2) < g'(x) < g'(3.1)$$

$$g'(3.1) = 0.97, g'(3.2) = 0.92$$

$$\therefore 0.92 < g'(x) < 0.97$$

$$\Rightarrow |g'(x)| \leq 0.98 \quad \forall x \in [3.1, 3.2]$$

condition (2) is satisfied.

(d) $f(x) = x + \ln x$

$$f(0.4) = -0.52 < 0$$

$$f(0.8) = 0.58 > 0$$

\therefore there is a root $\alpha: 0.4 < \alpha < 0.8$ such that $f(\alpha) = 0$

$$f'(x) = 1 + \frac{1}{x}$$

$$\therefore f'(x) > 0 \quad \forall x \in [0.4, 0.8]$$

$\therefore f(x)$ is strictly increasing and so that root α is unique.

(α) (i) $x_{r+1} = -\ln x_r$

$$g(x) = -\ln x$$

$$g'(x) = -\frac{1}{x}$$

$$g'(0.4) = -2.5; g'(0.8) = -1.25$$

\therefore condition (2) fails

- (ii) $x_{r+1} = e^{-x_r}$
 $g(x) = e^{-x}$
 $g'(x) = -e^{-x} < 0 \Rightarrow g(x)$ is strictly decreasing
 $g'(0.4) = -0.67$
 $g'(0.8) = -0.45$
 $g'(0.8) < g'(x) < g'(0.4)$ for $x: 0.4 < x < 0.8$
 $|g'(x)| \leq 0.7 < 1$
condition (2) is satisfied.
 $g(0.4) = 0.67, g(0.8) = 0.45$
 $0.4 < g(x) < 0.8$ for all $x: 0.4 < x < 0.8$
condition (1) is satisfied.
Fixed-point method can be applied.

- (iii) $x_{r+1} = \frac{x_r + e^{-x_r}}{2}$
 $g(x) = \frac{x + e^{-x}}{2}$
 $g(0.4) = 0.54; g(0.8) = 0.62$
 $g'(x) = \frac{1}{2}(1 - e^{-x}) > 0$ for all $x \in [0.4, 0.8]$.
 $g(x)$ is strictly increasing.
 $g(0.4) \leq g(x) \leq g(0.8)$
 $0.54 \leq g(x) \leq 0.62$
 $0.4 \leq g(x) \leq 0.8$
condition (1) is satisfied.
 $g'(0.4) = 0.16; g'(0.8) = 0.28$
 $g''(x) = \frac{1}{2}e^{-x} > 0$ for all x
 $g'(x)$ is strictly increasing on $[0.4, 0.8]$.
 $g'(0.4) \leq g'(x) \leq g'(0.8)$
 $0.16 \leq g'(x) \leq 0.28$
 $\therefore |g'(x)| \leq 0.3 < 1$
condition (2) is satisfied
Fixed-point method can be applied.

(β) For formula (ii), $M = 0.7$; for formula (iii), $M = 0.3$.

\therefore Formula (iii) is better.

Here is a comparison:

$$\text{Let } x_0 = \frac{0.4 + 0.8}{2} = 0.6$$

| Formula (ii) | | Formula (iii) | |
|---|----------------------|--|--------------------------------------|
| n | $x_{r+1} = e^{-x_r}$ | n | $x_{r+1} = \frac{x_r + e^{-x_r}}{2}$ |
| 0 | 0.6 | 0 | 0.6 |
| 1 | 0.5488 | 1 | 0.5744 |
| 2 | 0.5776 | 2 | 0.5687 |
| 3 | 0.5612 | 3 | 0.5674 |
| 4 | 0.5705 | 4 | 0.5672 |
| 5 | 0.5652 | | |
| 6 | 0.5682 | | |
| 7 | 0.5665 | | |
| 8 | 0.5675 | | |
| 9 | 0.5669 | | |
| 10 | 0.5673 | | |
| converges to 0.567 (3 d.p.) in 10 steps | | converges to 0.567 (3 d.p.) in 4 steps | |