

Hyperbolic Functions

First created by Mr. Francis Hung in 1989, retyped as MS WORD document on 20080609.

Last updated: 13 February 2022

Reference: Techniques of Mathematical Analysis by C.T. Tranter p. 160 - p.165, p.263 - p.266

Euler's Function: $e^{ix} = \cos x + i \sin x$; $e^{-ix} = \cos x - i \sin x$

$$\cos x = \frac{1}{2}(e^{ix} + e^{-ix}); \sin x = \frac{1}{2i}(e^{ix} - e^{-ix})$$

$$e^{x+iy} = e^x (\cos y + i \sin y)$$

Hyperbolic Function: $\cosh z = \frac{1}{2}(e^z + e^{-z})$

$$\sinh z = \frac{1}{2}(e^z - e^{-z})$$

$$\tanh z = \frac{\sinh z}{\cosh z} = \frac{e^z - e^{-z}}{e^z + e^{-z}}$$

$$\cosh z + \sinh z = e^z$$

$$\cosh z = 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \cdots + \frac{z^{2n}}{(2n)!} + \cdots;$$

$$\operatorname{sech} z = \frac{1}{\cosh z} = \frac{2}{e^z + e^{-z}}$$


$$\operatorname{cosech} z = \frac{1}{\sinh z} = \frac{2}{e^z - e^{-z}}$$

$$\coth z = \frac{1}{\tanh z} = \frac{e^z + e^{-z}}{e^z - e^{-z}}$$

$$\cosh z - \sinh z = e^{-z}$$

$$\sinh z = z + \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots + \frac{z^{2n-1}}{(2n-1)!} + \cdots$$

Graphs



Relation with real trigonometric functions:

$$\cosh z = \cos iz ;$$

$$\sinh z = -i \sin iz$$

$$\tanh z = -i \tan iz ;$$

$$\coth z = i \cot iz$$

$$\operatorname{sech} z = \sec iz ;$$

$$\operatorname{cosech} z = i \operatorname{cosec} iz$$

Identities:

$$\cosh^2 z - \sinh^2 z = 1$$

$$\operatorname{sech}^2 z = 1 - \tanh^2 z$$

$$\operatorname{cosech}^2 z = -1 + \coth^2 z$$

Exercise: Find $\sinh(-z)$; $\cosh(-z)$; $\sinh 0$; $\cosh 0$.

Addition Formulae: $\sinh(x \pm y) = \sinh x \cosh y \pm \cosh x \sinh y$

$$\cosh(x \pm y) = \cosh x \cosh y \pm \sinh x \sinh y$$

$$\tanh(x \pm y) = \frac{\tanh x \pm \tanh y}{1 \pm \tanh x \tanh y}$$

Double Angle Formulae: $\sinh 2z = 2 \sinh z \cosh z$

$$\cosh 2z = \cosh^2 z + \sinh^2 z = 2 \cosh^2 z - 1 = 2 \sinh^2 z + 1$$

Special manipulation: If x and y are real numbers,

$$\sin(x+iy) = \sin x \cosh y + i \cos x \sinh y$$

$$\cos(x+iy) = \cos x \cosh y - i \sin x \sinh y$$

$$\sinh(x+iy) = \sinh x \cos y + i \cosh x \sin y$$

$$\cosh(x+iy) = \cosh x \cos y + i \sinh x \sin y$$

Inverse hyperbolic function: z is a complex number, $\tanh^{-1} z = \frac{1}{2} \ln\left(\frac{1+z}{1-z}\right)$ for $-1 < z < 1$

$$\sinh^{-1} x = \ln\left(x + \sqrt{x^2 + 1}\right)$$

$$\operatorname{cosech}^{-1} x = \ln\left(\frac{1 \pm \sqrt{1+x^2}}{x}\right), x \neq 0$$

($x > 0$, take +; $x < 0$, take -)

$$\cosh^{-1} x = \ln\left(x + \sqrt{x^2 - 1}\right), x \geq 1$$

$$\operatorname{sech}^{-1} x = \ln\left(\frac{1 + \sqrt{1-x^2}}{x}\right), 0 < x \leq 1$$

$$\tanh^{-1} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right), -1 < x < 1$$

$$\operatorname{coth}^{-1} x = \frac{1}{2} \ln\left(\frac{x+1}{x-1}\right), |x| > 1$$

Differentiation:

$$\frac{d \sinh x}{dx} = \cosh x$$

$$\frac{d \operatorname{cosech} x}{dx} = -\operatorname{cosech} x \coth x$$

$$\frac{d \cosh x}{dx} = \sinh x$$

$$\frac{d \operatorname{sech} x}{dx} = -\operatorname{sech} x \tanh x$$

$$\frac{d \tanh x}{dx} = \operatorname{sech}^2 x$$

$$\frac{d \coth x}{dx} = -\operatorname{cosech}^2 x$$

$$\frac{d \sinh^{-1} x}{dx} = \frac{1}{\sqrt{1+x^2}}$$

$$\frac{d \cosh^{-1} x}{dx} = \frac{1}{\sqrt{x^2-1}}, \text{ for all } x > 1$$

$$\frac{d \tanh^{-1} x}{dx} = \frac{1}{1-x^2}, \text{ for } |x| < 1$$

$$\frac{d \coth^{-1} x}{dx} = \frac{1}{1-x^2}, |x| > 1$$

$$\frac{d \operatorname{sech}^{-1} x}{dx} = -\frac{1}{x\sqrt{1-x^2}}, 0 < x < 1$$

$$\frac{d \operatorname{cosech}^{-1} x}{dx} = \frac{-1}{|x|\sqrt{1+x^2}}, \text{ for } x \neq 0$$

Integration: The inverse process of differentiation

$$\int \sinh x dx = \cosh x + c$$

$$\int \operatorname{cosech} x \coth x dx = -\operatorname{cosech} x + c$$

$$\int \cosh x dx = \sinh x + c$$

$$\int \sec h x \tanh x dx = -\operatorname{sech} x + c$$

$$\int \operatorname{sech}^2 x dx = \tanh x + c$$

$$\int \cos \operatorname{cosech}^2 x dx = -\operatorname{coth} x + c$$

$$\int \frac{1}{\sqrt{1+x^2}} dx = \ln\left|x + \sqrt{x^2 + 1}\right| + c = \sinh^{-1} x + c$$

$$\int \frac{1}{\sqrt{x^2-1}} dx = \ln\left|x + \sqrt{x^2 - 1}\right| + c = \cosh^{-1} x + c$$

$$\int \frac{1}{1-x^2} dx = \frac{1}{2} \ln\left|\frac{1+x}{1-x}\right| + c = \tanh^{-1} x + c$$

$$\int \frac{1}{x^2-1} dx = -\frac{1}{2} \ln\left|\frac{x+1}{x-1}\right| + c = -\operatorname{coth}^{-1} x + c$$

$$\int \frac{1}{x\sqrt{1-x^2}} dx = -\ln\left|\frac{1+\sqrt{1-x^2}}{x}\right| + c = -\operatorname{sech}^{-1} x + c$$

$$\int \frac{1}{x\sqrt{1+x^2}} dx = -\ln\left|\frac{1+\sqrt{1+x^2}}{x}\right| + c = -\operatorname{cosech}^{-1} x + c$$

$$\cosh z = 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \cdots + \frac{z^{2n}}{(2n)!} + \cdots$$

$$\begin{aligned}\cosh z &= \frac{1}{2}(e^z + e^{-z}) \\ &= \frac{1}{2} \left(1 + z + \frac{z^2}{2!} + \cdots + \frac{z^n}{n!} + \cdots \right) + \frac{1}{2} \left(1 - z + \frac{z^2}{2!} - \cdots + (-1)^n \cdot \frac{z^n}{n!} + \cdots \right) \\ &= 1 + \frac{z^2}{2!} + \frac{z^4}{4!} + \cdots + \frac{z^{2n}}{(2n)!} + \cdots\end{aligned}$$

$$\sinh z = z + \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots + \frac{z^{2n-1}}{(2n-1)!} + \cdots$$

$$\begin{aligned}\sinh z &= \frac{1}{2}(e^z - e^{-z}) \\ &= \frac{1}{2} \left(1 + z + \frac{z^2}{2!} + \cdots + \frac{z^n}{n!} + \cdots \right) - \frac{1}{2} \left(1 - z + \frac{z^2}{2!} - \cdots + (-1)^n \cdot \frac{z^n}{n!} + \cdots \right) \\ &= z + \frac{z^3}{3!} + \frac{z^5}{5!} + \cdots + \frac{z^{2n-1}}{(2n-1)!} + \cdots\end{aligned}$$

$\cosh z = \cos iz$	$\sinh z = -i \sin iz$
$\cos iz = \frac{1}{2}(e^{i \cdot iz} + e^{-i \cdot iz})$	$-i \sin iz = -i \cdot \frac{1}{2i}(e^{i \cdot iz} - e^{-i \cdot iz})$
$= \frac{1}{2}(e^{-z} + e^z)$	$= -\frac{1}{2}(e^{-z} - e^z)$
$= \cosh z$	$= \sinh z$
$\tanh z = -i \tan iz$	$\coth z = i \cot iz$
$-i \tan iz = -i \cdot \frac{\frac{1}{2}(e^{i \cdot iz} - e^{-i \cdot iz})}{\frac{1}{2}(e^{i \cdot iz} + e^{-i \cdot iz})}$	$i \cot iz = i \cdot \frac{\frac{1}{2}(e^{i \cdot iz} + e^{-i \cdot iz})}{\frac{1}{2}(e^{i \cdot iz} - e^{-i \cdot iz})}$
$= -\frac{(e^{-z} - e^z)}{(e^{-z} + e^z)}$	$= -\frac{(e^{-z} + e^z)}{(e^{-z} - e^z)}$
$= \tanh z$	$= \coth z$
$\operatorname{sech} z = \sec iz$	$\operatorname{cosech} z = i \operatorname{cosec} iz$
$\sec iz = \frac{1}{\cos iz}$	$i \operatorname{cosec} iz = i \cdot \frac{1}{\sin iz}$
$= \frac{2}{e^{i \cdot iz} + e^{-i \cdot iz}}$	$= i \cdot \frac{1}{\frac{1}{2i}(e^{i \cdot iz} - e^{-i \cdot iz})}$
$= \frac{2}{e^{-z} + e^z}$	$= -\frac{2}{(e^{-z} - e^z)} = \operatorname{cosech} z$
$= \operatorname{sech} z$	
$\cosh^2 z - \sinh^2 z = 1$	$\operatorname{sech}^2 z = 1 - \tanh^2 z$
$\text{L.H.S.} = \frac{1}{4}(e^z + e^{-z})^2 - \frac{1}{4}(e^z - e^{-z})^2$	$1 - \tanh^2 z = \left(1 + \frac{e^z - e^{-z}}{e^z + e^{-z}}\right) \left(1 - \frac{e^z - e^{-z}}{e^z + e^{-z}}\right)$
$= \frac{1}{4}[(e^z + e^{-z})^2 - (e^z - e^{-z})^2]$	$= \frac{2e^z}{e^z + e^{-z}} \cdot \frac{2e^{-z}}{e^z + e^{-z}} = \left(\frac{2}{e^z + e^{-z}}\right)^2$
$= 1 = \text{R.H.S.}$	$= \operatorname{sech}^2 z$

$\text{cosech}^2 z = -1 + \coth^2 z$ $-1 + \coth^2 z = \left(\frac{e^z + e^{-z}}{e^z - e^{-z}} + 1 \right) \left(\frac{e^z + e^{-z}}{e^z - e^{-z}} - 1 \right)$ $= \frac{2e^z}{e^z - e^{-z}} \cdot \frac{2e^{-z}}{e^z - e^{-z}} = \left(\frac{2}{e^z - e^{-z}} \right)^2$ $= \text{cosech}^2 z$	$\sinh(-z) = \frac{1}{2}(e^{-z} - e^z) = -\frac{1}{2}(e^z - e^{-z}) = -\sinh z$ $\cosh(-z) = \frac{1}{2}(e^{-z} + e^z) = \cosh z$ $\sinh 0 = \frac{1}{2}(e^0 - e^{-0}) = 0; \cosh 0 = \frac{1}{2}(e^0 + e^{-0}) = 1$
--	--

$$\sinh(x \pm y) = \sinh x \cosh y \pm \cosh x \sinh y$$

$$\begin{aligned} & \sinh x \cosh y + \cosh x \sinh y \\ &= \frac{1}{2}(e^x - e^{-x}) \cdot \frac{1}{2}(e^y + e^{-y}) + \frac{1}{2}(e^x + e^{-x}) \cdot \frac{1}{2}(e^y - e^{-y}) \\ &= \frac{1}{4}(e^x - e^{-x})(e^y + e^{-y}) + \frac{1}{4}(e^x + e^{-x})(e^y - e^{-y}) \\ &= \frac{1}{4}(e^{x+y} - e^{-x+y} + e^{x-y} - e^{-x-y}) + \frac{1}{4}(e^{x+y} + e^{-x+y} - e^{x-y} - e^{-x-y}) \\ &= \frac{1}{2}(e^{x+y} - e^{-x-y}) \\ &= \sinh(x+y) \end{aligned}$$

The proof of $\sinh(x-y) = \sinh x \cosh y - \cosh x \sinh y$ is similar and omitted.

$$\cosh(x \pm y) = \cosh x \cosh y \pm \sinh x \sinh y$$

$$\begin{aligned} & \cosh x \cosh y + \sinh x \sinh y \\ &= \frac{1}{2}(e^x + e^{-x}) \cdot \frac{1}{2}(e^y + e^{-y}) + \frac{1}{2}(e^x - e^{-x}) \cdot \frac{1}{2}(e^y - e^{-y}) \\ &= \frac{1}{4}(e^x + e^{-x})(e^y + e^{-y}) + \frac{1}{4}(e^x - e^{-x})(e^y - e^{-y}) \\ &= \frac{1}{4}(e^{x+y} + e^{-x+y} + e^{x-y} + e^{-x-y}) + \frac{1}{4}(e^{x+y} - e^{-x+y} - e^{x-y} + e^{-x-y}) \\ &= \frac{1}{2}(e^{x+y} + e^{-x-y}) \\ &= \cosh(x+y) \end{aligned}$$

The proof of $\cosh(x-y) = \cosh x \cosh y - \sinh x \sinh y$ is similar and omitted.

$\tanh(x \pm y) = \frac{\tanh x \pm \tanh y}{1 \pm \tanh x \tanh y}$ $\begin{aligned} \tanh(x+y) &= \frac{\sinh(x+y)}{\cosh(x+y)} \\ &= \frac{\sinh x \cosh y + \cosh x \sinh y}{\cosh x \cosh y + \sinh x \sinh y} \\ &= \frac{\sinh x \cosh y}{\cosh x \cosh y} + \frac{\cosh x \sinh y}{\cosh x \cosh y} \\ &= \frac{\sinh x \sinh y}{\cosh x \cosh y} + \frac{\cosh x \cosh y}{\cosh x \cosh y} \\ &= \frac{\tanh x + \tanh y}{1 + \tanh x \tanh y} \end{aligned}$ $\text{The proof of } \tanh(x-y) = \frac{\tanh x - \tanh y}{1 - \tanh x \tanh y}$ <p>is similar and omitted.</p>	$\begin{aligned} \sinh 2z &= 2 \sinh z \cosh z \\ \sinh 2z &= \sinh(z+z) \\ &= \sinh z \cosh z + \cosh z \sinh z \\ &= 2 \sinh z \cosh z \\ \cosh 2z &= \cosh^2 z + \sinh^2 z = 2 \cosh^2 z - 1 = 2 \sinh^2 z + 1 \\ \cosh 2z &= \cosh(z+z) \\ &= \cosh z \cosh z + \sinh z \sinh z \\ &= \cosh^2 z + \sinh^2 z \dots\dots (*) \\ &= \cosh^2 z + \cosh^2 z - 1, \text{ using } \cosh^2 z - \sinh^2 z = 1 \\ &= 2 \cosh^2 z - 1 \dots\dots (**) \\ &= 1 + \sinh^2 z + \sinh^2 z, \text{ sub. the identity into } (*) \\ &= 2 \sinh^2 z + 1 \dots\dots (***) \end{aligned}$
---	---

$$\sin(x + iy) = \sin x \cosh y + i \cos x \sinh y$$

$$\sin(x + iy) = \frac{1}{2i} (e^{i(x+iy)} - e^{-i(x+iy)}) = \frac{1}{2i} (e^{-y+ix} - e^{y-ix})$$

$$\sin x \cosh y + i \cos x \sinh y$$

$$= \frac{1}{2i} (e^{ix} - e^{-ix}) \cdot \frac{1}{2} (e^y + e^{-y}) + i \cdot \frac{1}{2} (e^x + e^{-x}) \cdot \frac{1}{2} (e^y - e^{-y})$$

$$= \frac{1}{4i} (e^{y+ix} - e^{y-ix} + e^{-y+ix} - e^{-y-ix}) - \frac{1}{4i} (e^{y+ix} + e^{y-ix} - e^{-y+ix} - e^{-y-ix})$$

$$= \frac{2}{4i} (-e^{y-ix} + e^{-y+ix})$$

$$= \frac{1}{2i} (e^{-y+ix} - e^{y-ix}) = \sin(x + iy)$$

$$\cos(x + iy) = \cos x \cosh y - i \cos x \sinh y$$

$$\cos(x + iy) = \frac{1}{2} (e^{i(x+iy)} + e^{-i(x+iy)}) = \frac{1}{2} (e^{-y+ix} + e^{y-ix})$$

$$\cos x \cosh y - i \sin x \sinh y$$

$$= \frac{1}{2} (e^{ix} + e^{-ix}) \cdot \frac{1}{2} (e^y + e^{-y}) - i \cdot \frac{1}{2i} (e^{ix} - e^{-ix}) \cdot \frac{1}{2} (e^y - e^{-y})$$

$$= \frac{1}{4} (e^{y+ix} + e^{y-ix} + e^{-y+ix} + e^{-y-ix}) - \frac{1}{4} (e^{y+ix} - e^{y-ix} - e^{-y+ix} + e^{-y-ix})$$

$$= \frac{2}{4} (e^{y-ix} + e^{-y+ix})$$

$$= \frac{1}{2} (e^{y-ix} + e^{-y+ix}) = \cos(x + iy)$$

$$\sinh(x + iy) = \sinh x \cos y + i \cosh x \sin y$$

$$\sinh(x + iy) = \sinh x \cosh iy + \cosh x \sinh iy$$

$$= \sinh x \cos y + \cosh x \cdot \frac{1}{2} (e^{iy} - e^{-iy})$$

$$= \sinh x \cos y + i \cosh x \cdot \frac{1}{2i} (e^{iy} - e^{-iy})$$

$$= \sinh x \cos y + i \cosh x \sin y$$

$$\tanh^{-1} z = \frac{1}{2} \log \left(\frac{1+z}{1-z} \right) \text{ for } -1 < z < 1$$

Let $y = \tanh^{-1} z$, then $z = \tanh y$

$$z = \frac{e^y - e^{-y}}{e^y + e^{-y}} = \frac{e^{2y} - 1}{e^{2y} + 1}$$

$$(e^{2y} + 1)z = e^{2y} - 1$$

$$e^{2y} \cdot (1 - z) = 1 + z$$

$$e^{2y} = \frac{1+z}{1-z}$$

$$2y = \log \left(\frac{1+z}{1-z} \right)$$

$$y = \tanh^{-1} z = \frac{1}{2} \log \left(\frac{1+z}{1-z} \right) \text{ for } -1 < z < 1$$

$$\cosh(x + iy) = \cosh x \cos y + i \sinh x \sin y$$

$$\cosh(x + iy) = \cosh x \cosh iy + \sinh x \sinh iy$$

$$= \cosh x \cos y + \sinh x \cdot \frac{1}{2} (e^{iy} - e^{-iy})$$

$$= \sinh x \cos y + i \sinh x \cdot \frac{1}{2i} (e^{iy} - e^{-iy})$$

$$= \sinh x \cos y + i \sinh x \sin y$$

$$\sinh^{-1} x = \ln \left(x + \sqrt{x^2 + 1} \right)$$

Let $y = \sinh^{-1} x$, then $\sinh y = x$

$$\frac{1}{2} (e^y - e^{-y}) = x$$

$$e^{2y} - 2x e^y - 1 = 0$$

$$e^y = x + \sqrt{x^2 + 1} \text{ or } x - \sqrt{x^2 + 1}$$

$\therefore e^y > 0$ and $x - \sqrt{x^2 + 1} < x - x = 0$

$$\therefore e^y = x + \sqrt{x^2 + 1} \text{ only}$$

$$y = \sinh^{-1} x = \ln \left(x + \sqrt{x^2 + 1} \right)$$

$\cosh^{-1} x = \ln(x + \sqrt{x^2 - 1})$, for $x \geq 1$ Let $y = \cosh^{-1} x$, then $x = \cosh y$ $x = \frac{1}{2}(e^y + e^{-y})$, y is well defined only when $x > 0$ $e^{2y} - 2x e^y + 1 = 0$ $e^y = x + \sqrt{x^2 - 1}$ or $x - \sqrt{x^2 - 1}$ $\therefore (x + \sqrt{x^2 - 1})(x - \sqrt{x^2 - 1}) = 1$ $\therefore x - \sqrt{x^2 - 1} = \frac{1}{x + \sqrt{x^2 - 1}} = (x + \sqrt{x^2 - 1})^{-1}$ $e^y = x + \sqrt{x^2 - 1}$ or $(x + \sqrt{x^2 - 1})^{-1}$ $y = \ln(x + \sqrt{x^2 - 1})$ or $\ln(x + \sqrt{x^2 - 1})^{-1}$ $y = \ln(x + \sqrt{x^2 - 1})$ or $-\ln(x + \sqrt{x^2 - 1})$ y is well defined only when $x \geq 1$ or $x \leq -1$ It is convenient to adopt the positive sign and to write $y = \cosh^{-1} x = \ln(x + \sqrt{x^2 - 1})$	$\tanh^{-1} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$, $-1 < x < 1$ Let $y = \tanh^{-1} x$, then $x = \tanh y$ $x = \frac{e^z - e^{-z}}{e^z + e^{-z}} = \frac{e^{2z} - 1}{e^{2z} + 1}$ $x e^{2z} + x = e^{2z} - 1$ $x + 1 = e^{2z}(1 - x)$ $e^{2z} = \frac{1+x}{1-x} > 0 \Rightarrow -1 < x < 1$ $2z = \ln\left(\frac{1+x}{1-x}\right)$ $z = \tanh^{-1} x = \frac{1}{2} \ln\left(\frac{1+x}{1-x}\right)$, $-1 < x < 1$
$\operatorname{cosech}^{-1} x = \ln\left(\frac{1 \pm \sqrt{1+x^2}}{x}\right)$, $x \neq 0$ Let $y = \operatorname{cosech}^{-1} x$, then $\operatorname{cosech} y = x$ $x = \frac{2}{e^y - e^{-y}} = \frac{2e^y}{e^{2y} - 1}$ $x e^{2y} - 2 e^y - x = 0$ $e^y = \frac{1 \pm \sqrt{1+x^2}}{x} > 0$ When $x > 0$, $1+x^2 > 1 \Rightarrow \frac{1-\sqrt{1+x^2}}{x} < 0$ (rejected) When $x < 0$, $\frac{1+\sqrt{1+x^2}}{x} < 0$ (rejected) $y = \operatorname{cosech}^{-1} x = \begin{cases} \ln\left(\frac{1+\sqrt{1+x^2}}{x}\right) & \text{if } x > 0 \\ \ln\left(\frac{1-\sqrt{1+x^2}}{x}\right) & \text{if } x < 0 \end{cases}$	$\operatorname{sech}^{-1} x = \ln\left(\frac{1+\sqrt{1-x^2}}{x}\right)$, $0 < x \leq 1$ Let $y = \operatorname{sech}^{-1} x$, then $\operatorname{sech} y = x$ $x = \frac{2}{e^y + e^{-y}} = \frac{2e^y}{e^{2y} + 1}$ $x e^{2y} - 2 e^y + x = 0$ $e^y = \frac{1 \pm \sqrt{1-x^2}}{x} > 0 \Rightarrow -1 \leq x \leq 1, x \neq 0$ $\frac{x}{1+\sqrt{1-x^2}} \cdot \frac{1-\sqrt{1-x^2}}{x} = 1$ $\frac{1-\sqrt{1-x^2}}{x} = \left(\frac{1+\sqrt{1-x^2}}{x}\right)^{-1}$ When $x < 0$, $\frac{1 \pm \sqrt{1-x^2}}{x} < 0$ (rejected) $y = \ln\left(\frac{1+\sqrt{1-x^2}}{x}\right)$ or $-\ln\left(\frac{1+\sqrt{1-x^2}}{x}\right)$ It is convenient to adopt the positive sign and to write $y = \operatorname{sech}^{-1} x = \ln\left(\frac{1+\sqrt{1-x^2}}{x}\right)$, $0 < x \leq 1$

$$\coth^{-1} x = \frac{1}{2} \ln\left(\frac{x+1}{x-1}\right), |x| > 1; \text{ let } y = \coth^{-1} x \Rightarrow x = \coth y = \frac{e^y + e^{-y}}{e^y - e^{-y}} = \frac{e^{2y} + 1}{e^{2y} - 1}$$

$$x e^{2y} - x = e^{2y} + 1 \Rightarrow e^{2y}(x-1) = x+1 \Rightarrow e^{2y} = \frac{x+1}{x-1} > 0, x < -1 \text{ or } x > 1$$

$$y = \frac{1}{2} \ln\left(\frac{x+1}{x-1}\right), |x| > 1$$

$\frac{d \sinh x}{dx} = \cosh x$ $\frac{d \sinh x}{dx} = \frac{d}{dx} \left[\frac{1}{2} (e^x + e^{-x}) \right]$ $= \frac{1}{2} (e^x - e^{-x})$ $= \cosh x$	$\frac{d \cosh x}{dx} = \sinh x$ $\frac{d \cosh x}{dx} = \frac{d}{dx} \left[\frac{1}{2} (e^x - e^{-x}) \right]$ $= \frac{1}{2} (e^x + e^{-x})$ $= \sinh x$
$\frac{d \tanh x}{dx} = \operatorname{sech}^2 x$ $\frac{d \tanh x}{dx} = \frac{d}{dx} \left(\frac{e^x - e^{-x}}{e^x + e^{-x}} \right)$ $= \frac{(e^x + e^{-x})^2 - (e^x - e^{-x})^2}{(e^x + e^{-x})^2}$ $= \frac{4}{(e^x + e^{-x})^2}$ $= \operatorname{sech}^2 x$	$\frac{d \operatorname{cosech} x}{dx} = -\operatorname{cosech} x \coth x$ $\frac{d \operatorname{cosech} x}{dx} = \frac{d}{dx} \left(\frac{2}{e^x - e^{-x}} \right)$ $= -\frac{2}{(e^x - e^{-x})^2} \cdot (e^x + e^{-x})$ $= -\frac{2}{e^x - e^{-x}} \cdot \frac{e^x + e^{-x}}{e^x - e^{-x}}$ $= -\operatorname{cosech} x \coth x$
$\frac{d \operatorname{sech} x}{dx} = -\operatorname{sech} x \tanh x$ $\frac{d \operatorname{sech} x}{dx} = \frac{d}{dx} \left(\frac{2}{e^x + e^{-x}} \right)$ $= -\frac{2}{(e^x + e^{-x})^2} \cdot (e^x - e^{-x})$ $= -\frac{2}{e^x + e^{-x}} \cdot \frac{e^x - e^{-x}}{e^x + e^{-x}}$ $= -\operatorname{sech} x \tanh x$	$\frac{d \coth x}{dx} = -\operatorname{cosech} x$ $\frac{d \coth x}{dx} = \frac{d}{dx} \left(\frac{e^x + e^{-x}}{e^x - e^{-x}} \right)$ $= \frac{(e^x - e^{-x})^2 - (e^x + e^{-x})^2}{(e^x - e^{-x})^2}$ $= -\frac{4}{(e^x - e^{-x})^2}$ $= -\operatorname{cosech}^2 x$
$\frac{d \sinh^{-1} x}{dx} = \frac{1}{\sqrt{1+x^2}}$ <p>Let $y = \sinh^{-1} x$, then $\sinh y = x$</p> $\therefore \cosh^2 y - \sinh^2 y = 1$ $\therefore \cosh y = \pm \sqrt{1 + \sinh^2 y} = \pm \sqrt{1 + x^2}$ $\cosh y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{1}{\cosh y} = \pm \frac{1}{\sqrt{1-x^2}}$ <p>The slope of the graph of $y = \sinh x > 0 \forall x$</p> $\therefore \frac{d \sinh^{-1} x}{dx} = \frac{1}{\sqrt{1+x^2}}$	$\frac{d \cosh^{-1} x}{dx} = \frac{1}{\sqrt{x^2-1}} \text{ for all } x > 1$ <p>Let $y = \cosh^{-1} x$, then $\cosh y = x$</p> $\therefore \cosh^2 y - \sinh^2 y = 1$ $\therefore \sinh y = \pm \sqrt{\cosh^2 y - 1} = \pm \sqrt{x^2 - 1}$ $\sinh y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{1}{\sinh y} = \pm \frac{1}{\sqrt{x^2 - 1}}$ <p>The slope of the graph of $y = \cosh x > 0 \forall x > 1$</p> $\frac{d}{dx} \cosh^{-1} x = \frac{1}{\sinh y} = \frac{1}{\sqrt{x^2 - 1}} \text{ for } x > 1$

$\frac{d \tanh^{-1} x}{dx} = \frac{1}{1-x^2} \text{ for } x < 1$ <p>Let $y = \tanh^{-1} x$, then $\tanh y = x$</p> $\operatorname{sech}^2 y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{1}{\operatorname{sech}^2 y}$ $= \frac{1}{1 - \tanh^2 y}$ $= \frac{1}{1 - x^2}$ <p>The slope of the graph of $y = \tanh x > 0 \forall x$</p> $\therefore \frac{d \tanh^{-1} x}{dx} = \frac{1}{1-x^2} \text{ for } x < 1$	$\frac{d \coth^{-1} x}{dx} = \frac{1}{1-x^2}, x > 1$ <p>Let $y = \coth^{-1} x$, then $\coth y = x$</p> $-\operatorname{cosech}^2 y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{-1}{\operatorname{cosech}^2 y}$ $= \frac{-1}{\coth^2 y - 1}$ $= \frac{1}{1-x^2}$ <p>The slope of the graph of $y = \coth x < 0 \forall x$</p> $\therefore \frac{d \coth^{-1} x}{dx} = \frac{1}{1-x^2}, x > 1$
$\frac{d \operatorname{sech}^{-1} x}{dx} = -\frac{1}{x\sqrt{1-x^2}}, 0 < x < 1$ <p>Let $y = \operatorname{sech}^{-1} x$, then $\operatorname{sech} y = x$</p> $-\operatorname{sech} y \tanh y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{-1}{\operatorname{sech} y \tanh y}$ $= \frac{-1}{\pm x\sqrt{1-\operatorname{sech}^2 y}}$ $= \frac{-1}{\pm x\sqrt{1-x^2}}$ <p>The slope of the graph of $y = \cosh x > 0, 0 < x < 1$</p> <p>The slope of the graph of $y = \operatorname{sech} x < 0, 0 < x < 1$</p> $\therefore \frac{d \operatorname{sech}^{-1} x}{dx} = -\frac{1}{x\sqrt{1-x^2}}, 0 < x < 1$	$\frac{d \operatorname{cosech}^{-1} x}{dx} = \frac{-1}{ x \sqrt{1+x^2}} \text{ for } x \neq 0$ <p>Let $y = \operatorname{cosech}^{-1} x$, then $\operatorname{cosech} y = x$</p> $-\operatorname{cosech} y \coth y \frac{dy}{dx} = 1$ $\frac{dy}{dx} = \frac{-1}{\operatorname{cosech} y \coth y}$ $= \frac{-1}{\pm x\sqrt{1+\operatorname{cosech}^2 y}}$ $= \frac{-1}{\pm x\sqrt{1+x^2}}$ <p>The slope of the graph of $y = \sinh x > 0 \forall x$</p> <p>The slope of the graph of $y = \operatorname{cosech} x < 0, x \neq 0$</p> $\therefore \frac{d \operatorname{cosech}^{-1} x}{dx} = \frac{-1}{ x \sqrt{1+x^2}}, x \neq 0$
$\int \frac{1}{\sqrt{1+x^2}} dx = \ln x + \sqrt{x^2 + 1} + c = \sinh^{-1} x + c$ $\therefore \frac{d \sinh^{-1} x}{dx} = \frac{1}{\sqrt{1+x^2}}$ $\therefore \int \frac{1}{\sqrt{1+x^2}} dx = \sinh^{-1} x + c$ $= \ln x + \sqrt{x^2 + 1} + c$	$\int \frac{1}{\sqrt{x^2 - 1}} dx = \ln x + \sqrt{x^2 - 1} + c = \cosh^{-1} x + c$ $\therefore \frac{d \cosh^{-1} x}{dx} = \frac{1}{\sqrt{x^2 - 1}}, \text{ for all } x > 1$ $\therefore \int \frac{1}{\sqrt{x^2 - 1}} dx = \cosh^{-1} x + c$ $= \ln(x + \sqrt{x^2 - 1}) + c$
$\int \frac{1}{1-x^2} dx = \frac{1}{2} \ln \left \frac{1+x}{1-x} \right + c = \tanh^{-1} x + c$ $\therefore \frac{d \tanh^{-1} x}{dx} = \frac{1}{1-x^2}, \text{ for } x < 1$ $\therefore \int \frac{1}{1-x^2} dx = \tanh^{-1} x + c = \frac{1}{2} \ln \left(\frac{1+x}{1-x} \right) + c$	$\int \frac{1}{x^2-1} dx = -\frac{1}{2} \ln \left \frac{x+1}{x-1} \right + c = -\operatorname{coth}^{-1} x + c$ $\therefore \frac{d \coth^{-1} x}{dx} = \frac{1}{1-x^2}, x > 1$ $\int \frac{1}{x^2-1} dx = -\operatorname{coth}^{-1} x + c = -\frac{1}{2} \ln \left \frac{x+1}{x-1} \right + c$

$\int \frac{1}{x\sqrt{1-x^2}} dx = -\ln \left \frac{1+\sqrt{1-x^2}}{x} \right + c = -\operatorname{sech}^{-1} x + c$ $\therefore \frac{d \operatorname{sech}^{-1} x}{dx} = -\frac{1}{x\sqrt{1-x^2}}, \quad 0 < x < 1$ $\int \frac{1}{x\sqrt{1-x^2}} dx = -\operatorname{sech}^{-1} x + c$ $= -\ln \left \frac{1+\sqrt{1-x^2}}{x} \right + c$	$\int \frac{1}{x\sqrt{1+x^2}} dx = -\ln \left \frac{1+\sqrt{1+x^2}}{x} \right + c = -\operatorname{cosech}^{-1} x + c$ $\therefore \frac{d \operatorname{cosech}^{-1} x}{dx} = \frac{-1}{ x \sqrt{1+x^2}}, \text{ for } x \neq 0$ $\int \frac{1}{x\sqrt{1+x^2}} dx = -\operatorname{cosech}^{-1} x + c$ $= -\ln \left \frac{1+\sqrt{1+x^2}}{x} \right + c$
--	--