

Rotation of axis example

Created by Mr. Francis Hung on 20111116

Last updated: August 30, 2021

Given the central conics: $Ax^2 + Bxy + Cy^2 = D$, $A \neq C$, $B \neq 0$, $D \neq 0$

Perform the rotation of axes: $\begin{cases} x = x_1 \cos \theta - y_1 \sin \theta \\ y = x_1 \sin \theta + y_1 \cos \theta \end{cases}$, where $\theta = \frac{1}{2} \tan^{-1} \frac{B}{A-C}$.

Prove that after the transformation, the equation is in the form $ax_1^2 + by_1^2 = D$.

Proof:

$$\tan 2\theta = \frac{B}{A-C}$$

$$A(x_1 \cos \theta - y_1 \sin \theta)^2 + B(x_1 \cos \theta - y_1 \sin \theta)(x_1 \sin \theta + y_1 \cos \theta) + C(x_1 \sin \theta + y_1 \cos \theta)^2 = D$$

$$(A \cos^2 \theta + B \sin \theta \cos \theta + C \sin^2 \theta)x_1^2 + [-2A \sin \theta \cos \theta + B(\cos^2 \theta - \sin^2 \theta) + 2C \sin \theta \cos \theta]x_1y_1 + (A \sin^2 \theta - B \sin \theta \cos \theta + C \cos^2 \theta)y_1^2 = D$$

$$\left(A \frac{1+\cos 2\theta}{2} + \frac{B \sin 2\theta}{2} + C \frac{1-\cos 2\theta}{2} \right)x_1^2 + [(C-A)\sin 2\theta + B \cos 2\theta]x_1y_1 + \left(A \frac{1-\cos 2\theta}{2} - \frac{B \sin 2\theta}{2} + C \frac{1+\cos 2\theta}{2} \right)y_1^2 = D$$

$$[A+C+B \sin 2\theta + (A-C)\cos 2\theta]x_1^2 + 2[(C-A)\sin 2\theta + B \cos 2\theta]x_1y_1 + [A+C-B \sin 2\theta + (C-A)\cos 2\theta]y_1^2 = 2D$$

$$\left[A+C + \frac{B^2}{\sqrt{(A-C)^2 + B^2}} + \frac{(A-C)^2}{\sqrt{(A-C)^2 + B^2}} \right]x_1^2 + 2 \left(\frac{B(C-A)}{\sqrt{(A-C)^2 + B^2}} + \frac{B(A-C)}{\sqrt{(A-C)^2 + B^2}} \right)x_1y_1 + \left[A+C - \frac{B^2}{\sqrt{(A-C)^2 + B^2}} - \frac{(A-C)^2}{\sqrt{(A-C)^2 + B^2}} \right]y_1^2 = 2D$$

$$\left[A+C + \sqrt{(A-C)^2 + B^2} \right]x_1^2 + \left[A+C - \sqrt{(A-C)^2 + B^2} \right]y_1^2 = 2D$$

$$\left[\frac{A+C + \sqrt{(A-C)^2 + B^2}}{2} \right]x_1^2 + \left[\frac{A+C - \sqrt{(A-C)^2 + B^2}}{2} \right]y_1^2 = D$$

It is in the form $ax_1^2 + by_1^2 = D$.